



General Certificate of Education

Mathematics 6360

MS2A Statistics 2A

Mark Scheme

2007 examination - June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2007 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme and abbreviations used in marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
✓ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	C	candidate
PI	possibly implied	Sf	significant figure(s)
SCA	substantially correct approach	Dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.


MS2A

Q	Solution	Marks	Total	Comments																				
1	H ₀ : condition independent of treatment H ₁ : condition dependent upon treatment	B1																						
	Totals: 66, 84, 75, 75	B1																						
	<table border="1"> <thead> <tr> <th>O</th> <th>E</th> <th> O-E -0.5</th> <th></th> </tr> </thead> <tbody> <tr> <td>20</td> <td>33</td> <td>12.5</td> <td>4.7348</td> </tr> <tr> <td>55</td> <td>42</td> <td></td> <td>3.7202</td> </tr> <tr> <td>46</td> <td>33</td> <td></td> <td>4.7348</td> </tr> <tr> <td>29</td> <td>42</td> <td></td> <td>3.7202</td> </tr> </tbody> </table>	O	E	O-E -0.5		20	33	12.5	4.7348	55	42		3.7202	46	33		4.7348	29	42		3.7202	M1A1 M1 M1		for E _i attempted, correctly for use of Yates' correction final column If no Yates' correction: possible M1A1M0M1A0 If 0.5 incorrectly used: possible M1A1M1M1A0
	O	E	O-E -0.5																					
	20	33	12.5	4.7348																				
	55	42		3.7202																				
	46	33		4.7348																				
	29	42		3.7202																				
	X ² = 16.9	A1																						
	χ _{5%} ² (1) = 3.841 < 16.91	B1✓																						
Reject H ₀	A1✓																							
Evidence to suggest that the condition of the patients may be dependent upon the treatment that they received	E1✓	10																						
Total		10																						
2(a)(i)	$P(X = 3) = \frac{e^{-3.5} \times (3.5)^3}{3!} = 0.216$	M1 A1	2																					
(ii)	$P(Y \geq 5) = 1 - P(Y \leq 4)$ $= 1 - 0.2851$ $= 0.715$	M1 A1	2																					
(b)(i)	T ~ Po(9.5)	B1	1																					
(ii)	$P(7 \leq T \leq 10) = P(T \leq 10) - P(T \leq 6)$ $= 0.6453 - 0.1649$ $= 0.480$	M1 A1 A1	3	$\sum_{x=7}^{x=10} P(X = x) = 0.48$ Accept 0.48																				
(iii)	$p = (0.4804)^3 = 0.111$	M1 A1✓	2	on their (b)(ii)																				
Total			10																					

MS2A (cont)

Q	Solution	Marks	Total	Comments
3	$H_0 : \mu = 36$ $H_1 : \mu < 36$ $\bar{x} = \frac{1730}{50} = 34.6$ $s^2 = \frac{784}{49} = 16$ Test statistic: $z = \frac{34.6 - 36}{\frac{4}{\sqrt{50}}} = -2.47$ $z_{\text{crit}} = -2.3263$ Reject H_0 Sufficient evidence at the 1% level of significance to support David's claim	B1 B1 B1 M1 A1 B1 A1✓ E1✓	8	(-2.48 to -2.47) (dep M1)
	Total		8	
4(a)	$\bar{x} = 9.70$ $s = 0.0294$ ($s^2 = 8.67 \times 10^{-4}$) 95% Confidence interval for g : $= 9.70 \pm 2.262 \times \frac{0.0294}{\sqrt{10}}$ $= 9.70 \pm 0.021$ $= (9.68, 9.72)$	B1 B1 B1 M1 A1✓ A1	6	$\sigma = 0.0279$ $(\sigma^2 = 7.8 \times 10^{-4})$ or $\frac{0.0279}{3}$ on their \bar{x} and s
(b)	(9.78, 9.82)	B2✓	2	on their CI in (a) accept repeat of (a) using $(\bar{x} + 0.5)$ and s
	Total		8	

MS2A (cont)

Q	Solution	Mark	Total	Comments
5(a)	$1-3p \geq 0$ and $p \geq 0$	M1	3	
	$3p \leq 1$	A1		
	$0 \leq p \leq \frac{1}{3}$	A1		
(b)(i)	$E(X) = (1 \times p) + (2 \times p) + (3 \times p) + 4(1-3p)$ $= 4 - 6p$	B1	1	
(ii)	$E(X^2) = (1 \times p) + (4 \times p) + (9 \times p) + 16(1-3p)$ $= 16 - 34p$	M1	3	Attempting $E(X^2)$
	$\text{Var}(X) = (16 - 34p) - (4 - 6p)^2$	M1		
	$= 16 - 34p - 16 + 48p - 36p^2$ $= 14p - 36p^2$ $= 2p(7 - 18p)$	A1		
(c)(i)	Max value of quadratic graph occurs at:			
	 <p>$p = \frac{1}{2} \left(0 + \frac{7}{18} \right) = \frac{7}{36}$</p>	M1	2	Or $\frac{dV}{dp} = 14 - 72p$ and $\frac{d^2V}{dp^2} = -72$ For max V , $\frac{dV}{dp} = 0$ and $\frac{d^2V}{dp^2} < 0$ $\Rightarrow p = \frac{7}{36}$
(ii)	Max value of $\text{Var}(X)$			
	$= 2 \times \frac{7}{36} \times \left(7 - 18 \times \frac{7}{36} \right)$ $= 2 \times \frac{7}{36} \times \frac{7}{2}$ $= \frac{49}{36}$	M1		Attempted
	$\text{sd}_{\max}(X) = \frac{7}{6}$ (1.17)	M1 A1	3	(square root attempted) CAO
Total			12	

MS2A(cont)

Q	Solution	Marks	Total	Comments
6(a)(i)	$E\left(\frac{1}{X}\right) = \int_0^1 \frac{1}{x} 3x^2 dx = \int_0^1 3x dx$	M1	3	CAO
	$= \left[\frac{3x^2}{2} \right]_0^1 = 1.5$	A1 A1		
(ii)	$E\left(\frac{1}{X^2}\right) = \int_0^1 \frac{1}{x^2} 3x^2 dx = \int_0^1 3 dx$	M1	4	on their (i) and Var > 0
	$= [3x]_0^1 = 3.0$	A1		
	$\text{Var}\left(\frac{1}{X}\right) = 3.0 - (1.5)^2$	m1		
	$= 0.75$	A1✓		
(b)	$E\left(\frac{5+2X}{X}\right) = E\left(\frac{5}{X} + 2\right)$	M1	5	CAO
	$= 5E\left(\frac{1}{X}\right) + 2$	M1		
	$= 5 \times 1.5 + 2$	A1		
	$= 9.5$			
	$\text{Var}\left(\frac{5+2X}{X}\right) = \text{Var}\left(\frac{5}{X} + 2\right)$			
	$= 25 \times \text{Var}\left(\frac{1}{X}\right)$	M1		
$= 25 \times 0.75$				
	$= 18.75$	A1		CAO
	Total		12	
	TOTAL		60	