

**BUSINESS MATHEMATICS**

(Three hours and a quarter)

-----  
*Answer **Question 1** from Section A and **10 Questions** from Section B.*  
*All working, including rough work, should be done on the same sheet as, and adjacent to, the rest of the answer.*

*The intended marks for the **Questions** or parts of **Questions** are given in brackets [ ]*

***Mathematical formulae are given at the end of this Question paper.***

***The use of calculator (fx-82/fx-100) is allowed without memory.***

*Diagrams given in this question booklet are not in scale.*

-----

**Section A(30 Marks)**Answer **ALL** questions

**Directions:** Read the following questions carefully. For each question there are four alternatives **A, B, C** and **D**. Choose the correct alternative and write it in your answer sheet.

**Question 1.****(2×15=30 Marks)**(i) Summation notation for  $1+7+17+31$  is


**A**  $\sum_{i=1}^4 2i^2 - 1$

**B**  $\sum_{i=1}^4 2i - 1$

**C**  $\sum_{i=1}^4 4i^2 - 3$

**D**  $\sum_{i=1}^4 i$

(ii) The Factors of the polynomial of the given graph are



- A  $(x(x+2), (x+3))$
- B  $x, (x+2), (x-3)$
- C  $x(x-2), (x-3)$
- D  $x, (x-2), (x+3)$

(iii) If  $\tan \theta = \frac{12}{5}$ ,  $0 < \theta < \frac{\pi}{2}$ , then the value of  $\cos 2\theta$  is

- A  $\frac{169}{119}$
- B  $-\frac{169}{119}$
- C  $\frac{119}{169}$
- D  $-\frac{119}{169}$

(iv) Which of the following is a rational number?

- A  $(5 - \sqrt{3})^2$
- B  $(\sqrt{2} + 1)^2$
- C  $(\sqrt{8} + \sqrt{2})^2$
- D  $(\sqrt{3} - \sqrt{2})^2$

(v) The restrictions of the expression  $\frac{x^2 - 4}{x^2 + x - 6} \div \frac{x^2 + 3x}{x^2 + 7x + 12}$  are

- A  $x \neq \{-3, 0, 2\}$
- B  $x \neq \{-3, 0\}$
- C  $x \neq \{-3, 2\}$
- D  $x \neq \{-2, 0, 3\}$

(vi) For  $y = \frac{x-3}{2-x}$  Evaluate  $\frac{dy}{dx}\Big|_{x=1}$

- A -1
- B 3
- C 5
- D -3

(vii) If  $y = u^2 + 4u$  and  $u = \sqrt{x}$  then  $\frac{dy}{dx}$  is equal to

- A  $2\sqrt{x} + 4$
- B  $1 - \frac{2}{\sqrt{x}}$
- C  $1 + \frac{2}{\sqrt{x}}$
- D  $4x + 8\sqrt{x}$

(viii) The point on the curve  $y = \sin x$  at which the tangent is horizontal is

- A  $\left(1, \frac{\pi}{2}\right)$
- B  $\left(-\frac{\pi}{2}, -1\right)$
- C  $\left(-\frac{\pi}{2}, 1\right)$
- D  $\left(\frac{\pi}{2}, 1\right)$

(ix) Given  $y = x^x$  then  $\frac{dy}{dx}$  is

- A  $(1 + \log_e x)$
- B  $x^x (1 + \log_e x)$
- C  $\frac{1}{x^x (1 + \log_e x)}$
- D  $\frac{x^x}{1 + \log_e x}$

(x)  $\int \frac{x^3 + x^2 + x + 1}{x} dx$  is equal to

- A**  $x^2 + x + 1 + \frac{1}{x} + c$   
**B**  $\log_e (x^3 + x^2 + x + 1) + c$   
**C**  $\frac{x^3}{3} + \frac{x^2}{2} + x + \log_e x + c$   
**D**  $2x + 1 + \frac{1}{x^2} + c$

(xi)  $\int \frac{\cos(\log_e x)}{x} dx$  is equal to

- A**  $-\sin(\log_e x) + c$   
**B**  $\frac{\cos^2(\log_e x)}{2} + c$   
**C**  $-\cos(\log_e x) + c$   
**D**  $\sin(\log_e x) + c$

(xii) If  $A = \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix}$  then  $A \cdot (\text{Adj}A)$  is

- A**  $\begin{pmatrix} 4 & -3 \\ -1 & 2 \end{pmatrix}$   
**B**  $\begin{pmatrix} 5 & 0 \\ 0 & 5 \end{pmatrix}$   
**C**  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$   
**D**  $\begin{pmatrix} 7 & -12 \\ 4 & -7 \end{pmatrix}$

(xiii) Find  $\frac{dy}{dx}$  when  $y = \log_5 x^3$

- A  $\frac{3}{x}$
- B  $\frac{3}{x} \log_{10} e$
- C  $\frac{3}{x} \log_e 5$
- D  $\frac{3}{x} \log_5 e$

(xiv) The mean marks in mathematics of class A, and class B are 50 and 60 respectively. The mean marks of the combined class is 54. If the strength of the class A is 60, then the strength of class B is

- A 90
- B 50
- C 40
- D 55

(xv) The vertical asymptote of  $y = \frac{x-2}{x^2+2x-8}$  is

- A  $x = 4$
- B  $x = -4$
- C  $x = 2, x = -4$
- D  $x = 2, x = 4$

**Section B (70 marks)**

Answer any 10 questions. All questions in this section have equal marks. Unless otherwise stated, you may round answers to decimal places.

**Question 2**

a) Simplify  $\frac{x-3}{x^2-9} - \frac{x+5}{x^2+8x+15}$  [3]

b) Using mathematical Induction , prove that

$$\frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + \dots + \frac{1}{n(n+1)} = \frac{n}{n+1}, \forall n \in N$$
 [4]

**Question 3**

- a) Determine a quadratic equation  $f(x)$  such that  $f(2) = 21$ ,  $f'(2) = 14$  and  $f''(2) = 6$  [3]
- b) A rectangular shopping complex is to be built on a 80m by 60m rectangular plot in such a way that there is a path  $x$  meter wide surrounding the building. The building can occupy up to 70% of the plot's area. What is the range of possible integral values for the width of the path ( $x$ ) [4]

**Question 4**

- (a) Convert the exponential function  $y = 20(x)^5$  into base  $-e$  form [2]
- (b) The following table shows the relationship between the ages of husbands ( $x$ ) and wives ( $y$ ) in a town. Using this, find the regression line of  $Y$  on  $X$ . hence estimate the age of wife when the age of husband is 30. [5]

Ages of husband ( $x$ )	25	22	28	26	35	20
Age of wife ( $y$ )	18	15	20	19	22	14

**Question 5**

- (a) Evaluate  $\sum_{i=1}^{20} i(2-3i)^2$  [4]
- (b) Evaluate  $\int x \sin x dx$  [3]

**Question 6**

- a) Find the square root of  $8 + 2\sqrt{15}$  [3]
- b) The population of a town in the year 2005 was 0.15 million. The population increases by 15% p.a.
- i) Estimate the population of the town in the year 2009
  - ii) When will the population of the town become double the population of 2005? (Give both the answer to the nearest integer) [4]

**Question 7**

(a) Using  $\sin(A + B) = \sin A \cos B + \cos A \sin B$ , Prove that  $\sin 3x = 3 \sin x - 4 \sin^3 x$  [4]

(b) Evaluate  $\int_1^e \frac{\log_e x}{x} dx$  [3]

**Question 8**

(a) For what values of  $x$  is the graph of  $y = x^3 + x$  below the graph of  $y = 6 - 4x^2$  [4]

(b) Solve the equations by matrix method

$$\begin{aligned} 8x + 3y - 2 &= 0 \\ 5x - 4y - 13 &= 0 \end{aligned} \quad [3]$$

**Question 9**

(a) Calculate the area bound by the  $y = x(x - 4)$  and the X – axis. This area is rotated about X-axis through four right angles to generate a solid. Calculate the volume of the solid generated. [5]

(b) Calculate the mean deviation about median of the following data  
65, 35, 50, 42, 55 [2]

**Question 10**

(a) Evaluate  $\int (\sqrt{x} - 2)^2 dx$  [2]

(b) Determine the dimensions of a rectangular playground with maximum area that can be enclosed with 676m of fencing [5]

**Question 11**

(a) Given  $X = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$ , does  $X^{-1}$  exist? Give a reason, Also find Adj X [4]

(b) At a certain instant, the area of a circular oil spill on the ocean increases at the rate of  $2cm^2 / sec$ . Determine the rate of increase of the radius when the area is  $50cm^2$  [3]

**Question 12**

(a) If  $y = \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}}$ , show that  $\frac{dy}{dx} = \sec^2 x$  [4]

(b) Given  $y = \frac{1-x}{1+x}$ , show that  $(1+x)^2 \frac{dy}{dx} + 2 = 0$  [3]

**Question 13**

(a) Find the equation of the tangent to the curve  $y = 5^x$  at  $x = 1$  [3]

(b) Describe the transformation that should be applied to the graph of  $y = e^x$  to obtain the graph of  $y = 5e^x - 4$  [2]

(c) The following table is taken from the family budget of a group of people from a village. Calculate the cost of living index for the year 2007 [2]

Expenses on	Weight	Price in 2005	Price in 2007
Rice	40	200	220
Oil	15	120	150
Dal	10	80	90
Kharang	35	150	200

**Question 14**

For the curve  $y = x^4 - 8x + 16$ , find

- The x and y intercepts
- The local maximum and minimum points
- Point of inflexion
- With the help of the above information, sketch the curve( Use the graph Paper) [7]



MATHEMATICS FORMULAE

**Functions and Equations**

- (1)  $(a \pm b)^2 = a^2 + b^2 \pm 2ab$
- (2)  $(a \pm b)^3 = a^3 \pm 3a^2b + 3ab^2 \pm b^3$
- (3)  $a^2 - b^2 = (a + b)(a - b)$
- (4)  $a^3 \pm b^3 = (a \pm b)(a^2 \mp ab + b^2)$
- (5)  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

**Sequence and series**

- (1)  $\sum_{i=1}^n i = \frac{n(n+1)}{2}$
- (2)  $\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$
- (3)  $\sum_{i=1}^n i^3 = \left[ \frac{n(n+1)}{2} \right]^2$
- (4)  $t_n = ar^{n-1}$
- (5)  $t_n = a + (n - 1)d$
- (6)  $S_n = \frac{a(1-r^n)}{1-r}$  where  $r < 1$   
 $= \frac{a(r^n - 1)}{r - 1}$ , Where  $r > 1$
- (7)  $S_n = \frac{n}{2} [2a + (n - 1)d]$

**Differentiation**

- (1)  $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
- (2)  $y = x^n, y' = nx^{n-1}$ ,
- (3)  $y = cf(x), y' = cf'(x)$
- (4)  $y = f(x) \pm g(x), y' = f'(x) \pm g'(x)$
- (5)  $F(x) = f(x)g(x),$   
 $F'(x) = f(x)g'(x) + f'(x)g(x)$
- (6)  $F(x) = \frac{f(x)}{g(x)},$

$$F'(x) = \frac{g(x)f'(x) - f(x)g'(x)}{[g(x)]^2}$$

- (7)  $f \circ g(x)' = f'g(x) \times (g'x)$
- (8)  $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$
- (9)  $v(t) = h'(t)$

**Coordinate Geometry**

- (1)  $(y - y_1) = m(x - x_1)$
- (2)  $\sqrt{(x - a)^2 + (y - b)^2}$

**Trigonometry**

- (1)  $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$
- (2)  $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$
- (3)  $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$
- (4)  $\sin^2 \theta + \cos^2 \theta = 1$

**Logarithmic Exponentials**

- (1)  $y = y_0(1 + r)^x$
- (2)  $y = y_0e^{kx}$
- (3)  $A = P(1 + r)^n$

**Integration**

- (1)  $\int f(x)g(x)dx = f(x)\int g(x)dx - \int \left[ \left( \frac{d}{dx} f(x) \right) \int g(x)dx \right] dx$
- (2)  $\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i)\Delta x$
- (3)  $V = \pi \int_a^b y^2 dx$
- (4)  $A = \int_a^b y dx$

**Measurement**

- (1) Cone:  $V = \frac{\pi}{3} r^2 h$
- (2) Cone:  $SA = \pi r l + \pi r^2$
- (3) Sphere:  $V = \frac{4\pi}{3} r^3$

- (4) Sphere:  $SA = 4\pi r^2$
- (5) Cylinder:  $SA = 2\pi rh + 2\pi r^2$
- (6) Cylinder:  $V = \pi r^2 h$
- (7) Circle:  $A = \pi r^2$
- (8) Circle:  $C = 2\pi r$
- (9) Triangle:  $A = \frac{bh}{2}$ ,  $A = \frac{\sqrt{3}}{4} x^2$

$$A = \sqrt{s(s-a)(s-b)(s-c)}$$

- (10) Rectangle:  $A = lw$ ,
- (11) Rectangle  $P = 2l + 2w$
- (12) Square:  $A = s^2$ ,
- (13) Square  $P = 4s$
- (14) Rectangular Prism:  $V = lwh$

Matrices

- (1)  $C_{ij} = (-1)^{i+j} M_{ij}$
- (2)  $AA^{-1} = A^{-1}A = I$
- (3) Inverse of  $A = A^{-1} = \frac{1}{\det A} \cdot adjA$

Data & Probability

- (1)  $\bar{x} = \frac{\sum fx}{n}$
- (2) Median =  $l_1 + \frac{l_2 - l_1}{f1} (m - c)$
- (3)  $\sigma = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n}}$
- (4)  $\sigma_{12} = \sqrt{\frac{n_1\sigma_1^2 + n_1\sigma_2^2 + n_1d_1^2 + n_2d_2^2}{n_1 + n_2}}$

$$(5) \sigma = \sqrt{\frac{\sum f(x_i - \bar{x})^2}{\sum f}} = \sqrt{\frac{\sum fx^2}{N} - \left(\frac{\sum fx}{N}\right)^2}$$

$$(6) \bar{x}_{12} = \frac{m\bar{x}_1 + n\bar{x}_2}{m + n}$$

$$(7) I = \frac{\sum \frac{P_1}{P_0} \times 100}{n}$$

$$(8) I = \frac{\sum p_1 w}{\sum p_0 w} \times 100$$

$$(9) \text{Cov}(X, Y) = \frac{1}{n} \sum (X - \bar{X})(Y - \bar{Y})$$

$$(10) r = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sqrt{\sum (x - \bar{x})^2 \sum (y - \bar{y})^2}}$$

$$(11) r = \frac{\sum (x - \bar{x})(y - \bar{y})}{n\sigma_x \sigma_y}$$

$$(12) b_{YX} = \frac{\text{cov}(X, Y)}{\sigma_x^2} = r \frac{\sigma_y}{\sigma_x}$$

$$(13) Y - \bar{Y} = \frac{\text{cov}(X, Y)}{\sigma_x^2} (X - \bar{X}) = r \frac{\sigma_y}{\sigma_x} (X - \bar{X})$$

$$(14) b_{xy} \times b_{yx} = r \frac{\sigma_y}{\sigma_x} \times r \frac{\sigma_x}{\sigma_y}$$

$$(15) \tau = \frac{2S}{n(n-1)}$$

$$(16) r = 1 - \frac{6\sum d^2}{n(n^2 - 1)}$$



