

# PHYSICS

## Paper – 1

### (THEORY)

*Three hours and a quarter*

*(The first 15 minutes of the examination are for reading the paper only.*

*Candidates must NOT start writing during this time).*

-----

*Answer **all** questions in Part I. From Part II, answer any four questions from Section A, any three questions from Section B and any two questions from Section C.*

*All workings, including rough work, should be done on the same sheet as, and adjacent to the rest of the answer.*

*The intended marks for questions are given in brackets [ ].*

*A list of physical constants is given at the end of the question paper.*

-----

#### **PART I (40 marks)**

*Answer **all** questions.*

#### **Question 1.**

(a) *Each question is followed by four possible choices of answers. Choose the correct answer and write it in your answer sheet.*

[10]

- (i) A moving electric charge produces
- A electric field only.
  - B magnetic field only.
  - C both electric field and magnetic field.
  - D neither of these two fields.


(ii) If the electric field at A and B are  $E_A$  and  $E_B$  and the distance between them is 'r' as shown in the figure given below, then

A  $E_A > E_B$ .

B  $E_A < E_B$ .

C  $E_A = \frac{E_B}{r}$ .

D  $E_A = \frac{E_B}{r^2}$ .



(iii) If an electric current of 2mA flows through a wire, then the number of free electrons passing a given point in a wire per second will be

A  $1.25 \times 10^{13}$ .

B  $1.25 \times 10^{16}$ .

C  $1.25 \times 10^{18}$ .

D  $1.25 \times 10^{21}$ .


(iv) A conducting loop carrying a current 'i' is placed in a uniform magnetic field  $\vec{B}$  pointing into the plane of the paper as shown in the figure. The loop will have a tendency to

A expand.

B contract.

C move towards +ve x-axis.

D move towards -ve x-axis.



- (v) The core of transformers are laminated because it
- A increases the magnetic saturation level of the core.
  - B decreases the residual magnetism of the core.
  - C increases the strength of the magnetic field.
  - D decreases the eddy-current in the core.
- (vi) All the following statements are true for electromagnetic waves *EXCEPT*
- A travels with the same speed as light in free space.
  - B travels with the same speed in all media.
  - C produced by accelerating charges.
  - D are transverse in nature.
- (vii) If an astronomical telescope has a magnifying power 10 and the focal length of the eye piece is 20 cm, then the focal length of the objective is
- A  $\frac{1}{200} \text{ cm}$ .
  - B  $\frac{1}{2} \text{ cm}$ .
  - C  $2 \text{ cm}$ .
  - D  $200 \text{ cm}$ .
- (viii) In the photoelectric effect on metals, an increase in the frequency of incident radiation increases
- A rate of emission.
  - B threshold frequency.
  - C work function of the metal.
  - D velocity of emitted electrons.

- (ix) In which region of the electromagnetic spectrum lies the Lyman series of hydrogen?
- A ultraviolet
  - B infra-red
  - C visible
  - D X-ray
- (x) The work function of a substance is 4.0 eV. The longest wavelength of light that can cause photoelectron emission from this substance is approximately
- A  $3.10 \times 10^{-7} \text{ nm}$ .
  - B  $3.10 \times 10^{-7} \text{ cm}$ .
  - C  $310 \text{ nm}$ .
  - D  $310 \text{ m}$ .
- (b) Choose the correct word/s given in the brackets and write them in your answer sheets. [6]**
- (i) The phenomenon in which energy is converted into mass is called ..... and the phenomenon in which mass is converted into energy is called .....  
(compton-effect, pair-production, photoelectric-effect, pair-annihilation)
- (ii) The n-type semi-conductor is a ..... semiconductor obtained by adding a ..... impurity. (extrinsic, intrinsic, pentavalent, trivalent)
- (iii) The electrolyte in a Lechlanche cell is ..... and its common name is .....  
(manganese dioxide, ammonium chloride, secondary cell, dry cell)
- (iv) A diamagnetic substance is ..... of temperature and its relative permeability is slightly ..... than unity. (dependent, independent, less, more)
- (v) The device used to increase ac voltage is a ..... and ..... be used to step-up dc voltage. (step-up transformer, step-down transformer, can, cannot)
- (vi) The ratio ..... is a time constant and its unit is ..... ( $\frac{L}{R}$ ,  $\frac{R}{L}$ ,  $\frac{H}{\Omega}$ , second)

(c) **Match the items of column A with the items in column B. Rewrite the correct pairs in your answer sheet.**

Column A	Column B
(i) Maxima	(a) varying fringe width
(ii) Interference	(b) infinite
(iii) Holes as majority charge carriers	(c) stationary orbit
(iv) Bohr's Theory	(d) X-ray
(v) Diffraction	(e) n-type semi-conductor
(vi) Resistance of a pure semiconductor	(f) zero
(vii) Minima	(g) same fringe width
(viii) Continuous spectrum	(h) p-type semi-conductor
	(i) even multiple of $\frac{\lambda}{2}$
	(j) odd multiple of $\frac{\lambda}{2}$

(d) **Write True or False and give reasons for the false statements.**


[4]

- (i) A reverse biased p-n junction diode acts as a low resistance instrument.
- (ii) A beam of white light passing through a hollow prism gives no spectrum.
- (iii) In a compound microscope, the aperture of the eye piece is smaller than that of the objective.
- (iv) The depletion layer in the p-n junction is formed by drift of holes.

(e) **Answer the following questions.**

- (i) The work function of photons for photoelectric emission from a metal is 0.05 eV. Find the threshold frequency. [2]
- (ii) With the help of a labelled diagram, show the formation of a rainbow. [2]
- (iii) Establish a relation between half-life and decay constant. [2]

- (iv) Obtain an expression for  $\frac{e}{m}$  of electron in terms of 'v' and 'V' when an electron is accelerated from rest through a potential difference of 'V' volts and it acquires a final velocity 'v'  $\frac{m}{s}$ . [2]
- (v) For neutrino and antineutrino, mention *one* such property which is: [2]
1. common.
  2. opposite.
- (vi) The figure given below shows a fission reaction. Write an equation corresponding to this process. [2]



- (vii) The electric field at a point due to a point charge is  $20 \frac{N}{C}$  and the electric potential at that point is 10 V. Calculate the distance of the point from the charge and the magnitude of the charge. [2]
- (viii) An air solenoid has 500 turns of wire in its 40 cm length. If the current in the wire is 1.0A, find the magnetic field at the axis inside the solenoid. [2]

**PART II****SECTION A (28 marks)**

Answer any **four** questions.

**Question 2.**

- (a) The tyres of aircrafts are not made of ordinary rubber (which is an insulator) but of a special rubber which is slightly conducting. Why? [1]
- (b) A regular pentagon of side 20 cm has a charge  $5 \mu\text{C}$  at each of its vertices. Calculate the electric potential at the centre of the pentagon. [3]
- (c) Establish the formula  $U = \frac{CV^2}{2}$ , for the potential energy of a charged conductor of capacitance 'C' carrying a charge 'Q' at a potential 'V'. [3]


**Question 3.**

- (a) A carbon resistor is marked in orange, green and red bands. What is its approximate resistance? [1]
- (b) Two bulbs whose resistances are in the ratio 1:2 are connected in parallel to a source of constant voltage. [2]
- (i) What will be the ratio of power dissipation in them?
- (ii) If connected in series, then?
- (c) Derive an expression  $\vec{\tau} = \vec{M} \times \vec{B}$  for a rectangular coil of area A carrying a current 'i' placed in a magnetic field. The angle between the direction of field and normal to the plane of coil is  $\theta$ . [4]


**Question 4.**

- (a) How can a galvanometer be converted into an ammeter of desired range? Explain with the help of a diagram. [2]

- (b) The diagram given below has  $T_2 > T_1$ . Explain.



- (c) Calculate the equivalent capacitance of the network shown below between the points 'A' and 'B', given  $C_1 = C_2 = 12\mu F$ ,  $C_3 = 7\mu F$ ,  $C_4 = C_5 = C_6 = 15\mu F$ . [3]



**Question 5.**

- (a) Define internal resistance of a cell. A battery of emf 10V and internal resistance  $3\Omega$  is connected to a resistor. The current in the circuit is 0.5 A. What is the resistance of the resistor? [3]
- (b) Derive the expression for comparison of the magnetic moments of two bar magnets of the same size and mass using a vibration magnetometer. [3]
- (c) State the Curie Law. [1]

**Question 6.**

- (a) Distinguish between diamagnetic, paramagnetic and ferromagnetic substances in terms of their relative permeability and susceptibility. [3]
- (b) What is meant by eddy current? [2]



- (c) Show the growth and decay of current in an L-R circuit with the help of a graph.

**Question 7.**

- (a) Where is the power dissipation in an a.c circuit, in the resistance, inductance, capacitance or in all? [1]
- (b) Why is the choke preferred to a rheostat in controlling the current in an a.c circuit? [2]
- (c) An alternating emf of 100 V (rms), 50 Hz is applied across a capacitor of  $10 \mu\text{F}$  and a resistor of  $100 \Omega$  in series. Calculate: [4]
- (i) the reactance of the capacitor and
- (ii) the impedance of the circuit.

**SECTION B (18 marks)**

*Answer any three questions.*

**Question 8.**

- (a) If the two slits in Young's apparatus are illuminated by two identical but independent monochromatic light sources, will you observe interference pattern on the screen? Give a reason to support your answer. [2]
- (b) Derive Snell's law of refraction for a parallel beam incident on the plane surface of a refracting medium using 'Huygen's Principle'. [4]

**Question 9.**

- (a) Calculate the refractive index of the material of an equilateral prism for which the angle of minimum deviation is  $60^\circ$ . [2]
- (b) With the help of a labelled diagram, derive the expression for the angular width of the central maximum of the diffraction pattern produced by a single slit illuminated with a monochromatic light. [4]

**Question 10.**

- (a) Define dispersive power (for light) of a medium. [1]
- (b) What are the necessary conditions for sustained interference? [2]

- (c) Two thin lens of focal lengths, +10 cm and -5 cm are kept in contact. What is the focal length of the combination?

**Question 11.**

- (a) Explain briefly how the illuminating powers of two sources of light are compared using Bunsen's grease spot photometer. [3]
- (b) Explain the wave particle duality of radiation. [3]

**SECTION C (14 marks)**

Answer any **two** questions.

**Question 12.**

- (a) Calculate the maximum frequency and corresponding wavelength of X-rays produced in a tube maintained at 13.26 KV. [3]
- (b) Explain briefly why there is a maximum frequency for the X-rays produced by an X-ray tube operating at a certain voltage. [2]
- (c) When  ${}^7_3\text{Li}$  is bombarded with a certain particle, two alpha particles are produced. Identify the bombarding particle. [2]

**Question 13.**

- (a) You are given two nucleides  ${}_3X^7$  and  ${}_3Y^4$ . [2]
- (i) Are they the isotope of the same element? Why?
- (ii) Which one of the two is likely to be more stable? Why?
- (b) In  $\beta$ -decay, a neutron is converted into a proton, so neutron-proton ratio decreases. Justify your answer for the equation of  $\beta$ -decay:  ${}_zX^A \rightarrow {}_{z+1}Y^A + {}_{-1}\beta^0 + \bar{\gamma}$  [2]
- (c) Neutrons can be slowed down even by ordinary water which has hydrogen nuclei ( ${}_1H^1$ ) having mass equal to almost that of neutron. Then why is heavy hydrogen used for this purpose in a reactor? [3]

**Question 14.**

- (a) Draw a diagram of a p-n junction diode as a half-wave rectifier and explain its working briefly. [3]
- (b) Explain briefly with reference to semiconductor physics: [4]
  - (i) Reverse biased p-n junction.
  - (ii) How are the radiations given out by light emitting diode (LED)?

**[PHYSICAL CONSTANTS]**

Planck's constant

$$h = 6.63 \times 10^{-34} \text{ J}\cdot\text{s}$$

Electron charge

$$e = 1.6 \times 10^{-19} \text{ C}$$

$$K = \frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ Nm}^2 / \text{C}^2$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ TmA}^{-1}$$

$$C = 3 \times 10^8 \text{ m/s}$$

$$1\mu\text{C} = 1 \times 10^{-6} \text{ C}$$

$$1\text{pF} = 1 \times 10^{-12} \text{ F}$$