CAMBRIDGE INTERNATIONAL EXAMINATIONS

Pre-U Certificate

www. tirenepapers.com

MARK SCHEME for the May/June 2014 series

9790 BIOLOGY

9790/01

Paper 1 (Structured), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, Pre-U, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

Notes:

The following abbreviations may be used in mark schemes:

alternative and acceptable answers for the same marking point

; separates marking points allow/accept/A answers that can be accepted

AVP any valid point – marking points not listed on the mark scheme but which are

worthy of credit

AW/owtte credit alternative wording/or words to that effect

ecf error carried forward

ignore/I statements which are irrelevant – applies to neutral answers

not/reject/R answers which are not worthy of credit

ORA or reverse argument

(words) bracketed words which are not essential to gain credit

words underlined words must be present in answer to score a mark

Page 3	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

Section A

Question Number	Key
1	С
2	1, 3, 5
3	В
4	D
5	Α
6	Α
7	D
8	С
9	С
10	Α
11	В
12	В
13	В
14	В
15	С
16	В
17	В
18	С
19	С
20	D

Page 4	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

Section B

21 (a) (i) for any one time period

1 comparative/manipulated, data quote for decrease in number of (different) seed species (during the drought);

manipulated data June 77 to Dec 77 = 5 species

2 comparative / manipulated, data quote for decrease in (seed) abundance (during the drought);

manipulated data June 77 to Dec 77 = 0.58 cm³ m⁻¹ units to be given for abundance once

for mps 1 and 2 allow data for pre-drought to during drought if sampling dates indicated

March 76/pre-drought, to June 77 = 4 species $6.49 \,\mathrm{cm}^3 \,\mathrm{m}^{-1}$ March 76/pre-drought, to Dec 77 = 9 species $7.07 \,\mathrm{cm}^3 \,\mathrm{m}^{-1}$

if mps 1 and 2 not gained, allow one mark for

1/2 number of (different seed) species and seed abundance decreased;

[2]

(ii) any one relevant e.g.

ref. to easier/faster, than counting;

A more difficult/too small, to count

numbers do not indicate, size of seed/biomass/ORA;

A seeds can be of different sizes

more useful indication of, food/energy, availability;

[max 1]

(iii) 1 time of year (to collect samples);

A ref. to seasonal changes

2 (aim to) sample, within a short time period;

A take samples at the same time

A take samples during same weather conditions

- 3 ref. suitable, sampling area/sample size;
- 4 sampling method, random/systematic;

A described

- 5 qualified/further detail;
 - e.g. computer-generated random locations

choose representative areas on island

description of avoiding particular areas, e.g. craters/lakes

use of belt transects (for systematic)

detail of quadrats (for systematic/random, sampling)

- 6 correct identification of/use key to identify, seed species;
- 7+8 AVP;;
 - e.g. (repeat) samples to be taken at appropriate time intervals collect comparative data (from other years/non-drought years) relevant suggestion ref. seed collection/making estimates

[max 4]

Page 5	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

(b) (i)

percentage change	+6.71		+5.73	
-------------------	-------	--	-------	--

both correct = 1 mark

[1]

(ii) directional (selection);

[1]

- (c) accept birds or ground finches for G. fortis throughout
 - 1 ref. scarce food supply, limits population size / causes decrease in population size;
 - 2 variation in phenotypic variation

or

(individuals show) differences/variation, in mass

or

3

(individuals show) differences/variation, in bill, size/length/depth/width;

mp 2 must be as a general statement

- selection pressure(s) operating/environmental change is selective pressure;
- 4 selective pressure, named/described;
 - e.g. drought/seed supply/seed type (e.g. larger/tougher, seeds most commonly available)
- 5 (intraspecific) competition;
- 6 detail of effect;
 - e.g. bigger bills able to break open large, fruits / seeds larger *G. fortis* more able to compete for food
 - G. fortis with small bills unable to obtain sufficient food
- 7 idea of differential survival;
 - e.g. *G. fortis* with, higher mass/bigger bills, selected for/(better adapted to) survive/AW

(most) G. fortis with smaller, mass/bills, selected against/(more likely to) die

- 8 ref. those selected for/surviving, able to, reproduce/pass on (advantageous) alleles/ORA;
- 9 measurable change over time in phenotype/frequency in population of large, size/beak, increases over time;
- 10 AVP; e.g. different set of selection pressures operating when food abundant not all small *G. fortis* died as able to feed on, other food sources/insects/nectar/pollen

post-drought selection of mates based on similar (large) size

[max 6]

[Total: 15]

Page 6	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

22 (a) endocytosis/phagocytosis/pinocytosis;

further detail; e.g. membrane, invaginates/AW

membrane fusion

vacuole/vesicle, formed

or

diffusion;

R facilitated diffusion

further detail; e.g. across phospholipid bilayer

across hydrophobic core/AW from a high to a low concentration

down a, diffusion/concentration, gradient

[2]

(b) (i) initial/primary, RNA transcripts have, exons and introns/coding and non-coding portions;

introns removed;

A non-coding/AW, sections/regions, removed

exon splicing/AW;

ref. spliceosomes/ribozymes/appropriate enzymes;

[max 2]

- (ii) 1 if (translation on) free ribosomes, remain in cell/AW;
 - 2 (if translation on ribosomes on RER) cytokines/AW, can be exported from cell;
 - 3 polypeptide enters (lumen of) RER (as synthesis occurs);
 - 4 RER involved in, polypeptide/protein/cytokine, transport (through RER)/(post-translational) modification;

A description of modification

- 5 ref. (RER) formation of vesicle to transport to Golgi;
- 6 (transport to) Golgi body for, modification/processing/packaging;

A description e.g. folding, glycosylation

- 7 secretory/Golgi, vesicles transported to cell surface membrane for exocytosis/AW;
- 8 AVP; e.g. signal sequence (of amino acids)/signal sequence recognition protein (SRP), allows attachment to ER membrane

[max 3]

Page 7	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

(c) (i) 1 Langerhans cell/keratinocyte/macrophage, as, antigen-presenting cell/APC;

A displaying hapten in context of recognition

2 ref. T-cells and, recognition/binding;

A clonal selection

A T-cells have complementary receptors/specific T-cells

- 3 activation/sensitisation/described;
- 4 proliferation/mitosis;
 - A clonal expansion
- 5 T (cytotoxic and helper), memory cell formation;
- 6 description of T-cytotoxic cell action;
- 7 explanation of faster response for second and subsequent contacts;
 - e.g. increased number of (specific) T-cells increases chance of coming across APC

more T-cytotoxic cells present to destroy, skin cells/keratinocytes, with hapten

8 T-helper cells secrete cytokine to stimulate, T-cytotoxic cell response/macrophages;

9+10 AVP;;

e.g. ref. cytokine production by keratinocytes attract, macrophages/T-cells, to area

ref. rash linked to effect e.g. pus in blisters = dead, cells/macrophages suggestion of no initial reaction by T-cells because of short life span of keratinocytes

[max 6]

(ii) any one relevant e.g.

immunocompromised/described;

may not have, specific T-cells / T-cells with, quinone / hapten, receptors;

may have T-cells but low in number and not come across APC;

may need several doses to build up sufficient numbers of T-cells;

inability of cells to convert urushiol;

[max 1]

(d) urushiol more rapidly acted upon by, enzyme/catechol oxidase, only/AW; ref. (enzyme) inhibitor linked to red wine/red wine contains an inhibitor; use of comparative data to support;

A comparative description of V_{max}

further detail of inhibition; e.g. co

; e.g. competitive inhibition

similar shape to, substrate/urushiol competes with substrate for active site

fewer enzyme-substrate complexes formed

increasing substrate concentration decreases inhibition

[max 3]

Page 8	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

(e) passes through, epidermis/named leaf cells/cuticle;

to, vascular tissue/phloem;

translocated/mass flow;

symplastic movement;

A if in context of via leaf cells to phloem

from, leaf/source, to sink;

AVP; e.g. detail of mass flow

detail of symplast pathway

[max 3]

[Total: 20]

- **23** (a) (light-dependent stage) provides NADPH and ATP/ORA; further detail;
 - e.g. NADPH provides, electrons/reducing power, for GP to TP ATP required for, GP to TP/regeneration of RuBP

[2]

- (b) 2 molecules of, GP/TP, leave the cycle/AW; allow idea of 6 carbons with reference to GP/TP further detail;
 - e.g. for synthesis of glucose (phosphate) for synthesis of , fatty acids/amino acids

[2]

(c) endosymbiosis;

qualified;

e.g. chloroplasts were small prokaryotes that, lived in host cells/were parasites/were prey

mutually beneficial relationship

[2]

[Total: 6]

Page 9	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

- 24 (a) 1 (glycine) amino acid with, smallest/H, R-group/side chain;
 - 2 run along the, interior/central, axis of the helical polypeptide/AW;
 - 3 allow the, triple helix/three polypeptides, to be, closely associated/tightly packed together/AW;


look for idea of three compact helices

I ref. to alpha helices

- 4 allows H-bonds to form between polypeptides (for strength);
- 5 further detail;
 - e.g. bond forms between the amide (-C(=O)NH-) hydrogen atom of glycine in one polypeptide chain and the carbonyl oxygen atom of X in an adjacent polypeptide chain

[max 3]

(b) (i)



box drawn around R group;

[1]

(ii) OH and H on carbon, 1 reversed/AW; OH and H on carbon 4 reversed/AW; accept from annotations on Fig. 24.3

[max 1]

(c) (i) any one relevant e.g.

no risk of transmissible disease;

(possibility of) large-scale production;

enables production where animal sources are more scarce;

allows (genetic) manipulation to obtain different collagen types;

to avoid, allergy/immune response, problems with, animal products/foreign proteins;

A reduced inflammatory response

[max 1]

(ii) cannot carry out post-translational modification; do not have RER/Golgi;

A do not have necessary, cellular structures/organelles do not have the necessary, genes/enzymes;

[max 1]

Page 10	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

- (d) collagen structure:
 - 1 unable to hydroxylate, proline/lysine;
 - 2 hydrogen bonds not formed;
 - 3 no glycosylation for hydoxylysine (without the OH-);
 - 4 further detail; e.g. lack of stability/compromised structure, of triple helix polypeptide chains more easily separated weaker links between molecules to form collagen fibrils

collagen function:

- 5 ref. collagen, as a structural protein/high tensile strength;
- 6 important in, connective tissue for wound healing/strong gums;
- 7 ref. to presence in, tunica externa/tunica media/sub-endothelial layer, of, arteries/veins;
- 8 weakened blood vessel wall may cause, blood leakage/bursting (hence symptoms);
- 9 ref. role of collagen in blood vessels; e.g. in artery walls to resist high blood pressure
 [max 5]
- (e) (i) (rat cDNA probe binds to a) complementary (single-stranded) sequence in the sample being analysed/AW;

(rat cDNA) probe binding to human DNA indicates region of similar nucleotide sequence exists:

sufficient similarity to support idea of a gene that has lost expression/AW; if rat cDNA probe did not bind then would suggest no gene;

[max 3]

- (ii) 1 frameshift (mutation);
 - 2 from insertion site onwards new sequence of codons read/AW;
 - 3 leads to new sequence of amino acids specified (from point of mutation); needs idea of more than one amino acid changed
 - 4 altered primary stucture alters, tertiary structure/folding, of polypeptide;
 - 5 further detail; e.g. loss/change of shape, of active site, so loss of function
 - 6 change in codons may lead to a nonsense mutation/produce a stop codon;
 - 7 (so) shortened polypeptide/polypeptide produced may not function; in context of premature stop codon
 - 8 AVP; e.g. ref. more drastic effect if insertion early on in sequence

[max 4]

(f) (no need to synthesise as) able to obtain sufficient vitamin C, in the diet/from exogenous source :

A relevant named foods

[1]

[Total: 20]

Page 11	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

25 (a) oestrogen/oestradiol:

repair/thickening, of endometrium/uterine lining;

ref. low/decreased, concentrations of, FSH/LH;

A negative feedback effect on, FSH/LH

(higher concentrations) stimulate/AW, secretion/release, of LH (from anterior pituitary);

A positive feedback effect on LH

A LH surge

progesterone:

stimulates growth/maintenance/increases vascularisation/AW, of, endometrium/uterine lining;

inhibits release of FSH/LH/(anterior) pituitary hormones;

A idea of inhibition by oestrogen released from corpus luteum

FSH/follicle stimulating hormone:

development of follicles;

stimulates, production/secretion, of oestrogen;

LH/luteinising hormone:

stimulates follicle growth (together with FSH);

(high/increasing, levels bring about) ovulation;

stimulates development of corpus luteum (from empty Graafian follicle);

[max 5]

(b) (i) metaphase I;

[1]

(ii) award marks on annotated diagrams

- 1 condensation of chromosomes/described (e.g. coiling and supercoiling of DNA);
- 2 ref. to random orientation of chromosomes;
- 3 homologous chromosomes pair up/synapsis/AW;
- 4 detail; e.g. along length, held by proteins
- 5 crossing over;
- 6 described; e.g. exchange between non-sister chromatids of sections of DNA
- 7 ref. separation of chromosomes, still held together in places;
- 8 chiasmata:
- 9 attachment of, kinetochores/centromeres, to spindle/microtubules/spindle fibres;
- 10 homologous pair/bivalent, movement towards, spindle equator/metaphase plate;

[max 4]

Page 12	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2014	9790	01

(c)

cell type	timing	number of chromosomal DNA molecules per cell
oogonium	newly-formed	46
oogonium	late interphase immediately prior to onset of meiosis	92;
primary oocyte	shortly before puberty begins	92;
secondary oocyte	at ovulation	46;
ovum	after binding of a sperm, but before the sperm nucleus enters the cell	23;

[4]

[Total: 14]

- **26** assume herbivore unless stated otherwise, look for ORA response must include ref. to small and large intestine to score max
 - 1 have a, larger/developed, caecum;
 - 2 ref. to, most/much, digestion occurring in caecum;
 - 3 have, functioning/AW, appendix;
 - 4 (so can carry out) hindgut fermentation;
 - 5 ref. to (mutualistic) bacteria/gut flora/AW;
 - 6 for cellulose breakdown/have cellulase;
 - 7 microorganism metabolism/AW, releases volatile fatty acids;
 - 8 detail of benefit to herbivore; e.g. fatty acids for energy heat for thermoregulation
 - 9 longer, small intestine/large intestine/digestive system;
 - 10 (as) need more, surface area/time, for digestion/absorption;
 - 11 AVP;
 - e.g. little absorption of digested food in small intestine (on first time round) longer intestines as, little digestion in stomach (compared to carnivores) longer/developed, large intestines/colons, to extract water from fibre

[max 5]

[Total: 5]