

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Pre-U Certificate

[www.XtremePapers.com](http://www.XtremePapers.com)

**MARK SCHEME for the May/June 2013 series**

**1347 MATHEMATICS (STATISTICS WITH PURE  
MATHEMATICS)**

**1347/02**

Paper 2 (Statistics), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, Pre-U, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Pre-U – May/June 2013</b>	<b>1347</b>	<b>02</b>

<b>1 (i)</b>	Mean = 2.068 million Median = 2.035 million Values not very different, so no evidence of skew, fairly symmetric	B1 B1 B1	<b>[3]</b>	2.068 or 2068000, accept 2.07 or 2070000 2.035 or 2035000, accept 2.04 or 2040000 Symmetric or equivalent conclusion about the distribution of increases															
	<b>(ii)</b> Interquartile range = $2.52 - 1.35 = 1.17$  Either: $2.52 + 1.5 \times 1.17 = 4.275$  Or: $(3.81 - 2.52) \div 1.17 = 1.103$  3.81 is less than $1.5 \times$ IQR above the upper quartile, so it is not an outlier	M1  M1  A1		IQR calculated correctly, or implied from subsequent working $1.5 \times$ their IQR or 1.755 or $1.29 \div$ their IQR  A valid and correct calculation															
	<b>(iii)</b> The recent increases are all quite small compared with the earlier ones.	B1	<b>[1]</b>	Early values larger than later ones															
<b>2 (i)</b>	1,2 1,3 1,4 1,5 2,3 2,4 2,5 3,4 3,5 4,5	B1		Listing the ten pairs (allow twenty if given in both orders or as a two-way table), must be seen not implied															
	<table border="1"> <tr> <td>Total</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> <td>7</td> <td>8</td> <td>9</td> </tr> <tr> <td>Prob</td> <td>0.1</td> <td>0.1</td> <td>0.2</td> <td>0.2</td> <td>0.2</td> <td>0.1</td> <td>0.1</td> </tr> </table>	Total	3	4	5	6	7	8	9	Prob	0.1	0.1	0.2	0.2	0.2	0.1	0.1	B1 M1 M1 A1	<b>[5]</b>
Total	3	4	5	6	7	8	9												
Prob	0.1	0.1	0.2	0.2	0.2	0.1	0.1												
<b>(ii)</b>	Expectation = $3 \times 0.1 + 4 \times 0.1 + 5 \times 0.2 + \dots = 6$ $3^2 \times 0.1 + 4^2 \times 0.1 + 5^2 \times 0.2 + \dots = 39$ Variance = $39 - 36 = 3$ Standard deviation = $\sqrt{3} = 1.732$	M1 A1 M1 A1 B1	<b>[5]</b>	Or by symmetry, method may be implied 6.0 or 6 39 seen or valid method for variance 3, from valid method seen $\sqrt{3}$ or 1.73 or better (ft from variance)															
<b>3 (i)</b>	$P(\text{robin}) = 0.87$ $X =$ number of children who saw a robin $X \sim B(4, 0.87)$  $P(X < 2) = (0.13)^4 + 4(0.87)(0.13)^3 = 0.00793117$	B1  M1 A1	<b>[3]</b>	Parameters 4 and 0.87 seen or implied  A valid calculation of $P(X < 2)$ or $P(X < 2)$ 0.0079 or greater accuracy															
	<b>(ii)</b> The probability of a robin is 0.87 in the places where these four children live The four gardens are independent  'The gardens are a random sample from the population' $\Rightarrow$ SR2	B1  B1		$p = 0.87$ or constant probability  Independence / random / not all from the same area															
			<b>[2]</b>																

Page 3	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2013	1347	02

<b>(iii)</b>	$H_0: P(\text{see house sparrows}) = 0.65$ $H_1: P(\text{see house sparrows}) < 0.65$  $X =$ no of families who see house sparrows Assuming $H_0: X \sim B(20, 0.65)$ Using tables, either: $P(X < 5) = 0.0003$ or $cv = 7$ $0.0003 < 0.01$ or $0.0003 < 1\%$ or $7 > 5$  Reject $H_0$ (accept $H_1$ )  Evidence supports the claim that the probability of seeing house sparrows in city areas is less than 65%	B1 B1  M1  A1 B1  M1  A1  <b>[7]</b>	‘Probability of seeing house sparrows’ $< 0.65$ or $< 65\%$  Using $B(20, 0.65)$ , or implied from subsequent working $0.0003$ or $7$ from correct method Correct comparison seen  Reject $H_0$ or equivalent, correct for their tail probability or $cv$ Correct conclusion in words
<b>4 (i)</b>	$S_{xx} = 4171 - \frac{(143)^2}{5} = 81.2$ $S_{yy} = 432 - \frac{(46)^2}{5} = 8.8$ $S_{xy} = 1342 - \frac{143 \times 46}{5} = 26.4$  $r = \frac{26.4}{\sqrt{81.2 \times 8.8}} = 0.9876$ (0.988)  $r$ is very near 1, so a good fit to (an upward sloping) line  <b>(ii)</b> $b = \frac{26.4}{81.2} = 0.3251\dots$ $a = \frac{46}{5} - 0.3251 \times \frac{143}{5}$ $= 9.2 - 0.3251 \times 28.6 = -0.0985$ $y = 0.325x - 0.0985$ $x = 20 \Rightarrow \hat{y} = 6.40$ or 6  Extrapolation beyond range of data Small sample / only based on one sample	B1  B1  B1  M1  A1  <b>[5]</b>  M1  M1  A1  M1  A1  B1 B1 <b>[7]</b>	81.2  8.8  26.4  Calculating $r$ from their $S_{xx}$ , $S_{yy}$ and $S_{xy}$ (numerical working or their $r$ value correct to 3 sf or better) Drawing a valid conclusion (confirming that a linear fit is appropriate, as stated in question)  Calculating $b$ from <i>their</i> $S_{xx}$ , $S_{xy}$ (calculation seen or 0.32 to 0.33)  Calculating $a$ from $\Sigma x$ , $\Sigma y$ and <i>their</i> $b$ (seen or implied) $-0.10$ to $-0.09$ for $a$ (without wrong working) Line correct with coefficients 0.32 to 0.33 and $-0.10$ to $-0.09$ used correctly $6.3$ to $6.5$ (before rounding) or integer 6  Extrapolation Any valid objection
<b>5 (i)</b>	$H_0: \text{population mean} = 0.2$ $H_1: \text{population mean} > 0.2$  <b>(ii)</b> $X =$ reaction time Assuming $H_0: X \sim N(0.2, 0.001)$ $0.2 + 1.645 \times \sqrt{0.001} = 0.252$  Would not expect a reaction time $> 0.252$ if $H_0$ is true. If the reaction time is greater than 0.252 it provides evidence that Sam is right and $\mu > 0.2$ , otherwise there is insufficient evidence to reject $H_0$ .	B1  B1 <b>[2]</b>  M1 A1  B1  <b>[3]</b>	‘population mean’ or $\mu$ (but not just ‘mean’) $> 0.2$  Use of 1.645 $cv = 0.252$  Explaining how ( <i>their</i> ) $cv$ is used

Page 4	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2013	1347	02

(iii)	<p>P(Type II error)  <math>= P(X &lt; 0.252)</math> where <math>X \sim N(0.24, 0.001)</math>  <math>= P(Z &lt; 0.379) = 0.6477</math></p>	<p>M1  M1  A1  <b>[3]</b></p>	<p><math>X \sim N(0.24, 0.001)</math> used or implied  <math>P(X &lt; \text{or } &lt; \text{cv})</math>  0.647 to 0.648 as final answer without wrong working, cao</p>
6 (i)	<p><math>H_0</math>: No association between attractiveness of men and their wives  <math>H_1</math>: Some association</p>	<p>B1  <b>[1]</b></p>	<p><math>H_0</math>: No association (allow independence)  <math>H_1</math>: Association, or equivalent</p>
(ii)	<p><math>\frac{18}{50} \times \frac{16}{50} \times 50</math> or <math>18 \times 16 \div 50 = 5.76</math></p>	<p>M1  A1  <b>[2]</b></p>	<p><math>18 \times 16, 288, 0.32, 0.36</math> or <math>0.1152</math> seen  Correct method (answer given in question)</p>
(iii)	<p><math>\frac{(10 - 5.76)^2}{5.76} = \frac{4.24^2}{5.76} = \frac{17.9776}{5.76} = 3.12</math></p>	<p>B1  B1  <b>[2]</b></p>	<p><math>(10 - 5.76)^2</math>  <math>\div 5.76</math></p>
(iv)	<p><math>\nu = (3 - 1)(3 - 1) = 4</math> degrees of freedom  Critical value = 13.28  <math>38.24 &gt; 13.28 \Rightarrow</math> reject <math>H_0</math> (accept <math>H_1</math>)  More attractive men usually have more attractive wives and less attractive men have less attractive wives.</p>	<p>M1  A1  M1  A1  <b>[4]</b></p>	<p>4 df  13.28  Reject <math>H_0</math> or equivalent, correct for their cv  A correct conclusion in words</p>
7	<p>Weight loss 8 15 17 10 21 24 32 6 19 46  Rank 2 4 5 3 7 8 9 1 6 10  Type N N N N N N N E E E</p> <p>Rank sum E = 17 (<math>R_m = 17</math>)  <math>m(m + n + 1) - R_m = 16</math>  <math>W = 16</math></p> <p><math>H_0</math>: same distribution <math>\alpha = 5\%</math>  <math>H_1</math>: different distributions two-sided</p> <p><math>m = 3, n = 7 \Rightarrow \text{cv} = 7</math></p> <p><math>16 &gt; 7 \Rightarrow</math> accept <math>H_0</math>  Insufficient evidence to claim that there is a difference in the weight losses.</p>	<p>B1  M1  M1  A1  B1  M1  A1  <b>[7]</b></p>	<p>Ranks correct (at least for E)</p> <p><math>R_m = 17</math>  <math>m(m+n+1) - R_m</math>  <math>W = 16</math>, cao</p> <p>cv = 7</p> <p>Accept <math>H_0</math> (ft <i>their</i> <math>W</math> and cv)  Correct conclusion in words, from correct working for <i>their</i> <math>W</math> and <math>c</math>.</p>

Page 5	Mark Scheme	Syllabus	Paper
	Pre-U – May/June 2013	1347	02

<b>8</b>	<b>(i)</b> Estimate $\mu$ using $\bar{x} = \frac{2144}{16} = 134$ Estimate $\sigma^2$ using $s^2 = \frac{8640}{15} = 576$	B1 M1 A1 [3]	134 8640 $\div$ either 15 or 16, 540, 24 or 23.2 576
	<b>(ii)</b> Population distribution is normal and unknown population variance Small sample	B1 B1 [2]	Normal and $\sigma^2$ unknown Sample size
	<b>(iii)</b> 15 degrees of freedom $\Rightarrow t = 2.131$  $134 \pm 2.131 \times \sqrt{(576/16)}$ $= 134 \pm 2.131 \times 6$ $= (121.2, 146.8)$	M1 A1  B1 M1 A1 [5]	$n - 1 = 15$ 2.131  576/16 seen or implied Correct method for <i>their</i> $t$ or $z$ value, condone omission of 16 Confidence limits correct (3 sf or better)
	<b>(iv)</b> $\bar{X} \sim N(145, 100)$  $P(\bar{X} < 125) = P(Z < \frac{125 - 145}{10})$ $P(Z < -2) = 1 - 0.9772$ $= 0.0228$	B1  M1 M1 A1 [4]	$N(145, 100)$  Correct method for their distribution $Z$ value $-2$ or $+2$ 0.0228