#### PURE MATHEMATICS, PAPER-II



# FEDERAL PUBLIC SERVICE COMMISSION COMPETITIVE EXAMINATION FOR RECRUITMENT TO POSTS IN BPS-17 UNDER THE FEDERAL GOVERNMENT, 2010

## **PURE MATHEMATICS, PAPER-II**

TIME ALLOWED: 3 HOURS

Student Bounty.com **MAXIMUM MARKS:100** 

NOTE:

- (i) Attempt FIVE questions in all by selecting at least THREE questions from SECTION-A and TWO questions from SECTION-B. All questions carry EQUAL marks.
- (ii) Use of Scientific Calculator is allowed.

## SECTION - A

- If f is continuous on [a,b] and if  $\infty$  is of bounded variation on [a,b], then  $f \in R(\infty)$  on [a, b] i.e. f **Q.1.** (a) is Riemann – integrable with respect to  $\infty$  on [a,b] (10)
  - Let  $\sum a_n$  be an absolutely convergent series having sum S. then every rearrangement of  $\sum a_n$ also converges absolutely & has sum S. (10)
- For what +ve value of P,  $\int_{0}^{1} \frac{dn}{(1-x)^{p}}$  is convergent? **Q.2.** (a) (10)

(b) Evaluate 
$$\int_{1}^{5} \frac{dx}{\sqrt[3]{x-2}}$$
 (10)

**Q.3.** (a) Find the vertical and horizontal asymptotes of the graph of function:

$$f(x) = (2x+3)\sqrt{x^2 - 2x + 3}$$
 (10)

(b) Let (i)  $y = f(x) = \frac{(x+2)(x-1)}{(x-3)^2}$ (ii)  $y=f(x) = \frac{(x-1)}{(x+3)(x-2)}$ 

(ii) 
$$y=f(x) = \frac{(x-1)}{(x+3)(x-2)}$$
 (10)

Examine what happens to y when  $x \to -\infty$  &  $x \to +\infty$ 

- **Q.4.** (a) Find a power series about 0 that represent  $\frac{x}{1-x^3}$ **(6)** 
  - (b) Let  $\sum_{s}$  be any series, Justify. (5+5+4)
    - (i) if  $\lim_{n\to\infty} \left| \frac{Sn+1}{Sn} \right| = r < 1$ , then  $\sum_{n=1}^{\infty} s_n$  is absolutely convergent.
    - (ii) if  $\lim_{n\to\infty} \left| \frac{Sn+1}{Sn} \right| = r$  and  $(r > 1 \text{ or } r = \infty)$ , then  $\int_{n}^{\infty} diverges$ .
    - (iii) if  $\lim_{n\to\infty} \left| \frac{Sn+1}{Sn} \right| = 1$ , then we can draw no conclusion about the convergence or divergence.

PURE MATHEMATICS, PAPER-II

Q.5. (a) Show that 
$$\int_{0}^{\Pi 12} Sin^{2m-1}\theta \cos^{2n-1}\theta d\theta = \frac{\Gamma(m)\Gamma(n)}{2\Gamma(m+n)}; m, n > 0$$

- (b) Prove that  $\beta(m,n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}; m,n,>0$
- "MENTBOUNTY.COM Let A be a sequentially compact subset of a matrix space X. Prove that A is totally **Q.6.** (a) bounded.
  - Let A be compact subset of a metric space (X,d) and let B be a closed subset of X such that  $A \cap B = \Phi$  show that d(A,B) > 0(10)

## SECTION - B

- Show that if tanZ is expanded into Laurent series about  $Z = \frac{11}{2}$ , then **Q.7.** (a) (10)
  - Principal is  $\frac{-1}{z \Pi/2}$
  - (ii) Series converges for  $0 < |Z \frac{\Pi}{2}| < \frac{\Pi}{2}$
  - (b) Evaluate  $\frac{1}{2\Pi i} \oint \frac{e^{zi}}{z^2(z^2+2z+2)} dz$  around the circle with equation |z|=3. (10)
- **Q.8.** (a) Expand  $f(x) = x^2$ ;  $0 < x < 2\Pi$  in a Fourier series if period is  $2\Pi$ . (10)
  - (b) Show that  $\int_{0}^{\infty} \frac{Cosxdx}{x^2 + 1} = \frac{\Pi}{a} e^{-x}; x \ge 0$ (10)
- Let f(z) be analytic inside and on the simple close curve except at a pole of **Q.9.** (a) order m inside C. Prove that the residue of f(Z) at a is given

by 
$$a_{-1} = \lim_{Z \to a} \frac{1}{(m-1)!} \frac{m^{-1}d}{dz^{m-1}} \{ (z-a)^m f(z) \}$$
 (10)

(b) If f(z) s analytic inside a circle C with center at a, then for all Z inside C.

$$f(z) = f(a) + f'(a)(z-a) + f''(\frac{a}{2!}(z-a)^2 + f'''(\frac{a}{3!}(z-a)^3 + \dots$$
 (10)

\*\*\*\*\*\*