Answer Sheet No	CATAL .
Sig. of Invigilator	THAT

MATHEMATICS HSSC-II

SECTION - A (Marks 20)

Time allowed: 25 Minutes

NOTE: Section—A is compulsory and comprises pages 1–2. All parts of this section are to be answered on the question paper itself. It should be completed in the first 25 minutes and handed over to the Centre Superintendent. Deleting/overwriting is not allowed. Do not use lead pencil.

(i)	y = 0	sin x is a								
	A.	Linear functi	on		B.	Constant fur	iction			
	С	Rational fund	ction		D.	Trigonometri	ic functio	n		
(ii)	x.2 +	$-xy + y^2 = 2 \text{ is}$	a/an							
	A.	A Implicit function				Inverse function				
	C.	Explicit function	tion		D.	Constant function				
(iii)	If <i>f</i> (If $f(\theta) = 2\sin\theta + 3\cos\theta$ then $f(\theta)$ is a/an								
	A.	A. Odd function				Even function				
	C.	Hyperbolic fo	unction		D.	None of thes	e			
(iv)	$\lim_{x\to a}$	$\frac{x^n - a^n}{x - a} = \underline{\hspace{1cm}}$								
	Α.	na^{n-1}	B.	na^n	C.	na^{n+1}	D.	None of these		
(v)	lim n->x	$\left(1 + \frac{1}{n}\right)^{n/2}$ is	·							
	Α.	$\sqrt[3]{e}$	B.	\sqrt{c}	C.	$\sqrt[1]{e'}$	D	None of these		
(vi)	Equa	tion of parabola	ı is							
	A.	$x^2 + y^2 = 1$	B.	x = y	C.	$x^2 = 4ay$	Đ.	None of these		
(vii)	If f ($x) = \sin x$ and	q(x) = .	$\sin^{-1}x$ then_						
	$(gof)x = sin^{-1}(\sin x)$				B.	X				
	C.	$\sin x + \sin^{-1}$	X		D.	None of thes	е			
(viii)	Li m δx →a	<u>.</u> .	$\frac{C(x)}{x}$ is C	called						
	A.	Velocity			B.	Derivative				
	C.	Integration			Ð.	None of thes	۵			

DO NOT WRITE ANYTHING HERE

SkudentBounty.com

(x)
$$\int e^x (\tan x + sec^2 x) dx \quad \text{then} \quad \underline{\hspace{1cm}}$$

A.
$$e^x \tan x + c$$

B.
$$e^x sec^2 x + c$$

$$C. \qquad e^x \cot^{-1} x + c$$

D. None of these

(xi) Distance of the line
$$-x+3y+1=0$$
 from $(0,2)$ is _____

A.
$$\frac{7}{\sqrt{10}}$$

B.
$$\frac{\sqrt{7}}{10}$$

C.
$$\sqrt{\frac{7}{10}}$$

D.
$$\frac{10}{\sqrt{7}}$$

(xii) If the lines
$$ax - y - 1 = 0$$
 and $x + y + 1 = 0$ are perpendicular to each other then

A.
$$-j$$

B.
$$\bar{k}$$

D.
$$\frac{-1}{j}$$

C.
$$-2a$$

(xvi) The value of c for
$$y = mx + c$$
 touch the ellipse at (x, y) is _____

$$\Delta \qquad \sqrt{a^2 m^2 - h^2}$$

$$\sqrt{a^2m^2-b^2}$$
 B. $\sqrt{3a^2+b^2}$ C.

$$\sqrt{a^2m^2+b^2}$$

(xvii) If the vector
$$\vec{a}$$
 and \vec{b} are perpendicular to each other then $\vec{a} \times \vec{b}$ is ______

A.
$$\sqrt{|a||b|}$$

B.
$$|a|b|$$

C.
$$|a|^2 |b|^2$$
 D.

(xviii) The vectors
$$\vec{a}$$
, \vec{b} and \vec{c} are said to be coplanar if \vec{a} . $(\vec{b} \times \vec{c})$ is ______

c.
$$\sqrt{2}$$

A.
$$(\pm ae, 0)$$

B.
$$(0,\pm ae)$$

C.
$$(\pm a, 0)$$

(xx) The area of the triangle having
$$\vec{a}$$
 and \vec{b} as two sides is given by _____

A.
$$\overrightarrow{a} \times \overrightarrow{b} = c$$

A.
$$\vec{a} \times \vec{b} = o$$
 B. $\frac{1}{2} |\vec{a} \times \vec{b}|$ C. $\vec{a} \cdot \vec{b}$ D.

For Examiner's use only:

Total Marks:

20

Marks Obtained:

- 2HA 1311 (L) ----



MATHEMATICS HSSC-II



Time allowed: 2:35 Hours

Total Marks Sections B

Student Bounty.com NOTE: Attempt any ten parts from Section 'B' and any five questions from Section 'C' on the se provided answer book. Use supplementary answer sheet i.e. Sheet-B if required. Write your ans neatly and legibly.

SECTION - B (Marks 40)

Attempt any TEN parts. All parts carry equal marks. Q. 2

 $(10 \times 4 = 40)$

(i) Evaluate
$$\lim_{\theta \to 0} \frac{1 - \cos p \theta}{1 - \cos q \theta}$$

Find the values of m and n $\,$ so that the given function $\,f\,$ is continuous: (ii)

$$f(x) = \begin{cases} mx & \text{if } x < 3\\ n & \text{if } x = 3\\ -2x + 9 & \text{if } x > 3 \end{cases}$$

(iii) If
$$y = \sqrt{x} - \frac{1}{\sqrt{x}}$$
, then show that $2x \frac{dy}{dx} + y = 2\sqrt{x}$.

(iv) Differentiate ab – ignition w.r.t
$$x$$
 if $y = sin\sqrt{x}$

(v) Prove that
$$e^{x+h} = e^x \{1 + h + \frac{h^2}{2} + \frac{h^3}{3} + \dots \}$$

(vi) Evaluate
$$\int (\ln x)^2 dx$$

Write the equation of the parabola with focus (-1,0) and vertex (-1,2). (vii)

(viii) Solve
$$\frac{dy}{dx} = \frac{y^2 + 1}{e^{-x}}$$

Find an equation of an ellipse with foci $(-3\sqrt{3},0)$ and vertices $(\pm 6,0)$. (ix)

Write the equation of the tangent to the conic $3x^2 - 7y^2 = 20$ at the points where y = -1(x)

Show that 10xy + 8x - 15y - 12 = 0 represents a pair of straight lines. (xi)

Find the values of "a" and "b" so that the vectors $3\underline{i} - \underline{j} + 4\underline{k}$ and $a\underline{i} + b\underline{j} - 2\underline{k}$ are parallel. (xii)

Find 'k' so that the line joining A(7,3); B(k,-6) and the line joining C(-4,5) and D(-6,4) are (xiii) perpendicular.

Find measure of the angle between the lines represented by $x^2 - xy - 6y^2 = 0$. (xiv)

SECTION - C (Marks 40)

Attempt any FIVE questions. All questions carry equal marks.

 $(5 \times 8 = 40)$

Q. 3 Prove that
$$\lim_{x \to 0} \frac{\sqrt{x+a} - \sqrt{a}}{x} = \frac{1}{2\sqrt{a}}$$

Expand a^x in the Maclaurin series. Q. 4

Q. 5 Evaluate
$$\int \frac{e^x(x^2+1)}{(x+1)^2} dx$$

Note:

Q. 6 If
$$x = a(\theta + \sin \theta)$$
; $y = a(1 - \cos \theta)$ then show that $y^2 \frac{d^2y}{dx^2} + a = 0$

The points A(-1,2), B(6,3) and C(2,-4) are vertices of a triangle. Show that the line joining the Q. 7 midpoint D of AB and mid point E of AC is parallel to BC and $DE = \frac{1}{2}BC$

Find the center, foci, eccentricity and vertices of $\frac{(x-1)^2}{2} - \frac{(y-1)^2}{6} = 1$. Q. 8

Find a vector perpendicular to each of the vectors $\vec{a} = 2\vec{i} + \vec{j} + \vec{k}$ and $\vec{b} = 4\vec{i} + 2\vec{j} - \vec{k}$ Q. 9



Roll No.

Sig. of Candidate.

_				
Δ.	nswei	r Sh	eet	No.

Student Bounty Com Sig. of Invigilator._

MATHEMATICS HSSC-II

SECTION - A (Marks 20)

Time allowed: 25 Minutes

NOTE: Section-A is compulsory and comprises pages 1-2. All parts of this section are to be answered on the question paper itself. It should be completed in the first 25 minutes and handed over to the Centre Superintendent. Deleting/overwriting is not allowed. Do not use lead pencil.

(i)	Functions are used to explain the relationship between								
	A.	Variable quan	tities			В.	Notations		
	С	Values				D	None of these		
Cita	Volui	me of a sphere de	epends	on its					
	À.	π				B.	4/3		
	С	Radius				D.	Centre		
(iii)	A fur	nction in which the	variab	ole app	ears as exp	onent i	is called a/an	<u>.</u>	
	А	Rational functi	ion			В.	Exponential fu	inction	
	С	Hyperbolic fun	ction			D.	Inverse function	on	
(IV)	L im	$\frac{\sin x''}{x}$ is							
	Α.	1				B.	$\frac{\pi}{90}$		
	С	180 7				D.	45 π		
(v)	$\frac{d}{dx}$	$\frac{1}{\sqrt{x^+}}$ is							
	Α.	$\sqrt[3]{x^4}$	В.		$\frac{1}{3}$	C.	$\frac{1}{2}$	D	None of these
(Vi)		$+y^2 = 4$ then $\frac{dy}{dx}$							
	A.	$-\frac{Y}{X}$	В.	$\frac{x}{v}$		C.	AV V	D.	None of these
(vii)		$x) \circ a^x$ then f^0							
	Α.	a` l na				B.	σ^X		
	C.	Lna				D.	None of these		
(viii)	ii) A function is said to be increasing function i			nction if					
	A.	$f(x_1) = f(x_2)$				В.	$f(x_1) > f(x_1)$		
		$f(x_2) < f(x_1)$				D.	None of these		
(ix)		dx							
	Α.	$a^X + c$	В.	a^{X}	l na+c	C.	$\frac{a^{X}}{l n a} + c$	D.	None of these

DO NOT WRITE ANYTHING HERE

A.
$$\frac{\pi}{3}$$
 rad B. $\frac{\pi}{6}$ rad C. π rad D. $\frac{\pi}{2}$ rad

B.
$$\frac{\pi}{6}$$
 rad

D.
$$\frac{\pi}{2}$$
 rac

SkudentBounty.com

Solution of differential equation $\frac{dy}{dx} = -tan x$ is _____ (xi)

A.
$$y = l n \cos x + c$$

B.
$$xy = l n \cos x$$

C.
$$x = l n \cos y + c$$

Each equal part of a plane is called _____ (xii)

Where does the point (0,5) lie? (iiix)

A.
$$x - axis$$

B.
$$y - axis$$

Equation of the line parallel to y-axis through (3,7) is _____ (xiv)

A.
$$x = 7$$

B.
$$y = 3$$

C.
$$y = 7$$

D.
$$x = 3$$

Equation of the line through (-6,5) with slope 7 is ____ (vv)

A.
$$x + 7y + 47 = 0$$

B.
$$7x + y + 47 = 0$$

C.
$$7x - y + 47 = 0$$

(ivx) Slope of the line (1+7k)x + (k-1)y - 4 + 20k = 0 is____

A.
$$\frac{k+1}{1+7k}$$

$$B. \qquad -\frac{7k+1}{k-1} \qquad C.$$

$$\sum_{k=1}^{\infty} \frac{k-1}{1+7k}$$

Two lines represented by $ax^2 + 2hxy + by^2 = 0$ are parallel if (xvii)

$$h^2$$
 $ah = 0$

$$h^2$$
 $ab = 0$

$$h^2 - ab < 0$$
 B. $h^2 - ab = 0$ C. $h^2 - ab > ab$ D.

$$h^2 - ab = ab$$

Conics are the curves obtained by cutting a right circular cone by a____ (iiivx)

A unit vector $\,\hat{n}$ perpendicular to $\,ec{a}\,$ and $\,ec{b}\,$ is $\,$ _____ (xix)

A.
$$\frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$$

$$\vec{a} \times \vec{a}$$

$$\sum \frac{\overrightarrow{a}.\overrightarrow{b}}{|\overrightarrow{a}.\overrightarrow{b}|}$$

$$\overrightarrow{a}.(\overrightarrow{b}\times\overrightarrow{c})$$

3

If three points are collinear then area of a triangle will be (xx)

For Examiner's use only:

Total Marks:

20

Marks Obtained:

2HA 1311 (ON) ---



MATHEMATICS HSSC-II



Time allowed: 2:35 Hours

Total Marks Sections B and

SHILDENT BOUNTS! COM NOTE: Answer any ten parts from Section 'B' and any five questions from Section 'C' on the separate provided answer book. Use supplementary answer sheet i.e. Sheet-B if required. Write your answers neatly and legibly.

SECTION - B (Marks 40)

Q. 2 Attempt any TEN parts. All parts carry equal marks.

 $(10 \times 4 = 40)$

(i) Evaluate
$$\lim_{x \to a} \frac{\sqrt{x+a} - \sqrt{a}}{x}$$

(ii) If
$$f(x) = \begin{cases} 3x & \text{if } x \le -2 \\ x^2 - 1 \text{if } -2 < x < 2 \end{cases}$$
 Discuss the continuity at $x = -2$

$$\begin{cases} 3x & \text{if } x \le -2 \\ 3 & \text{if } x \ge 2 \end{cases}$$

(iii) Prove that
$$y \frac{dy}{dx} + x = 0$$
 if $x = \frac{1 - t^2}{1 + t^2}$, $y = \frac{2t}{1 + t^2}$

(iv) If
$$y = tan(2 tan^{-1} \frac{x}{2})$$
 show $\frac{dy}{dx} = \frac{4(1+y^2)}{4+x^2}$

(v) Find the extreme value of
$$f(x) = 5 + 3x - x^3$$

(vi) Use differential to approximate the value of
$$(31)^{\frac{1}{25}}$$

(vii) Evaluate
$$\int tan^2x \ dx$$

(viii) Find the area bounded by
$$y = x(x^2 - 4)$$
 and the $x - axis$.

(ix) Find an equation of the parabola whose focus is
$$F(-3, 4)$$
 and directrix is $3x - 4y + 5 = 0$.

(x) Find the angle between the lines represented by
$$ax^2 + 2hxy + by^2 = 0$$

(xi) Find an equation of the ellipse with vertices
$$(0,\pm 5)$$
 and eccentricity $\frac{3}{5}$

(xii) Show that the product of the distances from the foci to any tangent to the hyperbola
$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$
 is constant.

(xiii) In any triangle
$$\triangle ABC$$
 prove that $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

(xiv) Find analytic expression of
$$u.(v \times w)$$

SECTION - C (Marks 40)

Attempt any FIVE questions. All questions carry equal marks. Note:

 $(5 \times 8 = 40)$

Q. 3 If
$$\theta$$
 is measured in radian then show that $\lim_{\theta \to 0} \frac{\sin \theta}{\theta} = 1$

Q. 4 Find from definition the differential co-efficient of
$$(ax+b)^n$$
 $wx.t^nx^n$ when n is a positive integer.

Q. 5 If
$$y = a\cos(\ln x) + b\sin(\ln x)$$
 then prove that $x^2 \frac{d^2y}{dx^2} + x\frac{dy}{dx} + y = 0$

Q. 6 Evaluate
$$\int \frac{3}{x(x^3-1)} dx$$
, $x \neq 0$, $x \neq -1$.

Q. 7 Find equations of altitudes of a triangle whose vertices are
$$A(-3,2)$$
, $B(5,4)$ and $C(3,-8)$.

Q. 8 Find the area of the region bounded by
$$10x^2 - xy - 21y^2 = 0$$
 and $x + y + 1 = 0$.

Q.9 Graph the solution region of the system of linear inequalities:
$$2x + y \le 10$$

$$x + y \leq 7$$

$$-2x + y \leq 4$$