
G	ITEFORUM				
Q. 1 – Q. 25 carry one mark each.					
Q.1	Image: Image				
Q.2	Which one of the following modifications is common toboth protein and DNA? (A) SUMOylation (B) Nitrosylation (C) Methylation (D) Ubiquitination				
Q.3	Protein A, which has strong affinity to Fc region of immunoglobulin, is extracted from(A) Saccharomyces cerevisiae(B) Staphylococcus aureus(C) Streptococcus pyogenes(D) Streptococcus sanguis				
Q.4	The first humanized monoclonal antibody approved for the treatment of breast canceris (A) Rituximab (B) Cetuximab (C) Bevacizumab (D) Herceptin				
Q.5	Which one of the following aminoacids in proteins does NOT undergo phosphorylation?(A) Ser(B) Thr(C) Pro(D)Tyr				
Q6.	The role of an aduvant isto Data Description Success (A) prolong the persistence of antigen (B) cross link the antigen (C) increase the size of antigen (D) avoid inflammation				
Q7.	Endogenous antigens are presented on to the cell surface along with (A) MHC-II (B) MHC-I (C) Fcyreceptor (D) complement receptor				
Q.8	Human genome sequencing project involved the construction of genomic library in(A) Bacterial artificial chromosome(B) pBR322(C) Bacteriophage(D) pcDNA3.1				
Q.9	The nucleotide analogue used in DNA sequencing by chain termination method is (A) 1',3'-dideoxy nucleoside triphosphate (B) 2',3'-dideoxy nucleoside triphosphate (C) 2',4'-dideoxy nucleoside triphosphate (D) 2',5'-dideoxy nucleoside triphosphate				
Q.10	 In nature, the horizontal gene transfer across bacteria is mediated by (A) Gene cloning followed by transformation (B) Conjugation and transformation (C) Conjugation only (D) Transformation only 				



Q.20 One of the eigen value of $P = \begin{bmatrix} 10 & -4 \\ 18 & -12 \end{bmatrix}$ is (A) 2 (B) 4 (C) 6 (D) 8

|BT-GATE-2013 PAPER|

Q. 21 – Q. 25 are of numerical answer type.

- veig veig 0.21 A callus of 5 g dry weight was inoculated on semi-solid medium for growth. The dry weight of the callus was found to increase by 1.5 fold after 10 days of inoculation. The growth index of the culture is
- Q.22 A chemostat is operated at a dilution rate of 0.6 h⁻¹. At steady state, the biomass concentration in the exit stream was found to be 30 g l^{-1} . The biomass productivity (g $l^{-1}h^{-1}$) after 3h of steady state operation will be
- Q.23 A batch bioreactor is to be scaled up from 10 to 10,000 liters. The diameter of the large bioreactor is 10 times that of the small bioreactor. The agitator speed in the small bioreactor is 450 rpm. Determine the agitator speed (rpm) of the large bioreactor with same impeller tip speed as that of the small bioreactor
- Q.24 Calculate the percentage sequence identity for the pairwise alignment given below.

HELLO-YELLOW

Q.25 In a batch culture, the specific rate of substrate utilization is 0.25 g (g cell mass)-1 h-1 and specific rate of product formation is 0.215 g (g cell mass)-1 h-1. Calculate the yield of product from the substrate(Yp/s)

Q. 26 to Q. 55 carry two marks each.

Q.26 Match the commercial microbial sources in **Group I** with the products in **Group II**.

Group I

- P. Corynebacteriumlilium
- Q. Klebsiellaoxytoca
- R. Aspergillusniger
- S. Alcaligeneseutrophus
- (A) P-3,Q-1,R-2,S-4
- (C) P-1,Q-3,R-2,S-4

Group II

- 1. 2,3-Butane di-ol
- 2. Poly-β-hydroxybutyric acid
- 3. Glutamic acid
- 4. Citric acid
- (B) P-3,Q-1,R-4,S-2
 - (D) P-1,Q-3,R-4,S-2

Q.27 Match the entries in the Group I with the elution conditions in Group II.

Group I

- P. Ion-exchange chromatography
- Q. Hydrophobic column chromatography
- R. Gel filtration chromatography
- S. Chromatofocusing
- (A) P-4,Q-1,R-2,S-3
- (C) P-3,Q-4,R-1,S-2

- 1. Isocratic solvent
- 2. Ampholytes

Group II

- Increasing gradient of salt
- 4. Decreasing gradient of polarity
- (B) P-4,Q-3,R-1,S-2 (D) P-3,Q-4,R-2,S-1
- Determine the correctness or otherwise of the following **Assertion** (a) and **Reason** (r). Q.28 Assertion: Immobilization of plant cells can enhance secondary metabolite production during bioroactor cultivation

|BT-GATE-2013 PAPER|

StudentBounty.com Reason: Immobilization protects the plant cells from shear forces in the bioreact

- (A) Both (a) and (r) are true and (r) is the correct reason for (a).
- (B) Both (a) and (r) are true but (r) is not the correct reason for (a).
- (C) (a) is true but (r) is false.
- (D) (a) is false but (r) is true.

Q.29 Match the cell structures in **Group I** with the organisms in **Group II**.

Group I

P. Endospores

Q. Bipolar flagella


S. Periplasmic flagella

(A) P-4, Q-3, R-1, S-2 (C) P-3, Q-4, R-1, S-2

Group II

1. Methanobacterium 2. Treponema R. Pseudomurine in cell wall 3. Spirillum 4. Clostridium (B) P-4, Q-3, R-2, S-1 (D) P-4, Q-1, R-3, S-2

Q.30 Match the antibioticsin**Group I** with the targets in**Group II**.



Q.31 In nature, Agrobacterium tumefaciens mediated infection of plant cells leads to P. crown gall disease in plants


- Q. hairy root disease in plants
- R. transfer of T-DNA into the plant chromosome
- S. transfer of Ri-plasmid into the plant cell
- (A) S only (B) P and R only (C) Q and S only (D) Q only


Q.32 Match the entries in **Group I** with the enzymes in **Group II**.

<u>Group I</u>	<u>Group II</u>
P. NAD+	1. Glutathione peroxidase
Q. Selenium	2. Nitrogenase
R. Pyridoxal phosphate	3. Lactate dehydrogenase
S. Molybdenum	4. Glycogen phosphorylase
(A) P-3, Q-2, R-4, S-1	(B) P-4, Q-1, R-3, S-2
(C) P-3, Q-1, R-4, S-2	(D) P-3, Q-4, R-2, S-1

Match the herbicides in **Group I** with the target enzymesin **Group II**. Q.33

<u>Group I</u>	<u>Group II</u>
P. Glyphosate	1. Nitrilase
Q. Bromoxynil	2. Acetolactatesynthetase
R. Sulphonylureas	3. Dehalogenase





Q.42 A complete restriction digestion of a circular plasmid (5000bp) was carried out with *HindIII,Bam*HIand*Eco*RIindividually. Restriction digestion yielded following fragments.

Plasmid + $HindIII \rightarrow 1200$ bp and 3800bp Plasmid + $BamHI \rightarrow 5000$ bp Plasmid + $EcoRI \rightarrow 2500$ bp

The number of sites for EcoRI,BamHIandHindIIIpresent on this plasmid are(A) EcoRI-2, BamHI-1, HindIII-2(B) EcoRI-1, BamHI-1, HindIII-2(C)EcoRI-3, BamHI-2, HindIII-1(D) EcoRI-2, BamHI-2, HindIII-1

Q. 43 – Q. 47 are of numerical answer type.

Q.43 The total number of fragments generated by the complete and sequential cleavage of the polypeptide given below by Trypsin followed by CNBr is ______. Phe-Trp-Met-Gly-Ala-Lys-Leu-Pro-Met-Asp-Gly-Arg-Cys-Ala-Gln

BT-GATE-2013 PAPER

- Q.44 In a genetic study, 80 people were found to have alleles for polydactyly. Only 36 were polydactylous. What is the extent of penetrance percentage?
- StudentBounty.com Q.45 One percent of the cars manufactured by a company are defective. What is the probability (upto four decimals) that more than two cars are defective, if 100 cars are produced?
- The maximum cell concentration (q I-1) expected in a bioreactor with initial cell Q.46 concentration of

1.75 g l-1 and an initial glucose concentration of 125 g l-1 is (Yx/s = 0.6 g cell/g)substrate) _____

Q.47 A fed batch culture was operated with intermittent addition of glucose solution at a flow rate of

200 ml h-1. The values of Ks, K_s , μ_m and D are 0.3 g l⁻¹, 0.4 h⁻¹ and 0.1 h⁻¹, respectively. Determine the concentration of growth limiting substrate (gl-1) in the reactor at quasisteady state.

Common Data for Questions 48 and 49:


A solution was prepared by dissolving 100 mg of protein X in 100 ml of water. Molecular weight of protein X is 15,000 Da; Avogadro's number = 6.022x 1023.

- Q.48 Calculate the molarity (μ M) of the resulting solution. (B) 6.6 (A) 66.6 (C) 0.67 (D) 0.067
- Q.49 The number of moleculespresent in this solution is (A) 40.15x1019 (B) 6.023x1019 (C) 4.015x1019 (D) 0.08x1019

Common Data for Questions 50 and 51:

The binding efficiency of three different receptorsR1, R2 and R3 were tested against a ligand using equilibrium dialysis, with a constant concentration of receptor and varying concentrations of ligand. The Scatchard plot of receptor titration with different concentration of ligand is given below

(ris moles of bound ligand per moles of receptor and c is concentration of free ligand)




Q.50 The number of ligand binding sites present on receptors R1 and R3, respectively are (A)1 and 4 (B)1 and 1 (C) 4 and 1 (D) 2 and 2

ら罪	TEFORUM pineering Success	BT-G	ATE-2013 PAPER	k www. Hildry com
Q.51	Which one of th	e receptors has the	highestaffinity for th	ne ligand?
	(A) R1	(B) R2	(C) R3	(D) R4
	S	tatement for Link	ed Answer Questic	R www. he ligand? (D) R4 ons 52 and 53: coli (genome size 4x103kb) genomic
	A DNA fragment library.	of 5000bp needs to	o be isolated from <i>E</i> .	<i>coli</i> (genome size 4x103kb) genomic
Q.52	The minimum fragment in genomic library		lent recombinant clo	nes required to represent this
	(A) 16x102		(C) 8x102	(D) 1.25x102
2.53	The number of 95% are	clones to represent	this fragment in ger	nomic library with a probability of
	(A) 5.9 x103	(B) 4.5x103	(C) 3.6 x103	(D) 2.4x103
	During sterilizat	ion of a fermentatio = 12.56, $\nabla_{\text{cooling}} = 7$.	ed Answer Questic on medium in a giver 48 and the total val is the design criteri	ue of ∇ required for whole
2.54	What is the valu (A) 31.96	ue of ∇ holding? (B) 42.32	(C) 52.43	(D) 61.18
9.55			a k value of 3.36min (C) 8.4	
			ptitude (GA) Ques 60 carry one mark	
	If $3 \le X \le 5$ and	$8 \le Y \le 11$ then w	hich of the following	options is TRUE?
Q.56			2 V E	
Q.56	$(A) \frac{3}{5} \le \frac{X}{Y} \le \frac{8}{5}$	(B) $\frac{3}{11} \le \frac{X}{Y} \le \frac{5}{8}$	

Q.57 The Headmaster ______ to speak to you.
Which of the following options is incorrect to complete the above sentence?
(A) is wanting
(B) wants
(C) want
(D) was wanting

Q.58 Mahatama Gandhi was known for his humility as(A) he played an important role in humiliating exit of British from India.



Country	Number of
	Tourists
USA	2000
England	3500
Germany	1200
Italy	1100
Japan	2400
Australia	2300

Which two countries contributed to the one third of the total number of tourists who visited India in

1000

2011?

	(A) USA and Japan	(B)	USA and Australia		
	(C)England and France		(D) Japan and Australia		
Q.64	4 If $ -2X+9 = 3$ then the possible value of $ -2X+9 = 3$ would be:				
	(A) 30	(B) -30	(C) -42	(D) 42	

France

EGATEFORUM

|BT-GATE-2013 PAPER|

Q.65 All professors are researchers

Some scientists are professors

www. www. ants. www. Stilldenter ants. Comm Which of the given conclusions is logically valid and is inferred from the above arguments

- (A) All scientists are researchers
- (C) Some researchers are scientists
- (B) All professors are scientists
- (D) No conclusion follows

