

**Duration: Three Hours**

**Maximum Marks: 150**

➤ **Read the following instructions carefully. ]**

1. This question paper contains all objective questions. Q.1 to 20 carry one mark each and Q. 21 to Q. 85 carry two marks each.
2. Answer all the questions.
3. Questions must be answered on Objective Response Sheet (ORS) by darkening the appropriate bubble (marked A, B,C,D) using HB pencil against the question number on the left hand side of the ORS. Each question has only one correct answer. In case you wish to change an answer erase the old answer completely.
4. Wrong answers will carry Negative marks. In Q. 1 to Q. 20, 0.25 mark will be deducted for each wrong answer. In Q. 21 to Q. 76, Q. 78, Q.80, Q82 and in Q.84, 0.5 mark will be deducted for each wrong answer. However, there is no negative marking in Q. 77, Q.79, Q.81, Q.83 and in Q. 85. More than one answer bubbled against a question will be taken as an incorrect response.
5. Write your registration number, your name and name of the examination centre at the specified locations on the right half of the ORS.
6. Using HB pencil, darken the appropriate bubble under each digit of your registration number and the letters corresponding to your paper code.
7. Calculator is allowed in the examination hall.
8. Charts, graph sheets or tables are Not allowed in the examination hall.
9. Rough work can be done on the question paper itself. Additionally bank pages are given at the end of the question paper for rough work.
10. This question paper contains 24 printed pages including pages for rough work. Please check all pages and report, if there is any discrepancy.

## ONE MARKS QUESTIONS (1-20)

1. The dimension of the subspace  $\{(x_1, x_2, x_3, x_4, x_5) : 3x_1 - x_2 + x_3 = 0\}$  of  $\mathbb{R}^5$  is
  - (a.) 1
  - (b.) 2
  - (c.) 3
  - (d.) 4

$$T(x, y, z) = (x \cos \theta - y \sin \theta, \sin \theta, z)$$

here  $0 < \theta < \pi/2$ . Then

- (a.) S is one to one but not T
  - (b.) T is one to one but not S
  - (c.) Both S and T are one to one
  - (d.) Neither S nor T is one to one
3. Let E a non-measurable subset of  $[0,1]$ . If

$f : [0,1] \rightarrow \mathbb{R}$  is defined by

$$f(x) = \begin{cases} -1 & x \in E \\ 2 & \\ 0 & \text{Otherwise} \end{cases}$$

Then

- (a.)  $f$  is measurable but not  $|f|$
  - (b.)  $|f|$  is measurable but not  $f$
  - (c.) Both  $f$  and  $|f|$  are measurable
  - (d.) Neither  $f$  nor  $|f|$  is measurable
4. Let  $L^2([0,1])$  denote the space of all

square integrable functions on  $[0,1]$ .

Define  $f_1, f_2 : [0,1] \rightarrow \mathbb{R}$  by

$$f_1(t) = \begin{cases} t^{-1/3}, & 0 < t \leq 1 \\ 0, & t = 0 \end{cases} \quad f_2(t) = \begin{cases} t^{-2/3}, & 0 < t \leq 1 \\ 0, & t = 0 \end{cases}$$

Then,

- (a.)  $f_1$  belongs to  $L^2([0,1])$  but Not  $f_2$
  - (b.)  $f_2$  belongs to  $L^2([0,1])$  but Not  $f_1$
  - (c.) Both  $f_1$  and  $f_2$  belong to  $L^2([0,1])$
  - (d.) Neither  $f_1$  nor  $f_2$  belongs to  $L^2([0,1])$
5. For the ordinary differential equation

$$(x-1) \frac{d^2 y}{dx^2} + (\cot \pi x) \frac{dy}{dx} + (\cos e c^2 \pi x) y = 0$$

which of the following statements is true?

- (a.) 0 is regular and 1 is irregular
- (b.) 0 is irregular and 1 is regular
- (c.) Both 0 and 1 are regular
- (d.) Both 0 and 1 are irregular

(a.)  $\frac{1}{(n!2^n)}$

(b.)  $\frac{n!}{(2^n)}$

(c.)  $(n!)2^n$

(d.)  $\frac{2^n}{(n!)}$

7. Let  $G$  be a cyclic group of order 8, then its group of automorphisms has order

(a.) 2

(b.) 4

(c.) 6

(d.) 8

8. Let  $M_3(\mathbb{R})$  be the ring of all  $3 \times 3$  real matrices. If  $I, J \subseteq M_3(\mathbb{R})$  are defined as

$$I = \left\{ \begin{pmatrix} a & b & c \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \mid a, b, c \in \mathbb{R} \right\},$$

$$J = \left\{ \begin{pmatrix} a & 0 & 0 \\ b & 0 & 0 \\ c & 0 & 0 \end{pmatrix} \mid a, b, c \in \mathbb{R} \right\}$$

Then

(a.)  $I$  is a right ideal and  $J$  a left ideal

(b.)  $I$  and  $J$  are both left ideals

(c.)  $I$  and  $J$  are both right ideals

(d.)  $I$  is a left ideal and  $J$  a right ideal

9. Consider the Hilbert space

$$l^2 = \left\{ (x_1, x_2, \dots) \mid x_i \in \mathbb{R}, i = 1, 2, \dots \text{ and } \sum_{i=1}^{\infty} x_i^2 < \infty \right\}$$

under the inner product

$$\langle (x_1, x_2, \dots), (y_1, y_2, \dots) \rangle = \sum_{i=1}^{\infty} x_i y_i.$$

$$\text{Let } S = \left\{ (x_1, x_2, \dots) \in l^2 \mid \sum_{n=1}^{\infty} \frac{x_n}{n} = 0 \right\}.$$
 Then

the number of interior points of  $S$  is

(a.) 0

(b.) Non zero by finite

(c.) Count ably infinite

(d.) Un count ably infinite

10. Let  $C([0,1])$  be the space of all real valued continuous functions on  $[0,1]$  with

$n$  and the  $P([0,1])$  of all polynomials on  $[0,1]$

(a.)  $P_n([0,1])$  is closed in  $C([0,1])$  but not

$$P([0,1])$$

(b.)  $P([0,1])$  is closed in  $C([0,1])$  but not

$$P_n([0,1])$$

(c.) Both  $P([0,1])$  and  $P_n([0,1])$

closed in  $C([0,1])$

(d.) Neither  $P([0,1])$  nor  $P_n([0,1])$  is

closed in  $C([0,1])$

11. In the region  $x > 0, y > 0$ , the partial differential equation

$$(x^2 - y^2) \frac{\partial^2 u}{\partial x^2} + 2(x^2 + y^2) \frac{\partial^2 u}{\partial x \partial y} + (x^2 - y^2) \frac{\partial^2 u}{\partial y^2} = 0$$

(a.) Changes type

(b.) Is elliptic

(c.) Is parabolic

(d.) Is hyperbolic

12. Consider the partial differential equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0$$
 satisfying the initial

condition  $u(x, 0) = \alpha + \beta x$ . If  $u(x, t) = 1$  along the characteristic  $x = t + 1$ , then

(a.)  $\alpha = 1, \beta = 1$

(b.)  $\alpha = 2, \beta = 0$

(c.)  $\alpha = 0, \beta = 0$

(d.)  $\alpha = 0, \beta = 1$

13. Consider the usual topology on  $\mathbb{R}$ . Let  $S = \{U \subseteq \mathbb{R} : U \text{ is either bounded open or empty or } \mathbb{R}\}$  and  $T = \{U \subseteq \mathbb{R} : U \text{ is either unbounded open or empty or } \mathbb{R}\}$ .

Then, on  $\mathbb{R}$

(a.)  $S$  is a topology but  $T$  is not

(b.)  $T$  is a topology but  $S$  is not

(c.) Both  $S$  and  $T$  are topologies

(d.) Neither  $S$  nor  $T$  is a topology

14. Let  $X, Y$  and  $Z$  be events which are

(a.)  $ab + bc + ca - abc$

(b.)  $ab + bc + ca - 3abc$

(c.)  $2(a + b + c) - abc$

(d.)  $ab + bc + ca$

15. Assume that 45 percent of the population favours a certain candidate in an election. If a random sample of size 200 is chosen, then the standard deviation of the number of members of the sample that favours the candidate is

(a.) 6.12

(b.) 5.26

(c.) 8.18

(d.) 7.04

16. Let X and Y be independent Poisson random variables with parameters 1 and 2 respectively.

Then, P is  $\left( X = 1 \mid \frac{X + Y}{2} = 2 \right)$

(a.) 0.426

(b.) 0.293

(c.) 0.395

(d.) 0.512

17. For a linear programming primal maximization problem P with dual Q, which of the following statements is correct?

(a.) The optimal values of P and Q exist and are the same

(b.) Both optimal values exist and the optimal value of P is less than the optimal value of Q

(c.) P will have an optimal solution, if and only if Q also has an optimal solution

(d.) Both P and Q cannot be infeasible

18. Let a convex set in 9-dimensional space be given by the solution set of the following system of linear inequalities

$$\sum_{j=1}^3 x_{ij} = 1, \quad i = 1, 2, 3$$

$$\sum_{j=1}^3 x_{ij} = 1, \quad j = 1, 2, 3$$

$$x_{ij} \geq 0, \quad i, j = 1, 2, 3$$

Then, the number of extreme points of this set is

(a.) 3

$$I(y(x)) = \int_0^{\pi/2} y(x) dx; y(0) = 0$$

$$y(\pi/2) = 1$$

Where the unknown  $y(x)$  possesses two derivatives in  $(0, \pi/2)$ . Then

(a.) The functional has an extremum which can not be achieved in the class of continuous functions

(b.) The corresponding Euler's equation does not have a unique solution satisfying the given boundary conditions

(c.) I is not linear

(d.) I is linear

20. Solution of the initial value problem

$$\frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_2(x) y = F(x), \quad 0 \leq x \leq 1$$

$$y(0) = c_0, \quad \left( \frac{dy}{dx} = x \right)_{x=0} = c_1$$

Where  $a_1(x), a_2(x)$  and  $F(x)$  are continuous functions on  $\{0, 1\}$ , may be reduced, in general to a solution of some linear

(a.) Fredholm integral equation of first kind

(b.) Volterra's integral equation of first kind

(c.) Fredholm integral equation of second kind

(d.) Volterra's integral equation of second kind

## TWO MARKS QUESTIONS (21-75)

21. Let V be the vector space of all real polynomials. Consider the subspace W spanned by

$$t^2 + t + 2, \quad t^2 + 2t + 5, \quad 5t^2 + 3t + 4 \quad \text{and} \quad 2t^2 + 2t + 4.$$

Then the dimension of W is

(a.) 4

(b.) 3

(c.) 2

(d.) 1

$$\langle f, g \rangle = \int_0^1 f(x)g(x)dx \quad \text{and}$$

$V = \text{span}\{t^2\}$ . Let  $h(t) \in V$  be such that  $\|(2t-1)-h(t)\| \leq \|(2t-1)-x(t)\|$  for  $x(t) \in V$ . Then,  $h(t)$  is

- (a.)  $\frac{5}{6}t^2$
- (b.)  $\frac{5}{3}t^2$
- (c.)  $\frac{5}{12}t^2$
- (d.)  $\frac{5}{24}t^2$

23. Let  $M = \begin{pmatrix} 1 & a & b \\ 0 & 2 & c \\ 0 & 0 & 1 \end{pmatrix}$ ,  $a, b, c \in R$ .

Then, M is diagonalizable, if and only if

- (a.)  $a = bc$
- (b.)  $b = ac$
- (c.)  $c = ab$
- (d.)  $a = b = c$

24. Let M be the real  $5 \times 5$  matrix having all of its entries equal to 1. Then,

- (a.) M is not diagonalizable
- (b.) M is idempotent
- (c.) M is nilpotent
- (d.) The minimal polynomial and the characteristic polynomial of M are not equal

25. Let  $\{v_1, v_2, \dots, v_{16}\}$  be an ordered basis for  $V = C^{16}$ . If T is a linear transformation on V defined by

$$T(v_i) = v_i + 1 \quad \text{for } 1 \leq i \leq 15 \quad \text{and} \\ T(v_{16}) = -(v_1 + v_2 + \dots + v_{16}).$$

Then,

- (a.) R is singular with rational eigen values
- (b.) T is singular but has no rational eigen values
- (c.) T is regular (invertible) with rational eigen values
- (d.) T is regular but has no rational eigen values

$$\frac{2\pi}{\pi}$$

(c.)  $\pi$

(d.)  $i\pi$

27. The sum of the poles of  $f(z) = \frac{\cot \pi z}{(z+a)^2}$ , where  $a$  is a constant,

$$f(z) = \frac{\cot \pi z}{(z+a)^2}, \quad \text{where } a \text{ is a constant,}$$

( $a \neq 0, \pm 1, \pm 2, \dots$ ) is

(a.)  $\frac{1}{\pi} \sum_{n=-\infty}^{\infty} \frac{1}{(n+a)^2} - \pi \cos ec^2 \pi a$

(b.)  $-\frac{1}{\pi} \sum_{n=-\infty}^{\infty} \frac{1}{(n+a)^2} - \pi \cos ec^2 \pi a$

(c.)  $\frac{1}{\pi} \sum_{n=-\infty}^{\infty} \frac{1}{(n+a)^2} - \pi \cos ec^2 \pi a$

(d.)  $\frac{1}{\pi} \sum_{n=-\infty}^{\infty} \frac{1}{(n+a)^2} - \pi \cos ec^2 \pi a$

28. Let  $f(z)$  be an entire function such that for some constant,  $\alpha, |f(z)| \leq \alpha |z|^3$  for  $|z| \geq 1$  and  $f(z) = f(iz)$  for all  $z \in C$ . Then

(a.)  $f(z) = \alpha z^3$  for all  $z \in C$

(b.)  $f(z)$  is a constant

(c.)  $f(z)$  is a quadratic polynomial

(d.) No such  $f(z)$  exists

29. Which of the following is not the real part of an analytic function?

(a.)  $x^2 - y^2$

(b.)  $\frac{1}{1+x^2+y^2}$

(c.)  $\cos x \cos hy$

(d.)  $x + \frac{x}{x^2 + y^2}$

30. The radius of convergence of

$$\sum_{n=0}^{\infty} \left(1 + \frac{1}{n}\right)^{n^2} z^n$$

(a.)  $e$

(b.)  $1/e$

(c.)  $1$

(d.)  $\infty$

31. Let,  $S, T \subseteq R^2$  be given by

$$S = \left\{ \left( s, \sin \frac{1}{2} \right) : 0 < x \leq 1 \right\} \cup \{(0, 0)\} \quad \text{and}$$

$$T = \left\{ \left( x, \sin \frac{1}{2} \right) : 0 < x \leq 1 \right\} \cup \{(0, 0)\}.$$

Then, under the usual metric on  $\mathbb{R}^2$ ,

- (a.) S is compact but not T
- (b.) T is compact but not S
- (c.) Both S and T are compact
- (d.) Neither S nor T is compact

32. Let  $S, T \subseteq \mathbb{R}$  be given by

$$S = \left\{ x \in \mathbb{R} : 2x^2 \cos \frac{1}{x} = 1 \right\} \quad \text{and}$$

$$T = \left\{ x \in \mathbb{R} : 2x^2 \cos \frac{1}{x} \leq 1 \right\} \cup \{0\}. \quad \text{Then,}$$

under the usual metric on  $\mathbb{R}$ ,

- (a.) S is complete but not T
- (b.) T is complete but not S
- (c.) Both S and T are complete
- (d.) Neither S nor T is complete

33. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \begin{cases} n, & \text{If } x = n, \quad x \in \mathbb{N} \\ 0, & \text{Otherwise} \end{cases} \quad \text{and}$$

$$T = \mathbb{N} \cup \left\{ n + \frac{1}{n} : n \in \mathbb{N} \right\}. \quad \text{Then, under the}$$

usual metric on  $\mathbb{R}$ ,  $f$  is uniformly continuous on

- (a.)  $\mathbb{N}$  but Not T
- (b.) T but not  $\mathbb{N}$
- (c.) Both  $\mathbb{N}$  and T
- (d.) Neither  $\mathbb{N}$  nor T

34. For each  $n \in \mathbb{N}$  and  $n > 1$  define

$$f_n: [0, 1] \rightarrow \mathbb{R} \text{ by}$$

$$f_n(x) = \begin{cases} |nx - 1| & \text{for } 0 \leq x < \frac{2}{n} \\ 1 & \text{for } \frac{2}{n} \leq x \leq 1 \end{cases}$$

Let  $g_1, g_2: [0, 1] \rightarrow \mathbb{R}$  be defined by

$$g_1(x) = \begin{cases} 1 & \text{for } 0 < x \leq 1 \\ 0 & \text{for } x = 0 \end{cases} \quad \text{and} \quad g_2(x) = 1$$

for  $0 \leq x \leq 1$ .

Then, on  $[0, 1]$

- (a.)  $f_n \rightarrow g_1$  point wise but not uniformly

35. Let  $f_n, g_n$  be defined by

$$f_n(x) = x^2(1 - \frac{1}{n}) \quad \text{and}$$

$$g_n(x) = \frac{1}{n^2(1+x^2)}$$

Then, on  $[0, 1]$

- (a.)  $\sum_{n=1}^{\infty} f_n(x)$  converges uniformly

$$\sum_{n=1}^{\infty} g_n(x)$$

- (b.)  $\sum_{n=1}^{\infty} g_n(x)$  converges uniformly but not

$$\sum_{n=1}^{\infty} f_n(x)$$

- (c.) Both  $\sum_{n=1}^{\infty} f_n(x)$  and  $\sum_{n=1}^{\infty} g_n(x)$  converge uniformly

- (d.) Neither  $\sum_{n=1}^{\infty} f_n(x)$  nor  $\sum_{n=1}^{\infty} g_n(x)$  converges uniformly

36. The function  $f: [0, \infty] \rightarrow \mathbb{R}$  defined by

$$f(x) = \int_0^x (2 \sin^4 t \cos^2 t) dt \text{ is}$$

- (a.) Not continuous
- (b.) Continuous but not uniformly
- (c.) Uniformly continuous but not Lipschitz continuous
- (d.) Lipschitz continuous

37. Let S be a non-measurable subset of  $\mathbb{R}$  and T be measurable subset of  $\mathbb{R}$  such that  $S \subset T$ . Denote the outer measure of a set U by  $m^*(U)$ . Then,

- (a.)  $m^*(T/S) = 0$  and  $m^*(S) = 0$
- (b.)  $m^*(T/S) > 0$  and  $m^*(S) > 0$
- (c.)  $m^*(T/S) > 0$  and  $m^*(S) = 0$
- (d.)  $m^*(T/S) = 0$  and  $m^*(S) > 0$

38. Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined by

$$f(x, y) = \begin{cases} \frac{x^2 y}{x^4 + y^2} & \text{for } (x, y) \neq (0, 0) \\ 0 & \text{for } (x, y) = (0, 0) \end{cases}$$

Then, the directional derivative of  $f$  at  $(0,0)$  in the direction of the vector

$$\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) \text{ is}$$

- (a.)  $\frac{1}{\sqrt{2}}$
- (b.)  $\frac{1}{2}$
- (c.)  $\frac{1}{2\sqrt{2}}$
- (d.)  $\frac{1}{4\sqrt{2}}$

39. Consider the hemisphere  $x^2 + y^2 + (z-2)^2 = 9$ ,  $2 \leq z \leq 5$  and the vector field

$$\vec{F}(x, y, z) = x\vec{i} + y\vec{j} + (z-2)\vec{k}.$$

The surface integral  $\iint (\vec{F} \cdot \vec{n}) d\sigma$ , evaluated over the hemisphere with  $\vec{n}$  denoting the unit outward normal is

- (a.)  $9\pi$
- (b.)  $27\pi$
- (c.)  $54\pi$
- (d.)  $162\pi$

40. Let  $y_1(x)$  and  $y_2(x)$  be two solutions of

$$(1-x^2)\frac{d^2y}{dx^2} - 2x\frac{dy}{dx} + (\sec x)y = 0$$

With Wronskian  $W(x)$ . If  $y_1(0) = 1$ ,

$$\left(\frac{dy_1}{dx}\right)_{x=0} = 0 \text{ and } W\left(\frac{1}{2}\right) = \frac{1}{3}, \text{ then}$$

$$\left(\frac{dy_2}{dx}\right)_{x=0} \text{ equals}$$

- (a.)  $1/4$
- (b.)  $1$
- (c.)  $3/4$
- (d.)  $4/3$

41. If  $y(x)$  is the solution of the differential

$$\text{equation } \frac{dy}{dx} = 2(1+y)\sqrt{y} \text{ satisfying}$$

$y(0) = 0$ ;  $y(\pi/2) = 1$ , then the largest interval (to the right of origin) on which the solution exists is

42.

(d.)  $[0, 2\pi]$  A particular solution of

$$x^2 \frac{d^2y}{dx^2} + 2x \frac{dy}{dx} + \frac{y}{4}$$

- (a.)  $\frac{1}{2\sqrt{x}}$
- (b.)  $\frac{\log x}{2\sqrt{x}}$
- (c.)  $\frac{(\log x)^2}{2\sqrt{x}}$
- (d.)  $\frac{(\log x)\sqrt{x}}{2}$

43.

The initial value problem

$$x \frac{d^2y}{dx^2} + \frac{dy}{dx} + xy = 0; y(0) = 1, \left(\frac{dy}{dx}\right)_{x=0} = 0$$

has

- (a.) A unique solution
- (b.) No solution
- (c.) Infinitely many solutions
- (d.) Two linearly independent solutions

44.

An integrating factor for

$$(\cos y \sin 2x)dx + (\cos^2 y - \cos^2 x)dy = 0$$

is

- (a.)  $\sec^2 y + \sec y \tan y$
- (b.)  $\tan^2 y + \sec y \tan y$
- (c.)  $1/(\sec^2 y + \sec y \tan y)$
- (d.)  $1/(\tan^2 y + \sec y \tan y)$

45.

Let  $F_4$ ,  $F_8$  and  $F_{16}$  be finite fields of 4, 8 and 16 elements respectively. Then,

- (a.)  $F_4$  is isomorphic to a subfield of  $F_8$
- (b.)  $F_9$  is isomorphic to a subfield of  $F_{16}$
- (c.)  $F_4$  is isomorphic to a subfield of  $F_{16}$
- (d.) None of the above

46.

Let  $G$  be the group with the generators  $a$  and  $b$  given by

$$G = \langle a, b : a^4 = b^2 = 1, ba = a^{-1}b \rangle.$$

If  $Z(G)$  denotes the centre of  $G$ , then

$G/Z(G)$  is isomorphic to

- (a.) The trivial group
- (b.)  $C_2$ , the cyclic group of order 2
- (c.)  $C_2 \times C_2$
- (d.)  $C_4$



- (a.) F is an infinite field  
 (b.) F is a finite field of 4 elements  
 (c.) F is a finite field of 8 elements  
 (d.) F is a finite field of 16 elements
48. Let  $f, g: R/\{0,1\} \rightarrow R/\{0,1\}$  be defined by  $f(x) = 1/(1-x)$  and  $g(x) = x/(x-1)$ , and let G be the group generated by  $f$  and  $g$  under composition of mappings. It is given that G has order 6. Then,  
 (a.) G and its automorphisms group are both Abelian  
 (b.) G and its automorphisms group are both non-Abelian  
 (c.) G is abelian but its automorphisms group is non-abelian  
 (d.) G is non-abelian but its automorphisms group is Abelian
49. Let  $R = \{\alpha_0 + \alpha_1 i + \alpha_2 j + \alpha_3 k : \alpha_0, \alpha_1, \alpha_2, \alpha_3 \in \mathbb{Z}_3\}$  be the ring of quaternions over  $\mathbb{Z}_3$ , where  $i^2 = j^2 = k^2 = ijk = -1; ij = -ji = k; ki = -ik = j$ . Then  
 (a.) R is field  
 (b.) R is a division ring  
 (c.) R has zero divisors  
 (d.) None of the above
50. Consider the sequence of continuous linear operators  $T_n: l^2 \rightarrow l^2$  defined by  $T_n(x) = (0, 0, \dots, 0, x_{n+1}, x_{n+2}, x_{n+3}, \dots)$  for every  $x = (x_1, x_2, \dots) \in l^2$  and  $n \in \mathbb{N}$ . Then, for every  $x \neq 0$  in  $l^2$   
 (a.) Both  $\|T_n\|$  and  $\|T_n(x)\|$  converge to 0  
 (b.) Neither  $\|T_n\|$  nor  $\|T_n(x)\|$  converges to 0  
 (c.)  $\|T_n\|$  converges to 0 but not  $\|T_n(x)\|$   
 (d.)  $\|T_n(x)\|$  converges to 0 but not  $\|T_n\|$
51. Let the continuous linear operator  $T: l^2 \rightarrow l^2$  defined by  $T(x_1, x_2, \dots) = (0, x_1, 0, x_3, 0, x_5, 0, \dots)$ . Then  
 (a.) T is compact but not  $T^2$   
 (b.)  $T^2$  is compact but not T
52. Let  $f(x)$  be a real-valued function such that  $\frac{d^3 f}{dx^3} = 1$  for all  $x$ . If  $p(x)$  is the quadratic polynomial which interpolates  $f(x)$  at  $x = 1, 2$  and  $x = 3$ , then  $f(1) - p(1)$  equals  
 (a.) 0  
 (b.)  $1/3$   
 (c.)  $1/6$   
 (d.)  $2/3$
53. Let  $h(x)$  be a twice continuously differentiable function on  $[1, 2]$  with fixed point  $\alpha$ . Then, the sequence of iterates  $x_{n+1} = h(x_n)$  converges to  $\alpha$  quadratically, provided  
 (a.)  $\frac{dh}{dx}(\alpha) \neq 0$   
 (b.)  $\frac{dh}{dx}(\alpha) \neq 0, \frac{d^2 h}{dx^2}(\alpha) = 0$   
 (c.)  $\frac{dh}{dx}(\alpha) \neq 0, \frac{d^2 h}{dx^2}(\alpha) \neq 0$   
 (d.)  $\frac{dh}{dx}(\alpha) = 0, \frac{d^2 h}{dx^2}(\alpha) \neq 0$
54. Consider the initial value problem (IVP):  $\frac{dy}{dx} = f(x, y(x)), y(x_0) = y_0$ . Let  $y_1 = y_0 w_1 k_1 + 3k_1$  approximate the solution of the above IVP at  $x_1 = x_0 + h$  with  $k_1 = hf(x_0, y_0), k_2 = hf(x_0 + (h/6), y_0 + (k_1/5))$  and  $h$  being the step-size. If the formula for  $y_1$  yields a second order method, then the value of  $w_1$  is  
 (a.) -1  
 (b.) -2  
 (c.) 3  
 (d.)  $1/6$
55. Let  $u(x, t)$  be the solution of the initial value problem  $\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} = 0; u(x, 0) = \sin x; \frac{\partial u}{\partial t}(x, 0) = 0$ .

- (b.)  $1/2$   
(c.)  $-1$   
(d.)  $1$
56. Let  $u(x, t)$  be the bounded of  $\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} = 0$  with  $u(x, 0) = \frac{e^{2x} - 1}{e^{2x} + 1}$ . Then  $\lim_{t \rightarrow +\infty} u(1, t)$  equals  
(a.)  $-1/2$   
(b.)  $1/2$   
(c.)  $-1$   
(d.)  $1$
57. Let  $u(x, y)$  be a solution of Laplace's equation on  $x^2 + y^2 \leq 1$ . If  $u(\cos \theta, \sin \theta) = \begin{cases} \sin \theta & \text{for } 0 \leq \theta \leq \pi \\ 0 & \text{for } \pi \leq \theta \leq 2\pi \end{cases}$   
Then  $u(0, 0)$  equals  
(a.)  $1/\pi$   
(b.)  $2/\pi$   
(c.)  $1/(2\pi)$   
(d.)  $\pi/2$
58. Let PQRS be a rectangle in the first quadrant whose adjacent sides PQ and QR have slopes 1 and  $-1$  respectively. If  $u(x, t)$  is a solution of  $\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial z^2} = 0$  and  $u(P) = 1, u(Q) = -1/2, u(R) = 1/2$ , then  $u(S)$  equals  
(a.) 2  
(b.) 1  
(c.)  $1/2$   
(d.)  $-1/2$
59. In the motion of a two-particle system, if the two particles are connected by a rigid weightless rod of constant length  $l$ , then the number of degree of freedom of the system is  
(a.) 2  
(b.) 3  
(c.) 5  
(d.) 6
60. A particle of unit mass moves in the xy-plane under the influence of a central force depending only on its distance from the origin. If  $(r, \theta)$  be the polar coordinates of

- (a.)  $\frac{1}{2}\dot{r}^2 - V$   
(b.)  $\frac{1}{2}(\dot{r}^2 + \dot{\theta}^2) + V$   
(c.)  $\frac{1}{2}(\dot{r}^2 + r\dot{\theta}) + V(r)$   
(d.)  $\frac{1}{2}(\dot{r}^2 + r^2\dot{\theta}^2) - V(r)$
61. Let  $q, \dot{q}$  and  $p$  denote respective generalized coordinates, and corresponding velocity and momenta of a one-dimensional system with the Hamiltonian  $H = \frac{1}{2}\left(p^2 - \frac{1}{q^2}\right)$ . Then the Lagrangian of the system is  
(a.)  $\frac{1}{2}\dot{q}^2 - \frac{1}{q}$   
(b.)  $\frac{1}{2}\dot{q}^2 + \frac{1}{q}$   
(c.)  $\frac{1}{2}\left(\dot{q}^2 + \frac{1}{q^2}\right)$   
(d.)  $\frac{1}{2}\left(\dot{q}^2 - \frac{1}{q^2}\right)$
62. Let  $\tau_1$  be the usual topology on  $R$ . Define another topology  $\tau_2$  on  $R$  by  $\tau_2 = \{U \subseteq R \mid U^c \text{ is either finite or empty or whole of } R\}$  where  $U^c$  denotes the component of  $U$  in  $R$ . If  $I: (R, \tau_1) \rightarrow (R, \tau_2)$  is the identity map, then  
(a.)  $I$  is continuous but not  $I^{-1}$   
(b.)  $I^{-1}$  is continuous but not  $I$   
(c.) Both  $I$  and  $I^{-1}$  are continuous  
(d.) Neither  $I$  nor  $I^{-1}$  continuous
63. Let  $\tau_1$  be the usual topology on  $R$ . Define another topology  $\tau_2$  on  $R$  by  $\tau_2 = \{U \subseteq R \mid U^c \text{ is either countable or empty or whole of } R\}$ . Then,  $Z$  is  
(a.) Closed in  $(R, \tau_1)$  but not in  $(R, \tau_2)$   
(b.) Closed in  $(R, \tau_2)$  but not in  $(R, \tau_1)$



64. Consider  $\mathbb{R}^2$  with the usual topology. The complement of  $N \times N$  is  
 (a.) Open but not connected  
 (b.) Connected but not open  
 (c.) Both open and connected  
 (d.) Neither open nor connected
65. Let  $T$  denote the number of times we have to roll a fair dice before each face appearing in the first six rolls. Then  $E(T|N=3)$  is  
 (a.) 9  
 (b.) 15  
 (c.) 16  
 (d.) 17
66. Let there be three types of light bulbs with lifetimes  $X$ ,  $Y$  and  $Z$  having exponential distributions with mean  $\theta$ ,  $2\theta$  and  $3\theta$  respectively. Then, the maximum likelihood estimator of  $\theta$  based on the observation  $X$ ,  $Y$  and  $Z$  is  
 (a.)  $(X + 2Y + 3Z) / 3$   
 (b.)  $3(X + 2Y + 3Z)$   
 (c.)  $\frac{1}{3} \left( X + \frac{Y}{2} + \frac{Z}{3} \right)$   
 (d.)  $\frac{1}{6} \left( X + \frac{Y}{2} + \frac{Z}{3} \right)$
67. Let  $Z$  be the vertical coordinate, between  $-1$  and  $1$ , of a point chosen uniformly at random on the surface of a unit sphere in  $\mathbb{R}^3$ . Then,  $P\left(-\frac{1}{2} \leq Z \leq \frac{1}{2}\right)$  is  
 (a.)  $5/6$   
 (b.)  $(\sqrt{3}) / 2$   
 (c.)  $3/4$   
 (d.)  $1/2$
68. Let the marks obtained in the half-yearly and final examinations in a large class have an approximately bivariate normal distribution with the following parameters
- |                     | Mean | Deviation |
|---------------------|------|-----------|
| Marks (half yearly) | 60   | 18        |
| Marks (final exam)  | 55   | 20        |
| Correlation :       | 0.75 |           |
- Then, estimate of the average final examination score of students who were above average on the half-yearly

- (d.)  $7/2$
69. Let  $V_1, V_2, \dots, V_5$  independent uniform  $(0, 1)$  and let  $V_{(1)} < V_{(2)} < \dots < V_{(5)}$  their order statistics. Then, for  $0 < x < y < 1$  the joint density  $f(x, y)$  of  $(V_{(2)}, V_{(4)})$  is  
 (a.)  $(5!)xy(1-x)(1-y)$   
 (b.)  $x(y-x)(1-y) / (5!)$   
 (c.)  $(5!)x(y-x)(1-y)$   
 (d.)  $xy(1-x)(1-y) / (5!)$
70. Consider the linear programming problem  

$$\max c_1x_1 + c_2x_2 + c_3x_3$$

$$s.t \quad x_1 + x_2 + x_3 \leq 4$$

$$x_1 \leq 2$$

$$x_3 \leq 3$$

$$3x_1 + x_3 \leq 7$$

$$x_1, x_2, x_3 \geq 0.$$
 If  $(1, 0, 3)$  optimal solution, then  
 (a.)  $c_1 \leq c_2 \leq c_3$   
 (b.)  $c_3 \leq c_1 \leq c_2$   
 (c.)  $c_2 \leq c_3 \leq c_1$   
 (d.)  $c_2 \leq c_1 \leq c_3$
71. Let the convex set  $S$  be given by the solution set of the following system of linear inequalities in the sixteen variables  $\{x_{ij} : i, j = 1, \dots, 4\}$ .  

$$\sum_{j=1}^4 x_{ij} = 3, \quad i = 1, \dots, 4$$

$$\sum_{i=1}^4 x_{ij} = 3, \quad j = 1, \dots, 4$$

$$x_{ij} \geq 0, \quad i, j = 1, \dots, 4$$
 Then, the dimension of  $S$  is equal to  
 (a.) 4  
 (b.) 9  
 (c.) 8  
 (d.) 12
72. Let  $I(y(x)) = \int_0^1 F\left(x, y, \frac{dy}{dx}\right) dx$ , satisfying  
 $y(0) = 0, y(1) = 1$

two derivatives every where in  $(0,1)$ . If the function  $F$  depends only on  $x$  and  $\frac{dy}{dx}$ , then the Euler's equation is an ordinary differential equation in  $y$  which, in general, is

- (a.) First order linear
- (b.) First order nonlinear
- (c.) Second order linear
- (d.) Second order nonlinear

73. The functional

$$I(y(x)) = \int_0^1 \left( y + \frac{d^2 y}{dx^2} \right) dx$$

Defined on the set of functions  $C^2([0,1])$  satisfying

$$y(0) = 1, \quad y(1) = 1, \quad \left( \frac{dy}{dx} \right)_{x=0} = 0 \quad \text{and}$$

$$\left( \frac{dy}{dx} \right)_{x=1} = -1$$

- (a.) Only one extremal
- (b.) Exactly two extremals
- (c.) Infinite number of extremals
- (d.) No extremals

74. Which of the following functions is a solution of the Volterra type integral equation

$$f(x) = x + \int_0^x (\sin(x-t) f(t)) dt$$

- (a.)  $x + \frac{x^3}{3}$
- (b.)  $x - \frac{x^3}{3}$
- (c.)  $x + \frac{x^3}{6}$
- (d.)  $x - \frac{x^3}{6}$

75. Which of the following functions is a solution of the Fredholm type equation

$$f(x) = x + \int_0^1 (xtf(t)) dt$$

- (a.)  $2x/3$
- (b.)  $3x/2$
- (c.)  $3x/4$

**Statement for Linked Answer Questions 76 & 77:**

Let  $T: C^3 \rightarrow C^3$  be defined by  $T(x_1, x_2, x_3) = (x_1 + x_2 + x_3, x_1 - x_2 - x_3, x_1 + x_2 - x_3)$  and  $M$  be its matrix with respect to standard ordered basis.

76. The eigen values of  $M$  are

- (a.)  $-1, i, -i$
- (b.)  $1, i, -i$
- (c.)  $1, i, i$
- (d.)  $-1, -i, -i$

77. The matrix  $M$  is similar to a matrix which is

- (a.) Unitary
- (b.) Hermitian
- (c.) Skew Hermitian
- (d.) Having trace 0

**Statement for Linked Answer Questions 78 & 79:**

Let  $H$  be an infinite dimensional Hilbert space and  $f$  be a continuous linear functional on  $H$  such that  $\|f\| = 1$ . Define  $W = \{x \in H : f(x) = 1\}$ . Then interior and the boundary of the closed unit ball  $U$  of  $H$  are denoted by  $U^\circ$  and  $\partial U$  respectively.

78. Which of the following is correct?

- (a.)  $U^\circ \cap W = \phi$  and  $W \cap \partial U = \phi$
- (b.)  $U^\circ \cap W \neq \phi$  and  $W \cap \partial U = \phi$
- (c.)  $U^\circ \cap W = \phi$  and  $W \cap \partial U \neq \phi$
- (d.)  $U^\circ \cap W \neq \phi$  and  $W \cap \partial U \neq \phi$

79. The number of points in  $W \cap U$  is

- (a.) 0
- (b.) 1
- (c.) Not one but countable
- (d.) Uncountable

**Statement for Linked Answer Questions 80 and 81:**

Consider the partial differential equation

$$x \frac{\partial u}{\partial y} - y \frac{\partial u}{\partial x} = u$$

80. The characteristic curves for the above equation in the  $(x, y)$  plane are

- (a.) Straight line with slopes 1
- (b.) Straight lines with slopes  $-1$
- (c.) Circles with centre at the origin

81. If  $u(x, y)$  is a solution to the above equation with  $u(x, 0) = \sin\left(\frac{\pi}{4}x\right)$ , then

$u\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$  equals

(a.)  $\frac{1}{\sqrt{2}}e^{\frac{\pi}{4}}$

(b.)  $\frac{\pi}{4}e^{\frac{\pi}{\sqrt{2}}}$

(c.)  $\frac{1}{\sqrt{2}}e^{\frac{1}{\sqrt{2}}}$

(d.)  $\frac{\pi}{4}e^{\frac{\pi}{4}}$

**Statement for Linked Answer Questions 82 & 83:**

Let  $p(x) = c_0 + c_1x$  minimize

$$\langle f(x) - p(x), f(x) - p(x) \rangle = \int_{-1}^1 (f(x) - p(x))^2 dx$$

over all polynomials of degree less than or equal to 1.

82. The best choice of coefficients  $c_0, c_1$  is

(a.)  $\langle f, 1 \rangle, \langle f, x \rangle$

(b.)  $\langle f, 1 \rangle, \frac{2}{3}\langle f, x \rangle$

(c.)  $\frac{1}{2}\langle f, 1 \rangle, \frac{3}{2}\langle f, x \rangle$

(d.)  $\frac{2}{3}\langle f, 1 \rangle, \frac{2}{3}\langle f, x \rangle$

83. If  $f(x) = x^2 + x$ , then  $p(x)$  is given by

(a.)  $\left(1 + \frac{x}{3}\right)$

(b.)  $\frac{1}{3}(1 + 3x)$

(c.)  $\frac{1}{3}(1 + x)$

(d.)  $\frac{2}{3}(1 + x)$

**Statement for Linked Answer Questions**

**84 & 85:**

Consider the following Linear Programming Problem P:

$$\max \quad c_1x_1 + c_2x_2 + \dots$$

$$s.t. \quad \sum_{j=1}^n a_{ij}x_j \leq b_i, \quad i = 1, \dots, m$$

$$x_j \geq 0, \quad j = 1, \dots, n,$$

With  $m$  constants in  $n$  non-negative variables.

84. Let  $x^* = (x_1^*, x_2^*, \dots, x_n^*)$  be an optimal extreme point solution to P with  $x_1^*, x_2^*, x_3^*, \dots, x_n^* > 0$ . Then out of the  $m$

constraints  $\sum_{j=1}^n a_{ij}x_j \leq b_i, i = 1, \dots, m$  the

number of constraints not satisfied with equality at  $x^*$  is

(a.) At most  $m-4$

(b.) At most  $n-4$

(c.) Equal to  $m-3$

(d.) Equal to  $m-2$

85. Treat  $c_i$ 's,  $a_{ij}$ 's fixed and consider the problem P for different values of  $b_i$ 's. Let P be unbounded for some set of parameters  $b_1, b_2, \dots, b_m$ . Then

(a.)  $n > m$

(b.) P is either unbounded or infeasible every choice of  $b_i$ 's

(c.)  $m > n$

(d.) P has an optimal solution for some choice of  $b_i$ 's