



Physics Equations Sheet

GCSE Combined Science: Trilogy (8464) and GCSE Combined Science: Synergy (8465)

FOR USE IN JUNE 2022 ONLY

HT = Higher Tier only equations

| | |
|--|--------------------------------|
| kinetic energy = $0.5 \times \text{mass} \times (\text{speed})^2$ | $E_k = \frac{1}{2} m v^2$ |
| elastic potential energy = $0.5 \times \text{spring constant} \times (\text{extension})^2$ | $E_e = \frac{1}{2} k e^2$ |
| gravitational potential energy = $\text{mass} \times \text{gravitational field strength} \times \text{height}$ | $E_p = m g h$ |
| change in thermal energy = $\text{mass} \times \text{specific heat capacity} \times \text{temperature change}$ | $\Delta E = m c \Delta \theta$ |
| power = $\frac{\text{energy transferred}}{\text{time}}$ | $P = \frac{E}{t}$ |
| power = $\frac{\text{work done}}{\text{time}}$ | $P = \frac{W}{t}$ |
| efficiency = $\frac{\text{useful output energy transfer}}{\text{total input energy transfer}}$ | |
| efficiency = $\frac{\text{useful power output}}{\text{total power input}}$ | |
| charge flow = $\text{current} \times \text{time}$ | $Q = I t$ |
| potential difference = $\text{current} \times \text{resistance}$ | $V = I R$ |
| power = $\text{potential difference} \times \text{current}$ | $P = V I$ |
| power = $(\text{current})^2 \times \text{resistance}$ | $P = I^2 R$ |
| energy transferred = $\text{power} \times \text{time}$ | $E = P t$ |

| | | |
|----|--|--------------------------|
| | energy transferred = charge flow × potential difference | $E = QV$ |
| HT | potential difference across primary coil × current in primary coil = potential difference across secondary coil × current in secondary coil | $V_p I_p = V_s I_s$ |
| | density = $\frac{\text{mass}}{\text{volume}}$ | $\rho = \frac{m}{V}$ |
| | thermal energy for a change of state = mass × specific latent heat | $E = mL$ |
| | weight = mass × gravitational field strength | $W = mg$ |
| | work done = force × distance (along the line of action of the force) | $W = Fs$ |
| | force = spring constant × extension | $F = ke$ |
| | distance travelled = speed × time | $s = vt$ |
| | acceleration = $\frac{\text{change in velocity}}{\text{time taken}}$ | $a = \frac{\Delta v}{t}$ |
| | (final velocity) ² – (initial velocity) ² = 2 × acceleration × distance | $v^2 - u^2 = 2as$ |
| | resultant force = mass × acceleration | $F = ma$ |
| HT | momentum = mass × velocity | $p = mv$ |
| | period = $\frac{1}{\text{frequency}}$ | $T = \frac{1}{f}$ |
| | wave speed = frequency × wavelength | $v = f\lambda$ |
| HT | force on a conductor (at right angles to a magnetic field) carrying a current = magnetic flux density × current × length | $F = BIl$ |