

Edexcel GCSE

Mathematics B 1388 Paper 5510

November 2006

advancing learning, changing lives

Mark Scheme (Results)

Mathematics B 1388

NOTES ON MARKING PRINCIPLES

1 Types of mark

M marks: method marks A marks: accuracy marks

B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations

cao - correct answer only
ft - follow through
isw - ignore subsequent working
SC: special case
oe - or equivalent (and appropriate)
dep - dependent
indep - independent

3 No working

If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers
score no marks.

4 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader. If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work. If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.

Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct. It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.

If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

5510 - Section A					
N	lo	Working	Answer	Mark	Notes
1			120	3	M1 for $300 \div (2+3)$ or 60 seen
			180		M1 for "60" × 2 or "60" × 3
					A1 for 120 and 180
2	(a)		t 3	1	B1 cao
	(b)		$14f^5g^9$	2	B2 (B1 for f^5 or g^9)
3		$\frac{49}{15} \div \frac{21}{10}$	$\frac{14}{9}$ oe	3	M1 for one number written as an improper fraction correctly
		$=\frac{49}{15}\times\frac{10}{21}$			M1 for " $\frac{49}{15}$ " × " $\frac{10}{21}$ " oe
		$=\frac{7}{3}\times\frac{2}{3}$			A1 for $\frac{14}{9}$ or $1\frac{5}{9}$ oe (eg $\frac{490}{315}$)
4		6x + 13 = 4x - 12	$\frac{-25}{2}$ oe	3	B1 for $6x + 13 = 4x - 12$ or $\frac{6x}{4} + \frac{13}{4} = x - 3$
		6x - 4x = -12 - 13			M1 (ft from $ax + b = cx + d$ $a,b,c,d \neq 0$) for correct method to isolate the terms in x
		2x = -25			A1 for $\frac{-25}{2}$ oe
5	(a)		$7.2(300) \times 10^{-4}$	1	B1 cao
	(b)		0.0915	1	B1 cao
6			enlargement	2	B2 for correct enlargement
			(-4, -4) (-10, -4) (-10, -8)		(B1 for enlargement of -1 or 2 or $-\frac{1}{2}$, centre (0,0) or
					triangle of correct size and orientation or 2 out of 3 vertices correct $\pm \frac{1}{2}$ sq or coordinates listed correctly)
					(SC : B1 for (-4, -4) (-11, -4) (-11, -8))

5510 - Section A				
No	Working	Answer	Mark	Notes
7		Heights	3	B3 for fully correct histogram
		2, 4, 1		(B2 for 2 bars correct)
				(B1 for clear use of frequency density)

5510 - Section B					
	No	Working	Answer	Mark	Notes
1		8.1 × 20	162	2	M1 8.1 × 20
					A1 cao
2	(a)	(5), 0, (-3), -4, (-3), 0, 5	table	2	B2 (B1 for 2 or 3 correct)
	(b)		graph	2	B1 ft for all points correctly plotted
					B1 (ft if at least B1 in (a)) for points joined with smooth curve
3		5 120	5.3	4	B2 for trial between 5 and 6
		6 210			(B1 for trial at 5 or 6)
		5.2 135.4			B1 for different trial between 5.2 and 5.3 exclusive
		5.3 143.57			B1 (dep on at least one previous B1) for 5.3
		5.4 152.06			
		5.5 160.875			
		5.25 139.4			NB Trials should be evaluated to at least 3 sf, truncated or rounded
4	(a)		1.57	1	B1 cao
	(b)		1.4	1	B1 cao
	(c)		0.46	1	B1 cao

5510 - Section B				
No	Working	Answer	Mark	Notes
5	angle $ABD = 90^{\circ}$ $OB^{2} = 8.7^{2} - 6.8^{2}$ $OB = 5.426$ $\cos CBD = \frac{9.5}{2 \times "5.43"}$	28.9°	6	B1 for angle $ABD = 90^{\circ}$ (may be stated or marked on the diagram or may be implied by use of Pythagoras or trigonometry) M1 for $OB^2 + 6.8^2 = 8.7^2$ or $(OB^2 =) 8.7^2 - 6.8^2$ oe M1 for $\sqrt{8.7^2 - 6.8^2}$ ($\sqrt{29.45} = 5.42678542$) M1 for $\cos CBD = \frac{9.5}{2 \times "5.43"}$ or $\sin BDC = \frac{9.5}{2 \times "5.43"}$ M1 (dep) for $CBD = \cos^{-1}\left(\frac{9.5}{2 \times "5.43"}\right)$ (= 0.875) or $BDC = \sin^{-1}\left(\frac{9.5}{2 \times "5.43"}\right)$ A1 for $28.79 - 28.921$ Alternative methods (a combination may be seen) B1 for angle $ABD = 90^{\circ}$ M1 for using trigonometry for correct method to find angle and then correct trig statement involving OB M1 for correct method to make OB the subject M1 for correct method to find CD and correct trig statement or statement of Cosine rule that involves angle CBD M1 for correctly rearranging so CBD is the subject A1 for $28.79 - 28.921$