

Mark Scheme (Results)

June 2011

GCSE Mathematics (5384H)
Paper 13H (Non-Calculator)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:
<http://www.edexcel.com/Aboutus/contact-us/>

June 2011

Publications Code UG028394

All the material in this publication is copyright

© Edexcel Ltd 2011

NOTES ON MARKING PRINCIPLES

1 **Types of mark**

M marks: method marks

A marks: accuracy marks

B marks: unconditional accuracy marks (independent of M marks)

2 **Abbreviations**

cao – correct answer only

ft – follow through

isw – ignore subsequent working

SC: special case

oe – or equivalent (and appropriate)

dep – dependent

indep - independent

3 **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.

If there is no answer on the answer line then check the working for an obvious answer.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

5 **Follow through marks**

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.

Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.

If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

10 Money notation

Accepted with and without the “p” at the end.

11 Range of answers

Unless otherwise stated, when any answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1).

5384H_13H				
Question	Working	Answer	Mark	Notes
1	$15 \div 10$ 80×1.5 60×1.5 30×1.5 36×1.5	120, 90, 45, 54	3	M2 for any one of $80 + 40$ or $60 + 30$ or $30 + 15$ or $36 + 18$ or 120 or 90 or 45 or 54 seen A1 cao OR M1 for $15 \div 10$ or $3 \div 2$ or sight of 1.5 M1(dep) for $80 \times '1.5'$ or $60 \times '1.5'$ or $30 \times '1.5'$ or $36 \times '1.5'$ A1 cao OR M1 for $80 \div 10$ or $60 \div 10$ or $30 \div 10$ or $36 \div 10$ or 8 or 6 or 3 or 3.6 M1(dep) for $'8' \times 15$ or $'6' \times 15$ or $'3' \times 15$ or $'3.6' \times 15$ A1 cao OR M1 for $80 \div 2$ or $60 \div 2$ or $30 \div 2$ or $36 \div 2$ or 40 or 30 or 15 or 18 M1(dep) for $'40' \times 3$ or $'30' \times 3$ or $'15' \times 3$ or $'18' \times 3$ A1 cao
2	(a)	20	1	B1 cao
	(b)	18	1	B1 cao
	(c)	Graph completed	2	B1 for line from (12.40, 20) to (1.20, 20) ± 2 mm B1 ft line from "(1.20, 20)" to axis after 1hr 30 mins ± 2 mm.
3		$2 \times 5 + 3 \times -1$	2	M1 for 2×5 and 3×-1 or 10 and -3 seen A1 cao

5384H_13H					
Question		Working	Answer	Mark	Notes
4			Shaded region	3	B1 for circular arc radius 3 cm centre P within square ($\pm 2\text{mm}$) B1 for line 2 cm from PS within square ($\pm 2\text{mm}$) B1cao (see overlay)
5			$\frac{1}{8}$	1	B1 for $\frac{1}{8}$ or 0.125 or 8^{-1} oe
6	(a)		Reflection	2	B2 for vertices of shape plotted at (-3, 2), (-3, 3), (-5, 3), (-6, 2.5), (-5, 2) (B1 for a reflection in any vertical or horizontal line)
	(b)		Translation; $\begin{pmatrix} -6 \\ -1 \end{pmatrix}$	2	B1 for translation B1 for 6 left and 1 down OR $\begin{pmatrix} -6 \\ -1 \end{pmatrix}$ Note: B0 if more than one transformation given
7		$1 \text{ m}^2 = 100 \times 100 \text{ cm}^2$ $4 \text{ m}^2 = 40\,000 \text{ cm}^2$	Decision and explanation	2	B2 for states Callum is not correct with full correct explanation (B1 for 100×100 or 10000 seen or for 200×200 or 40 000 seen or for $= 0.01 \times 0.01$ or 0.0001 seen or for 0.04 seen)

5384H_13H				
Question	Working	Answer	Mark	Notes
8	<p>P: T: B = 1: 3: 6 $54 \div 10 \times 6$</p> <p>OR</p> <p>e.g. $T=3P$ $B=2T$ So, $B=2(3P)=6P$ $P+T+B=P+3P+6P=10P$ $P = 54 \div 10 = \text{£}5.40$ $B = 6 \times \text{£}5.40$</p>	32.40	3	<p>M1 for 1 : 3 : 6 or any 3 numbers in any order in the ratio 1 : 3 : 6 M1 for $54 \div (1 + 3 + 6) \times 6$ A1 for 32.4(0)</p> <p>Alternatives M1 for 1: 3: 6 oe or $P + 3P + 6P (=10P)$ oe eg $T/3 + T + 2T (=10T/3)$ or eg $B/6 + B/2 + B (=10B/6)$ or 5.4(0) or 16.2(0) seen M1 for $54 \div 10 \times 6$ or $[54 \div "10/3"] \times 2$ or $54 \div "10/6"$ "oe A1 for 32.4(0)</p> <p>OR M1 for a partial decomposition of £54 in ratio 1:3:6 e.g. $(\text{£})5 + (\text{£})15 + (\text{£})30 (=£50)$ M1 for a decomposition of the remaining amount in the ratio 1:3:6, e.g. $40(p) + 120(p) + 240(p) (=400p)$ A1 for 32.4(0)</p>

5384H_13H				
Question	Working	Answer	Mark	Notes
9	$\frac{1500 - 250}{250} \times \frac{100}{1}$	500	3	B1 for "1500 - 250" or difference of 1250 seen M1 for $\frac{1500 - 250}{250} \times \frac{100}{1}$ A1 cao OR M1 for $\frac{1500}{250} \times \frac{100}{1}$ A1 for 600 B1 ft for "600" - 100
10	$4^2 + 3^2$ $\sqrt{25}$	5	3	M1 for sight of "6 - 2" (=4) and "6 - 3" (=3) M1 (dep) for "4" ² + "3" ² A1 cao

5384H_13H

Question		Working	Answer	Mark	Notes																		
11	(a)	$\frac{1 \times 2}{5 \times 3}$	$\frac{2}{15}$	1	B1 for $\frac{2}{15}$ oe																		
	(b)	$(2-1) + \frac{5}{15} - \frac{6}{15}$ or $\frac{35}{15} - \frac{21}{15}$ or <table border="1" data-bbox="398 778 611 922"> <tr><td></td><td>1</td><td>3</td></tr> <tr><td>2</td><td></td><td>6</td></tr> <tr><td>5</td><td>5</td><td>15</td></tr> </table> <table border="1" data-bbox="398 976 618 1120"> <tr><td></td><td>7</td><td>5</td></tr> <tr><td>7</td><td></td><td>35</td></tr> <tr><td>3</td><td>21</td><td>15</td></tr> </table>		1	3	2		6	5	5	15		7	5	7		35	3	21	15	$\frac{14}{15}$	3	M1 for attempt to find a common denominator or sight of $\frac{5}{15}$ or $\frac{6}{15}$ or $\frac{35}{15}$ or $\frac{21}{15}$ oe or a fully correct table A1 for sight of $\frac{5}{15} - \frac{6}{15}$ or $\frac{35}{15} - \frac{21}{15}$ oe A1 for $\frac{14}{15}$ oe Alternative M1 for 0.33(3...) or 0.4 or 2.33(3...) or 1.4 A1 for 0.33(3...) - 0.4 or 2.33(3...) - 1.4 A1 for 0.93(recurring)
	1	3																					
2		6																					
5	5	15																					
	7	5																					
7		35																					
3	21	15																					

5384H_13H					
Question		Working	Answer	Mark	Notes
12	(a)		9	1	B1 cao
	(b)		$\frac{1}{8}$	1	B1 for $\frac{1}{8}$ oe
	(c)		t^6	1	B1 cao
	(d)	$\frac{n^4}{n^5}$	n^{-1}	2	M1 for $\frac{n^4}{n^{3+2}}$ or $\frac{n^4}{n^5}$ or $\frac{n^{4-3}}{n^2}$ or $\frac{n^1}{n^2}$ or $\frac{n^{4-2}}{n^3}$ or $\frac{n^2}{n^3}$ or a correct second step following an incorrect first step eg $\frac{n^4}{n^6} = n^{-2}$ oe A1 for n^{-1} (allow $\frac{1}{n}$)

5384H_13H				
Question	Working	Answer	Mark	Notes
13	$4x + y = 10$ $4x - 6y = 38$ $7y = -28, y = -4$ $4x - 4 = 10, x = 3.5$ OR $12x + 3y = 30$ $\frac{2x - 3y = 19}{14x} +$ $= 49, x = 3.5$ $7 - 3y = 19, y = -4$ Alternative $y = 10 - 4x$ $2x - 3(10 - 4x) = 19$ $14x - 30 = 19; x = 3.5$ $4 \times 3.5 + y = 10; y = -4$	$x = 3.5$ $y = -4$	3	M1 for full method to eliminate x or y , allow one error in calculation M1(dep) for substitution of one variable into one of the equations, or by appropriate method after starting again A1 for 3.5 and -4 oe Alternative M1 for full method to rearrange and substitute to eliminate x or y , allow one error in calculation M1(dep) for substitution of one variable into one of the equations, or by appropriate method after starting again A1 for 3.5 and -4 oe
14	(a)	(10), 3, 2, (1), -6	2	B2 for all 3 correct values (B1 for 2 correct values)
	(b)		2	B2 for fully correct graph OR B1 ft for 5 points plotted correctly $\pm 2\text{mm}$ B1 for smooth curve drawn through points provided B1 awarded in (a)

5384H_13H				
Question	Working	Answer	Mark	Notes
15	$t(k-2) = tk - 2t = k$ $tk - k = 2t$ $k(t-1) = 2t$ $k = \frac{2t}{t-1}$	$k = \frac{2t}{t-1}$	4	<p>M1 for attempt to multiply LHS by (k-2) or sight of $t(k-2)$ or $tk - 2t$ or $tk - 2$ (ignore RHS)</p> <p>M1 for attempt to subtract k from LHS or sight of $tk - k$ (ignore RHS) or attempt to subtract tk to give $-2t = k - kt$ (ignore LHS)</p> <p>M1 for attempt to factorise for k e.g. $k(t-1)$ or $k(1-t)$</p> <p>A1 for $k = \frac{2t}{t-1}$ or $k = \frac{-2t}{1-t}$ oe</p>
16	$\pi x l = 2\pi x^2$ $h^2 + x^2 = 4x^2$ $h^2 = 3x^2$ Alternative $\pi x \sqrt{h^2 + x^2} = 2\pi x^2$ $\sqrt{h^2 + x^2} = 2x$ $h^2 + x^2 = 4x^2$ $h^2 = 3x^2$	$\sqrt{3}x$	4	<p>B1 for curved surface area of one of the shapes eg $\pi x l$ or $2\pi x^2$</p> <p>M1 for attempt to equate surface areas eg $\pi x l = 2\pi x^2$ or $l = 2x$</p> <p>M1 for attempt to connect h and x using Pythagoras theorem eg $h^2 + x^2 = 4x^2$</p> <p>A1 for $\sqrt{3}x$ or $\sqrt{3}x^2$</p> <p>Alternative</p> <p>B1 $h^2 + x^2 = l^2$ oe</p> <p>M1 for attempt to equate surface areas eg $\pi x \sqrt{h^2 + x^2} = 2\pi x^2$ oe</p> <p>M1 (dep) for attempt to square both sides of their formula eg $h^2 + x^2 = 4x^2$</p> <p>A1 for $\sqrt{3}x$ or $\sqrt{3}x^2$</p> <p>If no marks awarded, SC B1 for attempt to equate surface areas in terms of r rather than x</p>

5384H_13H					
Question		Working	Answer	Mark	Notes
17	(a)	AB = AO + OB	-2a + 3b	1	B1 for -2a + 3b or 3b - 2a oe
	(b)	OP = 2a + $\frac{2}{5}(3b - 2a)$ = $\frac{6}{5}a + \frac{6}{5}b$ = $\frac{6}{5}(a + b)$ Parallel	$\frac{6}{5}(a + b)$ is parallel to a + b	3	M1 for 2a ± $\frac{2}{5}(3b - 2a)$ OR 3b ± $\frac{3}{5}(2a - 3b)$ A1 for $\frac{6}{5}a + \frac{6}{5}b$ oe A1 for $\frac{6}{5}(a + b)$ is parallel to a + boe
18		$\frac{x \times 2(x+1)}{2} - \frac{2 \times 2(x+1)}{x+1} = 1 \times 2(x+1)$ $x(x+1) - 4 = 2(x+1)$ $x^2 + x - 4 = 2x + 2$ $x^2 - x - 6 = 0$ $(x-3)(x+2) = 0$	x = 3, -2	4	M1 for an attempt to multiply one term of the equation by 2 or (x + 1) or $2(x+1)$ with or without cancelling or attempt to write LHS with a common denominator M1 for attempt to multiply all terms by $2(x+1)$ with or without cancelling e.g. $\frac{x \times 2(x+1)}{2} - \frac{2 \times 2(x+1)}{x+1} = 1 \times 2(x+1)$ or $x(x+1) - 4 = 2(x+1)$ A1 for $x^2 + x - 4 = 2x + 2$ or $x^2 - x - 6 = 0$ A1 cao for 3 and -2

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code UG028394 June 2011

For more information on Edexcel qualifications, please visit
www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual




Llywodraeth Cynulliad Cymru
Welsh Assembly Government



Rewarding Learning