Surname

Centre Number

0

Number

Other Names



GCSE

4370/06

MATHEMATICS – LINEAR PAPER 2 **HIGHER TIER**

A.M. MONDAY, 12 November 2012

2 hours

Suitable for Modified Language Candidates

ADDITIONAL MATERIALS

A calculator will be required for this paper.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer all the questions in the spaces provided.

Take π as 3.14 or use the π button on your calculator.

INFORMATION FOR CANDIDATES

You should give details of your method of solution when appropriate.

Unless stated, diagrams are not drawn to scale.

Scale drawing solutions will not be acceptable where you are asked to calculate.

The number of marks is given in brackets at the end of each question or part-question.

You are reminded that assessment will take into account the quality of written communication (including mathematical communication) used in your answer to question 4.

For E	xaminer's us	e only
Question	Maximum Mark	Mark Awarded
1	3	
2	10	
3	4	
4	8	
5	7	
6	9	
7	10	
8	6	
9	3	
10	11	
11	3	
12	6	
13	2	
14	9	
15	7	
16	2	
TOTAL	MARK	

1370 060001

Formula List

Area of trapezium =
$$\frac{1}{2}(a+b)h$$

Volume of prism = area of cross-section × length

Volume of sphere $=\frac{4}{3}\pi r^3$ Surface area of sphere $=4\pi r^2$


Volume of cone = $\frac{1}{3}\pi r^2 h$ Curved surface area of cone = πrl

In any triangle *ABC* Sine rule $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ Cosine rule $a^2 = b^2 + c^2 - 2bc \cos A$ Area of triangle $= \frac{1}{2}ab \sin C$


The Quadratic Equation

The solutions of
$$ax^2 + bx + c = 0$$

where $a \neq 0$ are given by



$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$



- 4
- 2. The table below gives information from the Highway Code on stopping distances for cars.




Diagram not drawn to scale

(a) A warning sign for a crossroads is to be placed on a road. The road has a speed limit of 30 mph.
 Find the minimum distance that the warning sign should be placed from the crossroads.

Find the minimum distance that the warning sign should be placed from the crossroads. Use the data given above to find your answer.

(b) An average car is approximately 4 metres in length. How many car lengths is the stopping distance for a car travelling at 40 mph?

[2]

<u>--</u>ا


[1]


(c) Complete the table below.

Spe	eed
mph	km/h
30	
50	80
	112

[3] The stopping distances given in the Highway Code are given for good driving conditions (d)and alert drivers (drivers who are not tired). When a driver is tired, the thinking distance increases by 30% and the braking distance increases by 20%. Calculate the stopping distance, in metres, for a tired driver travelling at 50 mph in good driving conditions. [4]

(4370-06)





Rotate the rectangle shown below through 180° about the point (2, 1). *(b)*

7

4370 060007

Examiner only

Turn over.

4. You will be assessed on the quality of your written communication in this question.

Kate lives in the UK. She goes on holiday to Paris with two friends, Janie who lives in America and Ami who lives in Japan.

They meet in London for a few days and then fly to Paris together.

Janie exchanges 450 American dollars to pounds and Ami exchanges 30000 Japanese yen to pounds.

In London, Janie and Ami each spend £100 and exchange their remaining money to euros. Kate also exchanges £250 to euros.

 $\pounds 1 = 129.82$ Japanese yen $\pounds 1 = 1.57$ American dollars $\pounds 1 = 1.18$ euros

Use the exchange rates shown above to calculate how many euros in total the three girls take to Paris.

You must explain each step of your calculations and show all your working.

[8]

4370 060009

In Kingstone, the mean daily snowfall for a week was 5.6 cm. What would the mean daily snowfall have been if it had snowed 2 cm more on each day? [1] (b)In Greyfield, the snowfall for each of 10 days was measured. The results are summarised in the table below. Daily snowfall, s, in cm Number of days 4 $4.5 \leq s < 5.5$ 2 $5 \cdot 5 \leq s < 6 \cdot 5$ $6 \cdot 5 \leq s < 7 \cdot 5$ 1 $7.5 \leq s < 8.5$ 1 2 $8.5 \leq s < 9.5$ Calculate an estimate for the mean daily snowfall for the 10 days. (i)

9

5.

(a)


[4] State the modal class. Modal class [1] Write down the class in which the median lies. [1]

(ii)

(iii)

Examiner Calculate the volume of a cylinder with a diameter of 4.6 cm and a height of 8.4 cm. 6. *(a)* ------..... [3]

only



© WJEC CBAC Ltd.


Arm length, <i>l</i> cm	$50 < l \leq 55$	$55 < l \leqslant 60$	$60 < l \leqslant 65$	$65 < l \leqslant 70$	$70 < l \leqslant 75$
Frequency	4	18	38	30	10
(a) On the	e graph paper be	low, draw a frec	uency polygon t	to show this data	a. [2]
	Frequency	7			
	40				
	30				
	20				
	10				

12

(b) Complete the following cumulative frequency table.

Arm length, <i>l</i> cm	$l \leqslant 50$	$l \leqslant 55$	$l \leqslant 60$	$l \leqslant 65$	$l \leqslant 70$	<i>l</i> ≤ 75
Cumulative frequency	0	4				

[1]




- 8. A warehouse stores electrical goods in boxes. The boxes are all cuboids.
 - (a) One of the boxes has a depth of 46 cm, a width of 55 cm and a length of 62 cm. All the measurements are correct to the nearest centimetre.
 - (i) Write down the greatest and least possible values for each of these measurements in the table below.

Dimension	Least value	Greatest value
Depth 46 cm	cm	cm
Width 55 cm	cm	cm
Length 62 cm	cm	cm

[2]

(ii) Hence, calculate the **greatest** possible volume of the box.

	[2	2]
(b)	Another box has dimensions $x \text{ cm}$ by $y \text{ cm}$ by $z \text{ cm}$. Each of these measurements is correct to the nearest cm. Find an expression for the least possible volume of this box in terms of x , y and z . You do not need to simplify your expression.	
•••••		••
•••••		••
		••
•••••		••
•••••		••
	[2	 2]




(a)	Solve the following simultaneous equations using an algebraic method.	Exan on
	3x + 2y = 27 $2x - 5y = 37$	
	~ 	
••••••		
.		
•••••		
•••••		
•••••		
<u>.</u>		
(1)	[4] $3 + x + 2x - 1 = 12$	
(<i>b</i>)	Solve $\frac{3+x}{2} + \frac{2x-1}{11} = 13.$	
•••••		
•••••		
••••••		
•••••		
•••••	[4]	

16


		17	
7r-b = ar - c [3] A unit of measure used with textiles is the denier. Silk is said to measure 1 denier when 9000m of a single strand of the silk weights 1 g. 1 denier is the same as 1g per 9000m (a) Complete the statement. 1 denier is the same as g per 450 m [J] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre	<i>(c)</i>	Rearrange the following formula to make r the subject.	Exar or
A unit of measure used with textiles is the denier. Silk is said to measure 1 denier when 9000 m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre			
A unit of measure used with textiles is the denier. Silk is said to measure 1 denier when 9000 m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre			
A unit of measure used with textiles is the denier. Silk is said to measure 1 denier when 9000 m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre			
A unit of measure used with textiles is the denier. Silk is said to measure 1 denier when 9000m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre			
A unit of measure used with textiles is the denier. Silk is said to measure 1 denier when 9000m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre			
A unit of measure used with textiles is the denier. Silk is said to measure 1 denier when 9000m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre			
A unit of measure used with textiles is the denier. Silk is said to measure 1 denier when 9000m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre			
Silk is said to measure 1 denier when 9000 m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre	••••••	[3]	
Silk is said to measure 1 denier when 9000 m of a single strand of the silk weights 1 g. 1 denier is the same as 1 g per 9000 m (a) Complete the statement. 1 denier is the same as g per 450 m [1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre			
 (a) Complete the statement. 1 denier is the same as			
1 denier is the same as		1 denier is the same as 1 g per 9000 m	
[1] (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as	<i>(a)</i>	Complete the statement.	
 (b) Complete the following statement. Give your answer in standard form correct to two significant figures. 1 denier is the same as g per metre 		1 denier is the same as g per 450 m	
two significant figures. 1 denier is the same as g per metre	•••••	[1]	
	<i>(b)</i>		,
		1 denier is the same as g per metre	
	.		
	.		
	••••••		
		[2]	

Turn over.



(b) A circle has a diameter FH. The point G is a point on the circumference of the circle. Given that the length of the straight line GH = 6.8 cm and GFH = 32°, calculate the length of FH.

(4370-06)



20

[2]

(4370-06)

	21	
l 4.	The point B is at the centre of the circle. The points P and Q are on the circumference of the circle.	Examiner only
	$A \xrightarrow{4 \text{ cm}} B \xrightarrow{8 \text{ cm}} B$	
	Diagram not drawn to scale Calculate the area of the shaded sector.	
	[9]	

enten	ce number	1	2	3	4	5	6
	number of a, e and r	8	6	8	3	4	5
	number of in the ce	36	22	42	8	10	14
(b)	All of the lette Two letters ar Calculate the Give your ans You must sho	re selected at probability t swer correct t	random from hat at least of two decin	n the bag an one of the le	d not repla		[2]
(b)	Two letters ar Calculate the Give your ans	re selected at probability t swer correct t	random from hat at least of two decin	n the bag an one of the le	d not repla		[2]
(b)	Two letters ar Calculate the Give your ans	re selected at probability t swer correct t	random from hat at least of two decin	n the bag an one of the le	d not repla		[2]
(b)	Two letters ar Calculate the Give your ans	re selected at probability t swer correct t	random from hat at least of two decin	n the bag an one of the le	d not repla		[2]

