M07/4/BIOLO/HP3/ENG/TZ2/XX/M+



B DIPLOMA PROGRAMME PROGRAMME DU DIPLÔME DU BI PROGRAMA DEL DIPLOMA DEL BI

MARKSCHEME

May 2007

BIOLOGY

Higher Level

Paper 3

13 pages

This markscheme is **confidential** and for the exclusive use of examiners in this examination session.

-2-

It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of IBCA.

General Marking Instructions

Subject Details: Biology HL Paper 3 Markscheme

Mark Allocation

Candidates are required to answer ALL questions in each of TWO Options (total [20 marks]). Maximum total = [40 marks].

-3-

General

A markscheme often has more specific points worthy of a mark than the total allows. This is intentional. Do not award more than the maximum marks allowed for part of a question.

When deciding upon alternative answers by candidates to those given in the markscheme, consider the following points:

- Each marking point has a separate line and the end is signified by means of a semicolon (;).
- An alternative answer or wording is indicated in the markscheme by a "/"; either wording can be accepted.
- Words in (...) in the markscheme are not necessary to gain the mark.
- Words that are <u>underlined</u> are essential for the mark.
- The order of points does not have to be as written (unless stated otherwise).
- If the candidate's answer has the same "meaning" or can be clearly interpreted as being the same as that in the markscheme then award the mark.
- Mark positively. Give candidates credit for what they have achieved, and for what they have got correct, rather than penalizing them for what they have not achieved or what they have got wrong.
- Remember that many candidates are writing in a second language. Effective communication is more important than grammatical accuracy.
- Occasionally, a part of a question may require a calculation whose answer is required for subsequent parts. If an error is made in the first part then it should be penalized. However, if the incorrect answer is used correctly in subsequent parts then **follow through** marks should be awarded.
- Units should always be given where appropriate. Omission of units should only be penalized once. Ignore this, if marks for units are already specified in the markscheme.
- Do not penalize candidates for errors in significant figures, unless it is specifically referred to in the markscheme.

Option D — **Evolution**

D1.	(a)	400 (\pm 20) million years ago	[1]
	(b)	10 (events)	[1]
	(c)	vertebrate hemoglobin and myoglobin are more closely related to each other because of a more recent spilt over evolutionary time / invertebrate hemoglobin more distantly related because of a more ancient (distant) split in evolutionary time	[1]
	(d)	animals have a higher metabolic rate/greater oxygen needs which requires transport of oxygen to cells / plants do not move thus have a lower metabolic rate / plants produce their own oxygen (through photosynthesis)	[1]
	(e)	the genetic code is degenerate/more than one codon will code for an amino acid; so a mutation may result in the same amino acid being produced; some mutations are neutral;	
		multiple point mutations for the same base (can obscure the mutation history);	[2 max]
D2.	(a)	Lamarck argued that acquired characteristics are inherited; <i>e.g.</i> long neck of giraffe; <i>(accept other suitable example)</i> Lamarck also proposed the theory of use and disuse / use causes organs to develop and disuse reduces them;	[2 max]
	(b)	forward facing eyes / binocular/stereoscopic vision; large brains; flexible shoulder joints / shoulder blades on the dorsal side of the thorax; colour vision; manual dexterity / power grip; finger pads / nails not claws; skull_adapted for upright posturo;	
		pronation / rotating hand;	[2 max]

-4-

[6 max]

D3. (a) sickle cell anemia is an example of balanced polymorphism; heterozygous (Hb^A Hb^S/ sickle cell traits) individuals are more resistant to malaria than homozygous individuals; therefore heterozygous individuals survive in greater numbers than those with other genotypes; consequently heterozygous individuals (on average) produce more offspring; maintains the Hb^S/ recessive allele in the population; this offsets deaths due to sickle cell anemia in the population;
(h) hisch winch

- 5 -

(b) *biochemical*:

use DNA and RNA as genetic material;

have same genetic code;

have same 20 amino acids in proteins / amino acids in all organisms are left-handed; conservation of sequences of nucleotides in DNA/amino acids in proteins between related species;

anatomical:

vertebrate embryos at early stages show similar anatomical features; homologous structures show similar anatomical pattern despite widely varying

function (*e.g.* the pentadactyl limb);

vestigial structures have had a function in ancestral organisms (*e.g.* appendix in humans);

neoteny/maintenance of embryological characteristics of ancestral species;

geographical:

related organisms/fossils show similar geographical distribution; *e.g.* placental mammals found in Asia and marsupial/monotremes found in Australia; isolation explains evolution of flora and fauna in Australia; continental drift helps explain the distribution of animals/fossils;

Accept any other valid answer for each section. At least one answer from each section must be present for full marks.

Option E — Neurobiology and Behaviour

E1. (a)
$$\frac{11}{41} \times 100 = 25\% (\pm 3\%)$$
 [1]
(b) $\frac{50-33}{50} \times 100 = 34\% (\pm 3\%)$ (less for angled run);
 $\frac{17}{33} \times 100 = 52\% (\pm 3\%)$ (more for a straight run);
 $\frac{50-33}{\left(\frac{50+33}{2}\right)} = \frac{17}{41.5} = 41\% (\pm 3\%)$ (less for angled run); [1 max]

- 6 -

- (c) grackles are more successful with males when using an angled run; more successful with females in a straight run; when using a straight run, they capture a similar number of male and female crabs; overall capture (many) more in angled run (only males); [2 max]
- (d) males are more conspicuous/easier to see than females; males are larger / greater energy source/ less mobile (due to one large claw) / angle of attack obscured by large male claw in angled attack; grackles are more visible to the crabs on a straight run and better able to defend themselves / angled run may surprise crabs / straight run may alert them to attack / in angled run, birds do not need to flap their wings as hard as often and this makes it easier to attack the prey;
- E2. (a) sympathetic; parasympathetic; *Both answers are required for [1].*
 - (b) rod cells absorb all wavelengths of light/monochromatic whereas cones absorb distinct wavelengths/red, green, blue; cones give greater visual sensitivity/acuity than rods; rods are more dispersed in the retina than cones / give wider field of vision whereas cones more concentrated in fovea; rod cells are more sensitive to dim light whereas cones function well in bright light; a group of rod cells connected to same (sensory) neuron whereas cones each connect to one individual neuron; [2 max]
 - (c) Award [1] for each two of the following clearly drawn and correctly labelled. sensory/receptor cells; sensory neuron with cell body; interneuron/association neuron with cell body; motor neuron with cell body; muscle/effector; dorsal root / ventral root / cell bodies;

[1]

E3. (a) (Parkinson's disease is) death of neurons (in *substantia nigra*); neurons release dopamine; at inhibitory synapses which control muscle contraction; muscle contraction not controlled without dopamine;
 symptoms: uncontrollable shaking/muscle contraction not controlled/tremor/increased muscle rigidity; slow movements; poor balance;

-7-

- speech impairment;
- Award [2 max] if no reference to any symptoms given.
- (b) some psycho-active drugs increase synaptic transmission / act like neurotransmitters / are not broken down (at the receptors) / some psycho-active drugs interfere with the breaking down / re-uptake of the neurotransmitters
- (c) innate: [4 max]

definition / develop independently of the environment / is inherited/instinct; stereotyped behaviour / no change with practice/repetition; innate behaviour develops by natural selection because they make a species better adapted to the environment / increase reproductive success; taxis: movement to or away from a directional stimulus; named example of taxis and how it helps *e.g.* fly larvae move away from light, protecting them from predators; kinesis: response to non-directional stimulus; named example of kinesis and how it helps *e.g.* woodlice move toward moist/dark places to avoid dehydration;

learned: [4 max]

definition / depend on the environment/experiences;

natural selection will favour certain learned behaviours that increase reproductive success;

classical conditioning: conditioned response to a conditioned stimulus;

operant conditioning: trial and error / reinforcement given after operant response; imprinting: learning a response in a receptive period;

named example and how it helps *e.g.* ducklings follow first thing they see that moves / mother when born to avoid predators;

conditioned behaviour based on response to rewards that increase survival;

[6 max]

[3 max]

[1]

Option F — **Applied Plant and Animal Science**

F1.	(a)	1.0 μ M (±0.2) (units required)	[1]
	(b)	production of H_2O_2 increased less in C (control) plants than A1/A5 plants; production of H_2O_2 in D41/D42 plants increased less than in C (control) plants; production of H_2O_2 increased the most in A1/least in D41 plants;	[3 max]
	(c)	transgenic plants with functioning gene showed more H_2O_2 production so hypothesis supported / data for A1 and A5 is variable, so difficult to conclude whether the hypothesis is supported / transgenic plants with gene suppressed showed less H_2O_2 production so hypothesis supported	[1]
	(d)	disease resistance could spread to species related to rice <i>e.g.</i> weeds; reluctance of consumers to eat genetically modified food products; outcompete natural species and change gene pool; may increase allergenic effects in human/cattle;	[1 max]
F2.	(a)	decomposers/fungi/bacteria can increase levels of soil nutrients needed for plant growth; fungi/bacteria/ <i>Rhizobium</i> can increase levels of soil nutrients needed for plant growth/mutualistic effects; pathogens/disease affect the health/growth rate/leaf area; predators consume plant tissues and reduce photosynthesis/growth rate;	[2 max]
	(b)	identify the highest yielding cows; artificial insemination using semen of bulls whose offspring are high yielding; female calves from these crosses should give higher milk yields;	[2 max]

F3. (a) use: [4 max]

antibiotics control infection so increase growth rate/reduce loss of animals; growth hormones boost production by increasing growth rates/yield; most humans do not seem to be affected by eating milk/meat that contain antibiotics; animals have same rights to protection against disease as humans;

-9-

misuse: [4 max]

overuse of antibiotics increases the rate of evolution of antibiotic resistance in bacteria;

antibiotic residue in foods that humans consume may cause allergies; steroid hormones used to promote growth may get into food or environment and alter development in humans / long-term effect of hormone use is unknown; animals given these hormones may give unnaturally high yields which cause the animal to suffer;

[6 max]

 (b) monoculture is where the same crop is grown year after year; same nutrients are absorbed from the soil; these nutrients tend to become depleted / harvest removes nutrients from the field nutrient cycle; fertilizer must be applied to renew nutrients;

use of crop rotation / leaving land fallow;

[4 max]

G1. (a) 65% (±3) (units required) [1] (b) wild type out competed by P28 but wild type out competed SP22 [1] (i) (ii) wild type out competed by SP22 but wild type out competed P28 [1] new strain would out compete the wild type; (c) because the (15:15) light-dark cycle supports the longer circadian rhythms; because it is closer to P28 /does not compete as completely as P28; [2 max] (d) competitive exclusion principle states that only one species can occupy a niche in an ecosystem; principle supported – in three experiments one species remained after 27 days and in the other three experiments one species dominated; principle not supported – in three experiments two species remained after 27 days; not possible to say - in three experiments one species remained after 27 days and in the other three experiments two species remained / experiment needs to run for a longer period; [2 max]**G2.** (a) Nitrosomonas: nitrification/oxidizes/converts ammonia to nitrites; Rhizobium: converts/reduces/fixes atmospheric nitrogen to ammonia/nitrogen fixation; [2]

•	
cling /	
ns die; [2	? max]
cling / ns die; [2	2 1

Option G — Ecology and Conservation

G3. (a) distribution affected by both abiotic and biotic factors; animals are adapted to specific temperatures / animals that maintain internal body temperature have a wider distribution than those that do not / animals require special adaptations to live in extreme temperature environments; some animals are aquatic and can only live in/near water / only animals adapted to dry environments will live there / some adapted to fresh water others to salt water; animals require/defend territory for breeding/feeding sites; animals will congregate near food supply; competition/predation may cause animals to live in areas other than their preferred areas / interactions between species / mutualism; human activities such as urban/industrial development / intensive agriculture / deforestation / desertification / conservation / pollution / use of insecticides / molluscicides / trapping / poisoning / shooting larger animals; [5 max] (b) bacteria found/grow in raw sewage; can cause disease in animals/humans; bacteria digest organic matter and release nutrients;

bacteria consume oxygen due to respiration and cause oxygen depletion / increase in BOD;

animals die due to lack of O_2 ;

can cause eutrophication/algal blooms;

aesthetic quality of the stream is reduced/ unpleasant smell;

[4 max]

Option H — Further Human Physiology

H1.	(a)	20 mg (unit required)	[1]
	(b)	increasing dose leads to decreasing the total cholesterol; higher doses have less effect than 10 mg dose / greater change with low dose compared to higher doses;	[2]
	(c)	drug lowers the low density lipoproteins and raises the high density lipoproteins / has opposite effects	[1]
	(d)	a placebo is the control against which changes in the other groups are measured / to safeguard against possible psychological factors affecting results	[1]
	(e)	affects bile production which may affect digestion; affects hormone biosynthesis which may cause hormonal imbalances / affect metabolism; affects cholesterol which may affect membranes of cells;	[1 max]
H2.	(a)	transport of lipids/fats/fat soluble vitamins (from digestive system); transport of leucocytes/white blood cells; returns (excess) tissue fluid to venous blood system; returns (leaked) proteins to venous blood system;	[2 max]
	(b)	Award [1 max] for any two of the following. cellulose/fibres; lignin/ keratin; bile pigments; bacteria; intestinal cells;	[1 max]
	(c)	endopeptidases hydrolyze/cut/break peptide linkages in the middle of (polypeptide) chains whereas exopeptidases hydrolyze/cut/break peptide linkages at the end of the chain; endopeptidases initiate digestion of large polypeptides whereas exopeptidases complete digestion of polypeptides (digestion into amino acids);	[1 max]

H3. (a) body temperature monitored by hypothalamus; when temperature drops neurosecretory cells/hypothalamus produce thyrotropin releasing hormone/TRH; TRH travels to the (anterior lobe of) pituitary (via the portal vein); pituitary secretes thyroid stimulating hormone/TSH; TSH travels to thyroid gland; thyroid secretes thyroxin; thyroxin increases metabolic rate; heat produced by increased metabolism increases body temperature / heat produced by metabolism inhibits production of thyrotropin releasing hormone/TRH; blood levels of thyroxin monitored by both hypothalamus and anterior pituitary; thyroxin/T3/T4 levels regulated by negative feedback; high levels of thyroxin/T3/T4 inhibit TRH secretion; [6 max] Accept TRF or TSH releasing factor instead of TRH. Accept converse statements for increase in body temperature. partial pressure of oxygen lower at high altitude; (b) hemoglobin may not become fully saturated; tissues may not get enough oxygen / causing fatigue; altitude sickness / suffer from pulmonary edema/headaches/dizziness; body can acclimatize by increasing ventilation rate/red blood cells/myoglobin /

increased capillary density;

[4 max]