



# **MARKSCHEME**

**May 2009**

**CHEMISTRY**

**Higher Level**

**Paper 3**

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of IB Cardiff.*

## Subject Details:      Chemistry HL Paper 3 Markscheme

### Mark Allocation

Candidates are required to answer questions from **TWO** of the options [**2 x 25 marks**]. Maximum total = [**50 marks**].

1. A markscheme often has more marking points than the total allows. This is intentional. Do not award more than the maximum marks allowed for part of a question.
2. Each marking point has a separate line and the end is signified by means of a semicolon (;).
3. An alternative answer or wording is indicated in the markscheme by a slash (/) - either wording can be accepted.
4. Words in brackets ( ) in the markscheme are not necessary to gain the mark.
5. Words that are underlined are essential for the mark.
6. The order of marking points does not have to be as in the markscheme, unless stated otherwise.
7. If the candidate's answer has the same "meaning" or can be clearly interpreted as being of equivalent significance, detail and validity as that in the markscheme then award the mark. Where this point is considered to be particularly relevant in a question it is emphasized by writing **OWTTE** (or words to that effect).
8. Remember that many candidates are writing in a second language. Effective communication is more important than grammatical accuracy.
9. Occasionally, a part of a question may require an answer that is required for subsequent marking points. If an error is made in the first marking point then it should be penalized. However, if the incorrect answer is used correctly in subsequent marking points then **follow through** marks should be awarded. Indicate this with **ECF** (error carried forward).
10. Only consider units at the end of a calculation. Unless directed otherwise in the markscheme, unit errors should only be penalized once in the paper. Indicate this by writing **-1(U)** at the first point it occurs and **U** on the cover page.
11. Significant digits should only be considered in the final answer. Deduct **1 mark in the paper** for an **error of 2 or more digits** unless directed otherwise in the markscheme.

*e.g.* if the answer is 1.63:

2	<i>reject</i>
1.6	<i>accept</i>
1.63	<i>accept</i>
1.631	<i>accept</i>
1.6314	<i>reject</i>

Indicate the mark deduction by writing **-1(SD)** at the first point it occurs and **SD** on the cover page.

12. If a question specifically asks for the name of a substance, do not award a mark for a correct formula, similarly, if the formula is specifically asked for, do not award a mark for a correct name.
13. If a question asks for an equation for a reaction, a balanced symbol equation is usually expected, do not award a mark for a word equation or an unbalanced equation unless directed otherwise in the markscheme.
14. Ignore missing or incorrect state symbols in an equation unless directed otherwise in the markscheme.

**Option A — Modern analytical chemistry**

- A1.** (a) absorption spectrum: energy required to move/excite (electrons) from lower/ground state to higher energy level/excited state;

emission spectrum: radiation emitted by electrons from higher/excited state to lower/ground energy level;

**OR**

absorption spectrum: continuous spectrum with missing regions/lines corresponding to energies absorbed;

emission spectrum: regions/lines corresponding to energies given out/emitted; [2]

- (b) (i) ( $^1\text{H}$ /proton) NMR/nuclear magnetic resonance; [1]

(ii) AA(S)/atomic absorption; [1]

- (c) (i) stationary phase is thin layer of  $\text{SiO}_2$ /silica/silicon dioxide/ $\text{Al}_2\text{O}_3$ /alumina/aluminium oxide (on support) (and mobile phase a solvent);  
*Ignore reference to  $\text{H}_2\text{O}$  in stationary phase.*

distance travelled/separation (of components) depends on adsorption/bonding to stationary phase / polarity of component / relative solubility between two phases; [2]  
*No mark for absorption.*

- (ii) separation is better/more effective/faster/efficient / separated component can be more easily recovered / withstands strong solvents / develops better; [1]

(iii)  $R_f = \frac{40(\text{mm})}{46(\text{mm})} = 0.87$ ; [1]

*Allow 0.86 to 0.88.*

*No mark if  $R_f$  has units.*

- A2.** (a)  $\text{C}_2\text{H}_4\text{O}_2$ ; [1]  
*No mark for  $(\text{CH}_2\text{O})_2$ .*

- (b)  $m/z = 15$   
 $\text{CH}_3^+$ ;


$m/z = 45$

$\text{COOH}^+/\text{CO}_2\text{H}^+/\text{HCOO}^+/\text{OCOH}^+$ ; [2]  
*Penalize once if charges are missing.*

- (c) ethanoic acid/ $\text{CH}_3\text{COOH}$  / methyl methanoate/ $\text{HCOOCH}_3$ ; [1]  
*Accept acetic acid.*

- A3. (a) infrared light is split into two (separate) beams;  
 one passes through the sample;  
 one through a reference;  
 a detector compares the two signals; [3 max]  
*Accept a suitable diagram.*

- (b) (i) absence of peak between 3200–3600  $\text{cm}^{-1}$  / above 3000  $\text{cm}^{-1}$  / peak for OH;  
 presence of peak between 1700–1750  $\text{cm}^{-1}$  / peak for C=O;  
 absence of peak between 1610–1680  $\text{cm}^{-1}$  / peak for C=C; [2 max]



*Accept CH<sub>3</sub>CH<sub>2</sub>CHO.*

3:2:1; [2]

*Ignore order*

*ECF if structure is incorrect only if its NMR spectrum contains three peaks.*

- (iii) triplet/1:2:1; [1]  
*ECF if structure is incorrect.*

- A4. (a) (isolated) double bonds/pi orbitals / limited conjugation; [1]

- (b) increasing wavelength means lesser energy / *OWTTE*;  
 1,3,5-hexatriene has three conjugated double bonds, so least energy;  
 1,3-pentadiene has two conjugated double bonds, it needs more energy;


*Accept alternative wording for M2 and M3:*

1,3-pentadiene has two conjugated double bonds, 1,3,5-hexatriene has three /  
 conjugation increases from 1,3-pentadiene to 1,3,5-hexatriene;  
 thus need more energy;

double bonds in 1,4-pentadiene are not conjugated, highest energy; [4 max]

**Option B — Human biochemistry**

**B1. (a)**



*Accept CHO, CH<sub>2</sub>OH and OH groups on either side of the carbon chain provided OH on C3 is on the opposite side to OHs on C2, C4 and C5.*

[1]

- (b) C1 is asymmetric/chiral/has four different groups;  
forming two isomers where the OH-group is on a different side (of C1/ring);  
 $\alpha$ -glucose **and**  $\beta$ -glucose;  
*Accept suitable diagrams as an explanation for M2.*

[3]

- (c) amylose **and** amylopectin;

[1]

- B2. (a)** steroid/steroidal backbone/4 ring/tetra cyclic carbon structure skeleton;  
*Do not accept OH, hydroxyl, hydroxide, alcohol.*  
*Accept a correct sketch of the steroid backbone.*

[1]

- (b) phospholipids;  
triglycerides/triglycerols;

[2]

- (c) (i) high density lipoprotein **and** low density lipoprotein;

[1]

- (ii) HDL has a higher proportion/percentage of proteins/LDL has a lower proportion of proteins / *OWTTE*;

[1]

- (d) LDL can be retained in the arteries / block arteries / start formation of plaque(s) / increases risk of atherosclerosis/arteriosclerosis/cardiovascular/heart diseases;

[1]

- B3.** (a) vitamin A: not water-soluble because it has only one OH / is not very/less polar / contains long hydrocarbon group;  
vitamin C: water-soluble because it has 4/many OH (and 1 C=O)/extensive hydrogen bonding;

[2]

*Accept reference to polarity in one case but not in both.*

- (b) effect: vitamin A: xerophthalmia/night blindness / vitamin C: scurvy / bleeding gums / less resistance to infection / bleeding lesions on legs/thighs / scorbatus;

*Accept either of the following for the second mark:*

solution for vitamin A: providing food composed of liver/fresh (orange and green) fruits/vegetables/spinach/eggs/carrots / providing genetically modified food containing vitamin A;


**OR**

solution for vitamin C: providing food composed of fresh fruits/vegetables/ providing genetically modified food containing vitamin C;

providing nutritional supplements / adding vitamin A/C in commonly consumed foods / vitamin A in margarine / vitamin C in fruit juices;

[3 max]

- B4.** (a) As [S] increases, more enzyme molecules can link with substrate molecules (and rate increases);  
 once all enzyme molecules are occupied/enzyme sites used up, rate cannot increase anymore/has no effect; [2]
- (b) 0.7 (mmol dm<sup>-3</sup>); [1]  
 Allow 0.6–0.8. If wrong units, then no mark.
- (c) inhibitor molecules are very similar to the substrate **and** the enzymes cannot distinguish them / are able to bind to/occupy the active sites / *OWTTE*; [1]
- (d) same  $V_{max}$ ;  
 at high [S], impact of inhibitor reduced / possible to reach same maximum rate;  
 the curve is less steep (than without inhibition) / the enzyme is less effective / *OWTTE*;  
 so  $K_m$  higher (than without inhibition); [4]
- (e) curve on graph levelling off below original line; [1]



$K_m$  should be approximately the same,  $V_{max}$  should be smaller.



**Option C — Chemistry in industry and technology**

- C1.** (a) homogeneous mixture of metals/a metal and non-metal; [1]
- (b) alloying element(s) disrupts regular/repeating (metal) lattice;  
difficult for one layer to slide over another / atoms smaller than metal cations can fit into the (holes of) metal lattice disrupting bonding;  
can make metal harder/stronger/more corrosion resistant/brittle; [2 max]
- (c) makes steel less brittle/softer/more ductile; [1]
- (d) Fe and Al production use large quantities of energy / about ten times/much more energy needed to produce Al than (similar mass of) Fe;  
(mined) areas can leave scars on landscape (unless mining companies re-vegetate areas) / destroys animal/plant habitat;  
(mining) waste products/tailings/metal wastes can damage environment / purification of bauxite produces considerable waste of iron(III) oxide which can result in visual pollution;  
(large volume of) CO<sub>2</sub> produced which contributes to global warming; [2 max]
- C2.** (a) contains no lithium/metal / uses lithium salt in an organic solvent (as electrolyte);  
involves movement of lithium ions (between electrodes); [2]
- (b) *Anode (-):*  
 $\text{LiC}_6 \rightarrow \text{Li}^+ + 6\text{C} + \text{e}^-$  / Li<sup>+</sup> ions dissociate from anode (and migrate to cathode);
- Cathode (+):*  
 $\text{Li}^+ + \text{e}^- + \text{MnO}_2 \rightarrow \text{LiMnO}_2$  /  $\text{Li}^+ + \text{e}^- + \text{CoO}_2 \rightarrow \text{LiCoO}_2$  /  $\text{Li}^+ + \text{e}^- + \text{FePO}_4 \rightarrow \text{LiFePO}_4$  /  
 $\text{Li}^+ + \text{e}^- + \text{NiO}_2 \rightarrow \text{LiNiO}_2$  / Li<sup>+</sup> ions are inserted into metal oxide/phosphate (structure); [2]
- Award [1] if electrodes are reversed.*
- (c) *Similarity:*  
both convert chemical energy directly into electrical energy / both use spontaneous redox reactions (to produce energy) / both are electrochemical cells/voltaic cells/galvanic cells;
- Difference:*  
fuel cells are energy conversion devices **and** rechargeable batteries are energy storage devices / fuel cells require constant supply of reactants **and** batteries have stored chemical energy/provide power until stored chemicals are used up / batteries can be recharged **and** fuel cells do not need recharging (have a continuous supply of fuel) / fuel cells are more expensive than rechargeable batteries / the reactions in a rechargeable battery are reversible **and** in a fuel cell are not; [2]

C3. (a)

	<b>Crystalline solid</b>	<b>Nematic phase liquid crystal</b>	<b>Pure liquid</b>
Positional order	yes	no	no
Directional order	yes	yes	no

;

;


[2]

*Need all three across table for each mark.*

- (b) liquid crystal molecules work over a limited temperature range;  
 ability of liquid crystal molecules to transmit light depends on molecular orientation /  
*OWTTE*;  
 orientation of polar molecules controlled by applying voltage across a small section;  
 areas of the material/display that are dark and light can be controlled;

[2 max]

C4. (a) (free) radical (addition); [1]



*Radical signs required for mark, no mark for radical sign on H.*

one electron from double bond joins with the electron in radical to form O–C bond;  
other carbon on double bond contains a single electron/double bond changes to  
single bond with electron on end carbon;

[3]

*Accept arrows showing the movement of electrons for these marks.*

- (c) addition polymerization: unsaturated/containing C=C monomers add together without elimination/ removal of any atoms (to form polymer) / *OWTTE*;  
condensation polymerization: monomers have two reactive sites/are bifunctional and produce a larger molecule with elimination/removal of a smaller molecule / *OWTTE*;

**OR**

in addition polymerisation unsaturated / containing C=C monomers undergo addition **and** in condensation polymerisation the monomer molecules have two reactive sites/ functional groups/ are bifunctional;

in the addition polymerisation reaction there are no by-products while in the condensation polymerisation reaction a small molecule is eliminated/ produced as a by-product;

[2 max]

- (d) chain length: greater chain length, greater intermolecular forces, higher strength / melting point;

way groups are arranged/orientated: trans-orientation allows close approach between chains giving it greater strength / cis-orientation produces bent chains, does not allow close approach for strong bonding / *OWTTE* / Kevlar example explained;

cross-linking: stronger covalent bonds in cross-linking lead to stronger / higher melting point / more rigid polymers (than linear or branched polymer) / *OWTTE* / amount of cross-linking determines how rigid structure becomes / vulcanized rubber example: sulfur atoms create strong covalent links between chains / in phenol-methanal (Bakelite), benzene ring is bonded to  $\text{CH}_2$  in three / several positions (to produce rigid plastic);

branching: depending on branching, close packing possible or prevented / amylose has a straight chain structure while amylopectin has straight linkages as well as branches;

[3 max]

*Need explanation for mark – no marks for branching related to LDPE and HDPE or isotactic and atactic arrangements as question is about condensation, not addition polymerization.*

**Option D — Medicines and drugs**

- D1.** measure of the relative margin of safety of a drug (for a particular treatment for a typical population) / measure for safe effective treatment;

$$\frac{\text{lethal dose (LD}_{50}\text{)}}{\text{therapeutic or effective dose (ED}_{50}\text{)}} /$$


ratio of the lethal dose (LD<sub>50</sub>) to the therapeutic or effective dose (ED<sub>50</sub>) / the range of dosage of a drug/its concentration in a bodily system/blood;

*Definition of LD<sub>50</sub> and ED<sub>50</sub> not required for mark.*

wide therapeutic window exists for small effective dose and larger lethal dose / toxicity occurs at much higher concentrations than for successful treatment / a big difference between effective and lethal dose / drugs with wide therapeutic window are safer;  
narrow therapeutic window requires small doses as lethal dose is not large / *OWTTE*;

[4]


- D2.** (a) react aspirin with sodium hydroxide/OH<sup>-</sup> to produce the (ionic) salt;  
*No MI for reaction with CaCO<sub>3</sub> (as calcium salicylate is not water soluble).*



[2]

*Accept ionic equation.  
ECF for M2*

- (b)



*Accept CF<sub>3</sub>C<sub>6</sub>H<sub>4</sub>OCH(C<sub>6</sub>H<sub>5</sub>)CH<sub>2</sub>CH<sub>2</sub>NHCH<sub>3</sub>*

[1]

- D3.** (a) (i) *Oxidation:*  

$$\text{C}_2\text{H}_5\text{OH} + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{COOH} + 4\text{H}^+ + 4\text{e}^- ;$$
- Reduction:*  

$$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} ; \quad [2]$$
- Accept balanced equation with molecular formulas.*  
*If both equations are wrong, award [1] for  $\text{C}_2\text{H}_5\text{OH} \rightarrow \text{CH}_3\text{COOH}$  and  $\text{Cr}_2\text{O}_7^{2-} \rightarrow 2\text{Cr}^{3+}$ .*  
*If correct equations are used but oxidation and reduction reversed, award [1].*
- (ii) orange to green; [1]
- (b) peak at  $2950 \text{ cm}^{-1}$  / absorption occurs due to C–H bonds in ethanol;  
*No mark for absorption due to just ethanol, or O–H bond in ethanol (water vapour in breath also contributes).*
- intensity / height of peak / absorption / amount of transmittance depends on amount of ethanol / compare absorption to standard / reference/control sample / sample containing no alcohol; [2]
- D4.** (a) side chain/alkyl group;  
*Accept hydrocarbon chain.*  
 modify side chain / use different R groups; [2]  
*Ignore reference to functional groups.*
- (b) ring consists of one nitrogen atom and three carbon atoms (and three hydrogen atoms);  
 4 membered/square ring structure / bond angles of  $90^\circ$  ;  
 ring under stress/strain / increased (chemical) reactivity / ring opens (due to angle of  $90^\circ$  instead of about  $109^\circ$  );  
 bonds to/blocks action of enzyme/transpeptidase;  
 reaction with the enzyme not reversible / prevents cross linking of peptides / inhibits synthesis and growth of bacterial cell walls / *OWTTE*; [5]
- (c) may wipe-out helpful/useful/beneficial bacteria (in the alimentary canal);  
 destroyed bacteria may be replaced by more harmful bacteria;  
 leads to resistance / makes penicillin less effective;  
 resistant bacteria grow / pass on their immunity/mutation/trait to succeeding generations / *OWTTE*; [2 max]
- D5.** HIV invades/bind to white blood cells/T4/T cells / *OWTTE*;  
 HIV viruses can mutate;  
 HIV viruses have similar metabolism to (human) cells / uses host cells to replicate;  
 high price of (antiretroviral) drugs / socioeconomic / cultural issues; [4]

**Option E — Environmental chemistry**

- E1.** (a) *NO*:  
 $N_2$  and  $O_2$  react in the engine /  $N_2 + O_2 \rightarrow 2NO$ ;  
*No mark for the high temperature without reference to the action between  $N_2$  and  $O_2$ .*
- NO<sub>2</sub>*:  
 NO oxidizes/reacts in the air to  $NO_2$  /  $2NO + O_2 \rightarrow 2NO_2$ ;
- CO*:  
 incomplete combustion; [3]  
 Accept balanced chemical equation for C5–C12 hydrocarbons.  
 Do not accept C1–C4.
- (b) combustion is incomplete therefore more CO is produced;  
 more VOCs remain unreacted;  
 (lack of air) reduces [NO]; [3]  
 Allow [1 max] if all correct answers correspond to air/fuel ratio.
- E2.** (a)  $Cl\cdot + O_3 \rightarrow ClO\cdot + O_2$ ;  
 $ClO\cdot + O\cdot \rightarrow Cl\cdot + O_2$ ; [2]  
*Radical symbols not required for mark.*
- (b) (i) heterogeneous (catalysis); [1]
- (ii) ice particles melt releasing the pollutants;  
 light breaks the bonds producing radicals / Cl; [2]
- E3.**  $2NH_3 + H_2SO_4 \rightarrow (NH_4)_2SO_4$  /  $2NH_3 + H_2SO_3 \rightarrow (NH_4)_2SO_3$  /  $NH_3 + HNO_3 \rightarrow NH_4NO_3$ ;  
 Accept ionic equations.
- $NH_4^+ + 2O_2 \rightarrow 2H^+ + NO_3^- + H_2O$ ;
- Any three of the following for [1] each:*  
 adding  $NH_3$  to soil makes it less acidic / OWTTE;  
 salt particles formed are acidic/have  $pH < 7$ ;  
 salts are washed into the ground;  
 in earth they can be nitrified;  
 nitrification acidifies the soil; [5 max]

- E4.** (a) (i) decayed plant and animal tissue/matter / soil biomass / non-living organic components / *OWTTE*; [1]  
*No mark for humus (question asks for main constituent of SOM).*
- (ii) *Provision of nutrients:*  
contains N/amine/amino acid / H (of COOH) can be interchanged by ions of nutrients / ion exchange of nutrients can occur due to H (of COOH);
- Water retention:*  
has (polar) OH/NH;  
these can form hydrogen bonds (with water molecules in the soil); [3]
- (b) irrigation waters contain salts;  
upon evaporation these stay (causing salinization of the soil);  
plants do not grow on soil with high salt content;  
over irrigation/water run off can remove nutrients/  $\text{Ca}^{2+}$  /  $\text{K}^+$  /  $\text{NH}_4^+$  /  $\text{Mg}^{2+}$  ; [3 max]
- (c) *Polyaromatic hydrocarbons (PAHs):*  
tar / coal / crude oil spills / incomplete combustion of wood/vegetation/waste / industrial/power station emissions;  
*Do not accept hydrocarbons.*
- Organotin compounds:*  
PVCs with tin compounds / antifouling compounds (coatings or paints) / fungicides / pesticides / biocidal agents; [2]

**Option F — Food chemistry**

- F1.** (a) ester of glycerol/propan-1,2,3-triol **and** three fatty acids/long chain carboxylic acids; [1]
- (b) II;  
more straight molecule/greater surface area hence greater distortion of electron cloud / allows closer packing of fatty acids for trans / does not allow closer packing for cis isomers / *OWTTE*;  
trans greater van der Waals' forces / cis less van der Waals' forces; [3]  
*Accept London/dispersion forces.*
- (c) addition of H<sub>2</sub> (to C=C bonds);  
finely divided Zn/Cu/Ni; [2]
- F2.** (a) protein;  
fat/lipid; [2]  
*Do not accept water.*
- (b) excludes oxygen/air;  
prevents oxidation;  
avoids decomposition with aerobic bacteria/fungi; [2 max]  
*Do not accept excludes light.*
- (c) (i) preserves meat / cures meat / fixes colour / inhibits microorganisms; [1]
- (ii) antioxidant; [1]
- F3.** (a) purplish-red colour of meat is produced by myoglobin;  
*Accept heme.*
- Fe has oxidation state +2 in myoglobin;  
(upon standing) oxidizes to Fe<sup>3+</sup> which is brown; [3]
- (b) Br<sub>2</sub> reacts with the double bonds / amount of double bonds in the conjugated system decreases;  
absorbed energy shifts to violet/higher energy in the visible region is absorbed;  
resulting in complementary yellow colour; [3]




**F4.** (a) salts of EDTA / extracts of plants/rosemary/tea/ground mustard **[1]**

(b) absorbers/quenchers of (free) radicals / reducing agent; **[1]**

(c)  $\beta$ -carotene: **[2]**  
 presence of conjugated/alternate single and double bonds;  
 electron donor / reducing agent (on oxygen in food);

**BHT: [3 max]**

phenol with bulky hydrocarbon/alkyl groups;



building more stable radicals;

due to the interaction of the free electron with the electrons in the aromatic ring; **[5 max]**

**Option G — Further organic chemistry**

**G1.** (a) bonding electron pair spread over three (or more) nuclei or atoms/not restricted/  
confined between two nuclei or atoms / *OWTTE*;

[1]


(b) physical evidence [2 marks]:

same carbon–carbon bond lengths in benzene / all carbon–carbon bonds are equal in length / forms regular hexagon;  
instead of longer single bonds **and** shorter double bonds / intermediate between single and double;

**OR**

3 isomers of  $C_6H_4X_2$ ;

not 4;

**OR**

not (structural) isomers / same compounds;

**OR**

( $^1H$ /proton) NMR spectrum shows only one peak / all the Hs in the same chemical environment;

not 2 peaks / not 2 different chemical environments;

**OR**

electron density maps;

show even electron density over ring;

chemical evidence [2 marks]:

hydrogenation of  $C_6H_6$  (1,3,5-cyclohexatriene) expected to produce three times as much energy as cyclohexene;

benzene produces less (due to delocalization);

**OR**

benzene undergoes substitution rather than addition reactions;

as it is more stable (due to delocalization);

**OR**

enthalpy of combustion of  $C_6H_6$  less than expected;

different bond energies;

[4 max]

- (c) reaction is faster with  $C_6H_5CH_2Cl$ ;  
 C–Cl bond weaker/easier to break;  
 because no overlap between a lone electron pair of Cl with ring delocalized electrons;  
 attacking  $OH^-$  nucleophile not repelled by the delocalized electrons;

**OR**



reaction is slower with  $C_6H_5Cl$ ;  
 C–Cl bond stronger/harder to break;  
 because of overlap between a lone electron pair of Cl with ring delocalized electrons  
 / delocalization increases electron density on ring / delocalization reduces  $\delta^+$  charge  
 on C (attached to Cl atom);  
 attacking  $OH^-$  nucleophile repelled by delocalized electrons / attraction of  
 nucleophile decreases / less polar C does not attract  $OH^-$  as much;

[3 max]

- (d) 1-chloro-3-nitrobenzene / meta-chloronitrobenzene / m-nitrochlorobenzene /  
 3-chloro-1-nitrobenzene / 3-chloronitrobenzene / 1-nitro-3-chlorobenzene /  
 3-nitrophenylchloride;  
 $NO_2$  electron withdrawing/attracting;  
 deactivates ring / reduces electron density on ring;  
 slower rate compared to benzene;  
 greater charge distribution in the 3-position / lesser charge in the 2- and 4-positions;

[5]


G2.



Accept condensed structural formulas

[2]

- G3.** (a) reaction of Mg/magnesium with halogenoalkane/named compound;  
in an anhydrous/dry solvent /  $(C_2H_5)_2O$ /diethyl ether; [2]  
*Accept equation with condition specified.*



*Accept either of the two following alternatives for the second and third mark.*




**OR**



[3 max]

*If  $\text{CH}_3\text{COCH}_3$  and  $\text{CH}_3\text{MgBr}$  or  $\text{C}_2\text{H}_5\text{COCH}_3$  and  $\text{C}_2\text{H}_5\text{MgBr}$  combination given, then award only [1].*

- G4.**  $\text{CH}_3\text{COCl} + \text{NaOH} \rightarrow \text{CH}_3\text{COOH} + \text{NaCl}$  /  $\text{CH}_3\text{COCl} + \text{OH}^- \rightarrow \text{CH}_3\text{COOH} + \text{Cl}^-$  ;  
addition-elimination mechanism;



movement of electrons from  $\text{OH}^-$  to C, originating from negative charge or oxygen atom;  
movement of electrons in the  $\text{C}=\text{O}$ ;  
*Partial polarity on  $\text{C}=\text{O}$  and  $\text{C}-\text{Cl}$  not required for mark.*

intermediate;  
*Curly arrows not required for mark.*

[5]