



**FURTHER MATHEMATICS
STANDARD LEVEL
PAPER 1**

Monday 14 May 2001 (afternoon)

1 hour

INSTRUCTIONS TO CANDIDATES

- Do not open this examination paper until instructed to do so.
- Answer all the questions.
- Unless otherwise stated in the question, all numerical answers must be given exactly or to three significant figures as appropriate.
- Write the make and model of your calculator on the front cover of your answer booklets *e.g.* Casio *fx-9750G*, Sharp *EL-9400*, Texas Instruments *TI-85*.

*A correct answer with **no** indication of the method used will usually receive **no** marks. You are therefore advised to show your working. In particular, where graphs from a graphic display calculator are being used to find solutions, you should sketch these graphs as part of your answer.*

1. (a) Explain when the Yates continuity correction needs to be used, giving a reason.
 (b) In 200 tosses of a coin, 108 tails and 92 heads were observed. Test the hypothesis that it is a fair coin, at a significance level of 1% .

2. Let (G, \circ) be a group with identity element e . Given that $x \circ x = e$ for all $x \in G$, prove that (G, \circ) is an Abelian group.

3. The profit of an internet company at the end of a given year is 8000 dollars more than twice the profit for the previous year. If the profit at the end of the first year is \$30 000, find an expression for profit at the end of the n th year, for $n = 1, 2, \dots$.

4. Let $(\mathbb{R}, +)$ be the group of real numbers under addition, and (\mathbb{R}^+, \times) be the group of positive real numbers under multiplication. Prove that the two groups are isomorphic.

5. The points T , C and D lie on a circle with centre S . A tangent [OT] and a secant [OCD] are drawn from a point O to this circle. Prove that $OT^2 = OC \times OD$.

6. (a) Prove that the series $\sum_{n=0}^{\infty} \frac{(-1)^n}{(n+1)^7}$ converges, for $n \in \mathbb{N}$.
 (b) Approximate the sum of the series to an accuracy of six decimal places.

7. Let $I = \int_0^5 e^{-x^2} dx$. Find the number n , of intervals necessary to approximate correct to two decimal places, the value of I by the trapezium rule.

8. Prove that a line $y = mx + c$ is a tangent to an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ if $c^2 = a^2 m^2 + b^2$.
9. A manager of two coal mines wants to test the heat-producing capacity of coal from each mine. The heat-producing capacity (in millions of calories per ton) of random samples of coal from each mine is given in the following table.

Mine 1	8260	8130	8350	8070	8340	
Mine 2	7950	7890	7900	8140	7920	7840

The manager knows that the two population variances are equal.

- (a) Describe the test to be used with the choice of the test statistic, giving reasons for your answers.
- (b) At the 5% level of significance, test if the average heat-producing capacity of the coal from the two mines is equal.
10. Let G be a simple graph. Prove that G has a spanning tree if and only if G is connected.
-

