



**MATHEMATICS
HIGHER LEVEL
PAPER 2**

Friday 8 May 2009 (morning)

2 hours

Candidate session number

0	0							
---	---	--	--	--	--	--	--	--

INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all of Section A in the spaces provided.
- Section B: answer all of Section B on the answer sheets provided. Write your session number on each answer sheet, and attach them to this examination paper and your cover sheet using the tag provided.
- At the end of the examination, indicate the number of sheets used in the appropriate box on your cover sheet.
- Unless otherwise stated in the question, all numerical answers must be given exactly or correct to three significant figures.

ExamsBuddy



5. [Maximum mark: 6]

Find the angle between the lines $\frac{x-1}{2} = 1-y = 2z$ and $x = y = 3z$.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

ExamsBuddy



SECTION B

Answer **all** the questions on the answer sheets provided. Please start each question on a new page.

10. [Maximum mark: 18]

Let $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}$.

(a) Given that $X = B - A^{-1}$ and $Y = B^{-1} - A$,

(i) find X and Y ;

(ii) does $X^{-1} + Y^{-1}$ have an inverse? Justify your conclusion.

[5 marks]

(b) Prove by induction that $A^n = \begin{pmatrix} 1 & n & \frac{n(n+1)}{2} \\ 0 & 1 & n \\ 0 & 0 & 1 \end{pmatrix}$, for $n \in \mathbb{Z}^+$.

[7 marks]

ExamsBuddy

(c) Given that $(A^n)^{-1} = \begin{pmatrix} 1 & x & y \\ 0 & 1 & x \\ 0 & 0 & 1 \end{pmatrix}$, for $n \in \mathbb{Z}^+$,

(i) find x and y in terms of n ,

(ii) and hence find an expression for $A^n + (A^n)^{-1}$.

[6 marks]



11. [Maximum mark: 23]

The position vector at time t of a point P is given by

$$\vec{OP} = (1+t)\mathbf{i} + (2-2t)\mathbf{j} + (3t-1)\mathbf{k}, t \geq 0.$$

- (a) Find the coordinates of P when $t = 0$. [2 marks]
- (b) Show that P moves along the line L with Cartesian equations

$$x-1 = \frac{y-2}{-2} = \frac{z+1}{3}. \quad [2 \text{ marks}]$$

- (c) (i) Find the value of t when P lies on the plane with equation $2x + y + z = 6$.
- (ii) State the coordinates of P at this time.
- (iii) Hence find the total distance travelled by P before it meets the plane. [6 marks]

The position vector at time t of another point, Q, is given by


$$\vec{OQ} = \begin{pmatrix} t^2 \\ t \\ 1-t^2 \end{pmatrix}$$

- (d) (i) Find the value of t for which the distance from Q to the origin is minimum.
- (ii) Find the coordinates of Q at this time. [6 marks]
- (e) Let \mathbf{a} , \mathbf{b} and \mathbf{c} be the position vectors of Q at times $t = 0$, $t = 1$ and $t = 2$ respectively.
 - (i) Show that the equation $\mathbf{a} - \mathbf{b} = k(\mathbf{b} - \mathbf{c})$ has no solution for k .
 - (ii) Hence show that the path of Q is not a straight line. [7 marks]



12. [Maximum mark: 19]

Let f be a function defined by $f(x) = x + 2\cos x$, $x \in [0, 2\pi]$. The diagram below shows a region S bound by the graph of f and the line $y = x$.



ExamsBuddy

A and C are the points of intersection of the line $y = x$ and the graph of f , and B is the minimum point of f .

- (a) If A, B and C have x -coordinates $a\frac{\pi}{2}$, $b\frac{\pi}{6}$ and $c\frac{\pi}{2}$, where $a, b, c \in \mathbb{N}$, find the values of a , b and c . [4 marks]
- (b) Find the range of f . [3 marks]
- (c) Find the equation of the normal to the graph of f at the point C, giving your answer in the form $y = px + q$. [5 marks]
- (d) The region S is rotated through 2π about the x -axis to generate a solid.
 - (i) Write down an integral that represents the volume V of this solid.
 - (ii) Show that $V = 6\pi^2$. [7 marks]

