MARK SCHEME for the October/November 2010 question paper

for the guidance of teachers

0606 ADDITIONAL MATHEMATICS

0606/11

Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0606	11

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Accuracy mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
 B2, 1, 0 means that the candidate can earn anything from 0 to 2.

Page 3	Page 3 Mark Scheme: Teachers' version		Paper
	IGCSE – October/November 2010	0606	11

The following abbreviations may be used in a mark scheme or used on the scripts:

- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)

Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures this is regarded as an error in accuracy.
- OW –1,2 This is deducted from A or B marks when essential working is omitted.
- PA –1 This is deducted from A or B marks in the case of premature approximation.
- S –1 Occasionally used for persistent slackness usually discussed at a meeting.
- EX –1 Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0606	11

		1	
1	(i) $a = -12, b = -4$	B1, B1 [2]	B1 for each
	(ii) -4	√ B1 [1]	Follow through on their y value
2	(i) Graphs	B1 B1 [2]	B1 for one correct curve B1 for a second correct curve consistent with the first curve
	(ii) 3	√B1 [1]	Follow through on number of clear points of intersection
3	$\frac{\cos x(1+\sin x) + \cos x(1-\sin x)}{1-\sin^2 x}$ $\frac{2\cos x}{\cos^2 x}$ $2\sec x$	M1 DM1 M1 A1 [4]	M1 for attempt to get in terms of a single fraction DM1 simplifying numerator M1 simplifying denominator
4	$x = -1 \text{ or } 7 \text{ or } -\frac{1}{2} \text{ seen}$ Either $(x+1)(2x^2 - 13x - 7)$ or $(x-7)(2x^2 + 3x + 1)$ or $(2x+1)(x^2 - 6x - 7)$ leading to $(x+1)(x-7)(2x+1)$	M1 DM1 A1 DM1, A1 [5]	M1 for attempt to find a root DM1 for attempt to obtain quadratic factor A1 correct quadratic factor DM1 attempt to factorise quadratic factor
5	(i) $a = \pi + \frac{\pi}{3}, a = \frac{4\pi}{3}$	B1 [1]	Must be in terms of π
	(ii) $\frac{dy}{dx} = 2x \cos x + 2 \sin x$ at P , $\frac{dy}{dx} = 2$, \Rightarrow grad of normal $= -\frac{1}{2}$ normal: $y - \frac{4\pi}{3} = -\frac{1}{2}\left(x - \frac{\pi}{2}\right)$	M1, A1 M1	M1 for attempt to differentiate a product M1 for $m_1m_2 = -1$, must have used differentiation
	$\left(2y = \frac{19\pi}{6} - x\right)$	M1, A1 [5]	M1 for attempt at a normal equation, must have used differentiation, allow unsimplified

	Page 5	Mark Scheme: Teachers	Syllabus	Paper			
	IGCSE – October/November 2010			0606 11			
6	(i) 64-960	$0x + 6000x^2$	B1, B1, B1 [3]	B1 for each correct term, allow 2^6			
	(ii) $1 \times (\text{their } x \text{ term}) + \frac{10}{2} \times (\text{their } 64)$ = -640			M1 for 2 terms B1 for $\frac{10}{2}$ or 5			
7	(a) (i) $x =$	30°, 150°	[3] B1, B1 [2]	B1 for each			
	x =	$30^{\circ} = 120^{\circ}, 240^{\circ}$ $150^{\circ}, 270^{\circ}$ $B = \{30^{\circ}, 150^{\circ}, 270^{\circ}\}$	B1 √B1 [2]	B1 for $x = 150^\circ$, 270° or Follow through on their			
	$3x = 0^{\circ},$	± 1 or $\tan 3x = 0$ 180°, 360°, 540° 0°, 120°, 180°	M1 A1 √B1 [3]	M1 for dealing with sec and 3 <i>x</i> A1 for all solutions correct Follow through on their number of solutions			
8	Use of ra $\ln y = 6.3$ $\ln y = b$ $A = e^{(\text{their})}$	$\ln x + \ln A$	M1 M1 A1 B1 M1 A1, A1 [7]	M1 for attempt at gradie M1 for attempt at y inter A1 for $\ln y = 6.8$ B1 for $\ln y = b \ln x + \ln$ M1 for use of e A1 for A and A1 for b	rcept		
9	(i) $A = x^2$,	$\Rightarrow \frac{\mathrm{d}A}{\mathrm{d}x} = 2x$	B1 [1]				
	(ii) When $x = \frac{dx}{dt} = \frac{0}{10}$ $= 0.0003$	$\frac{003}{10}$	√ B1 M1 A1 [3]	Follow through on their M1 for 0.003 ÷ their 10	$\frac{\mathrm{d}A}{\mathrm{d}x}$		
	(iii) $V = 4x^3$ $\frac{dV}{dt} = 12$ $= 0.09$	$\frac{dV}{dx} = 12x^2$ $x^2 \times 0.0003$	B1, B1 M1 A1 [4]	B1 for each			

Page 6 Mark Scheme: Teachers' version		Syllabus	Paper
	IGCSE – October/November 2010		11

10	(i)	$\tan\frac{\pi}{6} = \frac{4}{PA}, \ PA = 4\sqrt{3}$	B1	B1 for <i>PA</i> (answer given)
		$PB = \frac{4}{\sin\frac{\pi}{6}} + 4, PB = 12$	B1	B1 for <i>PB</i> (answer given)
		allow equivalent methods	[2]	
	(ii)	Sector area = $\frac{1}{2}12^2 \times \frac{\pi}{3}$	$\sqrt{\mathbf{B}1}$	$\sqrt{B1}$ sector area, ft on their <i>PB</i>
		Area of kite = $2 \times \frac{1}{2} \times 4\sqrt{3} \times 4$	M1, A1	M1 for attempt to find area of kite or appropriate triangle
		Shaded area $= 47.7$	A1 [4]	
	(iii)	$P = \left(12 \times \frac{\pi}{3}\right) + 2\left(12 - 4\sqrt{3}\right) + 2(4)$	B1, B1, B1	B1 for each of the 3 terms
		= 30.7	B1 [4]	B1 for final answer
11	(i)	$2(1+x)^{\frac{1}{2}}(+c)$	M1, A1 [2]	M1 for $(1+x)^{\frac{1}{2}}$, A1 for 2
	(ii)	$\frac{dy}{dx} = \frac{2\sqrt{1+x} - 2x\frac{1}{2}(1+x)^{-\frac{1}{2}}}{1+x}$	M1 A2, 1, 0	M1 attempt at differentiation -1 each error
		$=\frac{2}{\left(\sqrt{1+x}\right)}-\frac{x}{\left(\sqrt{1+x}\right)^3}$	A1 [4]	A1 all correct
	(iii)	$\int \frac{x}{\left(\sqrt{1+x}\right)^3} \mathrm{d}x = \int \frac{2}{\left(\sqrt{1+x}\right)} \mathrm{d}x - \frac{2x}{\sqrt{1+x}}$	M1	M1 for idea of using (ii) 'in reverse'
		$=4\sqrt{1+x}-\frac{2x}{\sqrt{1+x}}(+c)$	A1	A1 all correct
		$\int_{0}^{3} \frac{x}{\left(\sqrt{1+x}\right)^{3}} \mathrm{d}x = (8-3) - (4), = 1$	M1, A1	M1 for attempt evaluation
		· · · ·	[4]	

	Page							Syllabus	Paper	
		IGCSE – October/Novem			ber 2	010		0606	11	
12		$v = \frac{4x^3}{3}$	9x(+c) 3, $y = 1$, so $c = -8$		M1, 2 M1, 2			attempt to integra	te	
), leads to $x = \pm 1.5$ 5, -17), (-1.5, 1)		M1 A1, <i>A</i>	A1 [3]	M1 for a	attempt to solve	$\frac{\mathrm{d}y}{\mathrm{d}x} = 0$	
	G	iradient c	AB: $(0, -8)$ of AB = -6, perp grad = x - 6y = 48	$\frac{1}{6}$	M1 M1 M1, 2	A1 [4]	M1 for a	 for attempt to find midpoint for attempt to find grad of perp must be working with perp 		
12	$\frac{d}{d}$	20 = 2A -	$e^{2x}-Be^{-x}$		B1 M1 A1 DM1 A1	[5]	A1 all co	attempt to different prrect attempt to solve		
	e^{3x}	3x = 2	$e^{2x} - 40e^{-x}$, $20e^{2x} = 40$ or 0.231)e ^{-x}	M1 M1 M1 A1	[4]	solution M1 for d	equating to zero a lealing with expo attempt to obtain oth	onentials	
		<i></i>	$e^{2x} + 40e^{-x}$ we, so min		M1 A1	[2]	other val	attempt at second lid method correct conclusio		

© UCLES 2010

www.XtremePapers.net