

	UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education
CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER
ADDITIONAL MATHEMATICS 0606/12 Paper 1 October/November 2013	

Candidates answer on the Question Paper.

Electronic calculator Additional Materials:

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question. The use of an electronic calculator is expected, where appropriate. You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question. The total number of marks for this paper is 80.

This document consists of 16 printed pages.



Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \cdot$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where *n* is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$.


2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\csc^2 A = 1 + \cot^2 A$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc \cos A$$
$$\Delta = \frac{1}{2} bc \sin A$$



2 Find the set of values of k for which the curve $y = (k + 1)x^2 - 3x + (k + 1)$ lies below the x-axis. [4]

[Turn over

3 Show that $\frac{1+\sin\theta}{\cos\theta} + \frac{\cos\theta}{1+\sin\theta} = 2\sec\theta$.

[4] For Examiner's Use $A = \left\{ x : \cos x = \frac{1}{2}, 0^{\circ} \le x \le 620^{\circ} \right\},\$

 $B = \{x: \tan x = \sqrt{3}, 0^{\circ} \le x \le 620^{\circ}\}.$

4

The sets *A* and *B* are such that

(i) Find n(A).

(ii) Find n(B). [1] (iii) Find the elements of $A \cup B$. [1] (iv) Find the elements of $A \cap B$. [1]

[1]

For Examiner's

Use

5 (i) Find $\int (9 + \sin 3x) dx$.

(ii) Hence show that $\int_{\frac{\pi}{9}}^{\pi} (9 + \sin 3x) dx = a\pi + b$, where *a* and *b* are constants to be found. [3]


[3] For Examiner's Use

6 The function $f(x) = ax^3 + 4x^2 + bx - 2$, where *a* and *b* are constants, is such that 2x - 1 is a factor. Given that the remainder when f(x) is divided by x - 2 is twice the remainder when f(x) is divided by x + 1, find the value of *a* and of *b*. [6]

For Examiner's Use

(a) (i)	Find how many different 4-digit numbers can be formed from the digits 1, 3, 5, 6, 8 and 9 if each digit may be used only once.	[1]	For Examiner's Use
(ii)	Find how many of these 4-digit numbers are even.	[1]	
tea	ream of 6 people is to be selected from 8 men and 4 women. Find the number of difference ms that can be selected if		
(i)	there are no restrictions,	[1]	
(ii)	the team contains all 4 women,	[1]	
(iii)	the team contains at least 4 men.	[3]	

7



(iii) Given that k is a positive constant, state the set of values of k for which |(x-2)(x+3)| = k has 2 solutions only.

[1]

[3]

For Examiner 's Use

9 (a) Differentiate $4x^3 \ln(2x+1)$ with respect to x.

(**b**) (**i**) Given that
$$y = \frac{2x}{\sqrt{x+2}}$$
, show that $\frac{dy}{dx} = \frac{x+4}{(\sqrt{x+2})^3}$. [4]

(ii) Hence find
$$\int \frac{5x+20}{(\sqrt{x+2})^3} dx$$
. [2] For Examiner's Use

11

(iii) Hence evaluate
$$\int_2^7 \frac{5x+20}{(\sqrt{x+2})^3} dx$$
.

[2]

10	Solutions to this question by accurate drawing will not be accepted.	For
	The points $A(-3, 2)$ and $B(1, 4)$ are vertices of an isosceles triangle ABC, where angle $B = 90^{\circ}$.	Examiner's Use
	(i) Find the length of the line AB . [1]	


(ii) Find the equation of the line *BC*.

[3]

(iii) Find the coordinates of each of the two possible positions of *C*.

[6]

For Examiner's Use



(b) Given that the matrix
$$\mathbf{C} = \begin{pmatrix} x & -1 \\ x^2 - x + 1 & x - 1 \end{pmatrix}$$
, show that det $\mathbf{C} \neq 0$. [4]
For Examiner's Use

15

12 (a) A function f is such that $f(x) = 3x^2 - 1$ for $-10 \le x \le 8$.

(ii) Write down a suitable domain for f for which f^{-1} exists. [1]

(b) Functions g and h are defined by

$$g(x) = 4e^{x} - 2 \text{ for } x \in \mathbb{R},$$

$$h(x) = \ln 5x \text{ for } x > 0.$$
(i) Find $g^{-1}(x)$.
[2]

(ii) Solve gh(x) = 18.

[3]

For

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.