UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the June 2005 question paper

0620 CHEMISTRY

0620/02

Paper 2 (Core Theory), maximum mark 80

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the *Report on the Examination*.

 CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the June 2005 question papers for most IGCSE and GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Grade thresholds for Syllabus 0620 (Chemistry) in the June 2005 examination.

	maximum	minimum mark required for grade:			
	mark available	А	С	E	F
Component 2	80	N/A	57	42	31

The threshold (minimum mark) for B is set halfway between those for Grades A and C. The threshold (minimum mark) for D is set halfway between those for Grades C and E. The threshold (minimum mark) for G is set as many marks below the F threshold as the E threshold is above it.

Grade A* does not exist at the level of an individual component.


IGCSE

MARK SCHEME

MAXIMUM MARK: 80

SYLLABUS/COMPONENT: 0620/02

CHEMISTRY (Core Theory)



	ı ag	· ·	IGCSE – JUNE 2005	0620	2
1	(a)	(i)			[1]
•	()		B+E		[1]
			A + C		[1]
		(iv)			[1]
			B+E		[1]
	/ b \	. ,			
	(b)	(1)	graphite NOT: charcoal		[1]
		(ii)	diamond/buckminsterfullerene NOT: graphite (but ALLOW: ecf from part (i)		[1]
	(c)		.OW: Na ⁺ I⁻ T: Na ⁺ + I⁻		[1]
	(d)	bec ALL	npound (no mark) ause two different (types of) atoms <u>joined/bonded</u> etc. OW: two different elements bonded T: atoms together		[1]
			G		Total = 9
2	(a)		ring/filtration T: distillation, NOT: decanting		[1]
	(b)		us turns pink/red T: orange		[1]
	(c)	(i)	steel		[1]
		(ii)	water NOT: steam		[1]
		(iii)	copper/iron		[1]
		(iv)	natural gas NOT: methane		[1]
	(d)	met	hane		[1]
	(e)	100	° C (100 = 0)		[1]
	(f)	(i)	calcium/Ca ²⁺		[1]
		(ii)	e ⁻ ALLOW: e NOT: electron		[1]
	(g)	(i)	carbon dioxide + water ACCEPT: correct formulae		[1]

Mark Scheme

Syllabus

Paper

Page 1

	Page 2		Mark Scheme	Syllabus	Paper
<u> </u>			IGCSE – JUNE 2005	0620	2
			carbon monoxide NOT: CO		[1]
					Total = 12
3	(a)	rand far a	om arrangement; part.		[2]
	(b)	рН9			[1]
	(c)	(i)	NH_3/H_3N		[1]
		(ii)	covalent		[1]
		(iii) weak forces <u>between</u> particles OR molecules/weak intermolecular forces		rces [1]	
	(d)	(i)	H ₂ SO ₄		[1]
		(ii)	nitrogen; soil		[2]
	(e)	Any	3 from:		
		parti move NOT NOT move	nonia <u>evaporates</u> from (ammonia) solution; sion; <u>cles/molecules</u> of ammonia/gases are in constant move e freely; ALLOW: move fast : particles of ammonia solution move freely : move from high to low concentration ement of <u>particles/molecules</u> is random. : ammonia spreads out;	ement/	[3]
	(f)	(i)	the air ALLOW: atmosphere		[1]
		(ii)	2 (NO ₂)		[1]
			reversible reaction ALLOW: equilibrium		[1]
			exothermic/heat given out		[1]
		(,			Total = 16
1	(a)	mon	omers		[1]
-	` ,			s a single C	
	(D)	NOT NOT	s not have a double bond/only contains single bonds/has : has a single bond : it is saturated : no spare bonds	s a single C –	C bond [1]
	(c)		ayed/graphical formula correct OW: correct dot and cross diagrams		[1]
	(d)		breaking down of long-chained hydrocarbons/formation hydrocarbons from larger	of smaller	[1]

. ug		IGCSE – JUNE 2005	0620	2			
	(ii)	high temperature ALLOW: heat (catalyst alone = 0)	0020	[1]			
	(iii)	C_8H_{18} ALLOW: other sensible combinations $2(C_2H_4) + C_6H_{14}$		[1]			
(e)	(i)	H_2		[1]			
	(ii)	any 2 of:					
	temperature gradient in fractionation column; smaller/lighter molecules (rise) higher in column OR smaller/lighter molecules more easily vaporised OR e.g. referring to larger/heaver molecules ALLOW: hydrocarbons in place of molecules NOT: lighter/heavier fractions different fractions condense at particular heights in column/fractions condense when temperature falls below their boiling points ALLOW: different fractions have different boiling points/condense at different temperatures						
	(iii)	petrol: fuel (for cars)/other suitable use		[1]			
	NOT: for cars etc. lubricating fraction: lubricating oils/waxes/polishes/other suitable use						
		NOT: for planes etc.		Total = 11			
(a)	(i)	molecule containing 2 atoms ALLOW: element containing 2 atoms		[1]			
	(ii)	whether it is solid, liquid or gas (all 3 needed)		[1]			
(b)	(i)	gas; liquid; solid (all 3 = 2 marks; 2 correct = 1 mark)					
	(ii)) red/brown/orange or combination of these		[1]			
	(iii)	130-210 (°C) (actual = +184 °C)					
(c)	iodi	ne + potassium chloride (1 each)		[2]			
(d)	(i)	8 electrons in each shell + atoms joined no bonding electrons = 1 IGNORE: inner shell electrons if correct (incorrect inner shell electrons = 1 max)		[2]			
	(ii)	water purification OR treatment/killing bacteria etc./bleaching agent (for paper)/ making refrigerants/making organic chlorine compounds (named)/making solvents/extracting titanium/detinning scrap tinplate/making hydrochloric acid/extraction of bromine from seawater/other suitable use					

Mark Scheme

Page 3

5

Syllabus

Paper

	Page 4			Mark Scheme	Syllabus	Paper	
				IGCSE – JUNE 2005	0620	2	
	(e)	(i)	Α			[1]	
		(ii)	С			[1]	
		(iii)		it contains ions; which can <u>move</u> /are free to <u>move</u> (OWTTE) reference to electrons = 0			
	(f)	(i)	Ρ	eriod 6		[1]	
		(ii)	8	5		[1]	
		(iii)	di N	atoms with (same number of protons and) different number of neutrons/ lifferent mass number/different nucleon number NOT: molecules with NLLOW: elements with			
		(iv)	12	25		[1]	
						Total = 19	
6	(a)	(i)	-1 N	on + sulphuric acid → iron sulphate + hydrogen I per error/omission OT: iron(III) sulphate OT: hydrogen sulphate		[2]	
		(ii)	р	ghted splint; op/small explosion etc. consequential marking)		[2]	
	(b)	(i)	Ca	athode		[1]	
		(ii)	(ii) allows conduction (of electricity)/allows charges or ions to flow through the solution			the solution [1]	
		(iii)	N de A	ains layer of copper/coated with copper OT: gets bigger ecreases in size/gets smaller/loses copper etc. LLOW: the copper dissolves OT: breaks up/flakes off		[1] [1]	
		(iv)	0	queous sodium hydroxide; light blue ppt; insoluble in e PR aqueous ammonia; light blue ppt; soluble in excess/ consequential marking)		blue solution [3]	
	(c)	not NO NO	o make them attractive/makes them shiny/protects the iron so it does not rust/does ot corrode OR oxidise easily/less reactive than iron [1] [OT: chromium is unreactive etc. [IOT: other properties of chromium e.g. hard [IOT: corrosive			rust/does [1]	
	(d)	iron	>	chromium > copper		[1]	
						Total = 13	