

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

	CANDIDATE NAME		
	CENTRE NUMBER	CANDIDATE NUMBER	
*			
0 4	CHEMISTRY		0620/61
444	Paper 6 Alternat	tive to Practical	May/June 2010
5 0			1 hour
3 9	Candidates ans	wer on the Question Paper.	
8 0	No Additional M	aterials are required.	

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
4		
5		
6		
7		
Total		

This document consists of 14 printed pages and 2 blank pages.




UNIVERSITY of CAMBRIDGE International Examinations

[Turn over

WWW_XTREMEPAPERS_NET

1 The diagram shows the apparatus used to prepare a gas. The gas is more dense than air.







2 Three bottles of liquids have lost their labels.

The liquids are known to be:

aqueous sodium iodide,

hexene,

dilute nitric acid.

Outline chemical tests you could use to distinguish between the liquids in the three bottles.

liquid	test	result
aqueous sodium iodide		
hexene		
dilute nitric acid		

[6]

[Total: 6]

Examiner's Use

For

copper carbonate

The volume of carbon dioxide produced was measured every minute for six minutes.

nitric acid

(a) Use the gas syringe diagrams to complete the table of results.

time/minutes	gas syringe diagram	total volume of carbon dioxide produced/cm ³
0	0 10 20 30 40 50 60	
1	0 10 20 30 40 50 60	
2	annin ann an an ann an ann ann ann ann a	
3	100 100 30 40 50 60 70 80 90 100	
4		
5		
6	απητητητητητητητητητητητητητητητητητητητ	

gas syringe

The temperature of the nitric acid was 20 °C.

For


Examiner's

Use

0620/61/M/J/10

³ The speed of reaction between excess copper carbonate and dilute nitric acid was investigated using the apparatus below.

20 0 2 0 3 4 5 6 1 time/minutes [4] (d) Sketch on the grid, the graph you would expect if the experiment was repeated using nitric acid at a temperature of 60 °C. [2] [Total: 12]



(b) Plot the results on the grid below and draw a smooth line graph.

120

5

For Examiner's Use

0620/61/M/J/10

WWW_XTREMEPHPERS_NET

4 A student investigated the reaction of aqueous sodium hydroxide with two different acids, acid **C** and acid **D**.

For Examiner's Use

Two experiments were carried out.

Experiment 1

By using a measuring cylinder, 20 cm³ of aqueous sodium hydroxide was poured into a conical flask and the initial temperature of the solution was measured.

A burette was filled with acid \mathbf{C} up to the 0.0 cm³ mark.

 5 cm^3 of acid **C** was added to the sodium hydroxide in the flask. The temperature of the mixture was measured.

Further 5 cm³ portions of acid **C** were added to the mixture in the flask, stirring with the thermometer until a total volume of 30 cm^3 of acid **C** had been added. The temperatures after each 5 cm³ portion had been added were measured.

(a) Use the thermometer diagrams to record the temperatures in the table of results.



For Examiner's Use

Table of results

7

volume of acid C added/cm ³	thermometer diagrams	temperature/°C
0	30 -25 -20	
5	40 -35 -30	
10		
15	40 35 30	
20		
25	35 	
30	35 30 25	

[2]

Experiment 2

The burette was emptied and rinsed with water. Experiment 1 was repeated using acid **D**.

(b) Use the thermometer diagrams to record the temperatures in the table of results.

volume of acid D added/cm ³	thermometer diagrams	temperature/°C
0	30 -25 -20	
5	30 25 20	
10	25 20	
15	30 25 20	
20	30 25 20	
25	25 20	
30	30 -25 -20	

Table of results

For


Examiner's Use

0620/61/M/J/10

(c) Plot the results for Experiments 1 and 2 on the grid and draw two smooth line graphs. Clearly label your graphs.

9

For Examiner's Use



[2]

0620/61/M/J/10

(e)	(i)	Which experiment produced the larger temperature change?	E
	(ii)		
(f)	Why	y was the burette rinsed with water in Experiment 2?	
(g)		dict the temperature of the reaction mixture in Experiment 2 after 1 hour. Explain your wer.	
		[Total: 18]	

For Examiner's Use

5 Solid E was analysed. E was an aluminium salt. The tests on the solid and some of the observations are in the following table. Complete the observations in the table.

	tests	observations
tests o	n solid E	
(a) Ap	pearance of solid E .	white crystalline solid
. ,	little of solid E was heated in a st-tube.	colourless drops of liquid formed at the top of the tube
• •	ittle of solid E was dissolved in distilled Iter.	
tes	e solution was divided into four st-tubes and the following tests were rried out.	
 (i) To the first test-tube of solution, drops of aqueous sodium hydroxide were added. 		
	Excess sodium hydroxide was then added to the test-tube.	[3]
(ii)	Test (i) was repeated using aqueous ammonia solution instead of aqueous	
	sodium hydroxide.	
(iii)	To the third test-tube of solution, dilute hydrochloric acid was added, followed by barium chloride solution.	no reaction
(iv)	To the fourth test-tube of solution, aqueous sodium hydroxide and aluminium powder were added. The mixture was heated.	effervescence pungent gas given off turned damp litmus paper blue

0620/61/M/J/10


(d)	What does test (b) tell you about solid E .
	[1]
(e)	Identify the gas given off in test (c)(iv).
(f)	What conclusions can you draw about solid E?
	[2]
	[Total: 9]

0620/61/M/J/10

WWW.XTREMEPAPERS.NET

For Examiner's Use **6** Hydrated cobalt chloride crystals, $CoCl_2.6H_2O$, were heated in the apparatus shown below.

For Examiner's Use



(a)	Indicate on the diagram, using an arrow, where heat is applied.	[1]
(b)	The crystals change colour from to to	[1]
(c)	What is the purpose of the ice?	
		[1]
(d)	Why is the tube open at point W ?	
		[1]
	[Tota	: 4]



7	Malachite is a naturally occurring form of copper carbonate. Outline how a sample of copper metal could be obtained from large lumps of malachite in the laboratory. Copper is one of the least reactive metals. Your answer should include any chemicals used and conditions.	For Examiner's Use
	[6]	
	[Total: 6]	

0620/61/M/J/10

BLANK PAGE

© UCLES 2010

0620/61/M/J/10

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

0620/61/M/J/10



16 BLANK PAGE