

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**  
International General Certificate of Secondary Education

## **MARK SCHEME for the May/June 2014 series**

### **0620 CHEMISTRY**

**0620/33**

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0620	33

- 1 (a) carbon dioxide (1) [1]
- (b) propene (1) [1]
- (c) krypton (1) [1]
- (d) nitrogen (1) [1]
- (e) fluorine (1) [1]
- (f) sulfur dioxide (1) [1]
- (g) hydrogen (1) [1]
- [Total: 7]
- 2 (a) any **three** from:  
 particles have more energy (1)  
 move faster (1)  
 collide more frequently (1)  
 more particles have energy greater than  $E_a$  [3]  
**guidance:** more colliding molecules have enough energy to react is worth (2)
- (b) particles move in all directions/randomly in both liquids and gases (1)  
 no bonds/very weak forces between particles in gases (1)  
 molecules can move apart/separate (to fill entire volume) (1)  
**OR**  
 bonds/forces/IMF between particles in liquids (1)  
 molecules cannot move apart/separate (so fixed volume in liquids) (1) [3]
- [Total: 6]
- 3 (a) (i) enzymes (1) [1]
- (ii) reduces growth of microbes/rate of reproduction of microbes is lower/  
 microbes are dormant (1)  
 fewer (enzymes) to decay food (1)  
**OR**  
 enzymes less efficient at lower temperatures (1)  
 slower reaction rate (1) [2]
- (b) correct linkage (1)  
 rest of molecule correct **and** continuation shown (1)  
 (other product is) water (1) [3]

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0620	33

- (c) any **three** from:  
 photosynthesis (1)  
 light/photochemical (1)  
 chlorophyll/chloroplasts (1)  
 carbon dioxide and water needed (1)  
 (glucose and) oxygen (1) [3]

[Total: 9]

- 4 (a) (i) heat limestone/calcium carbonate (1)  
 fractional distillation (1)  
 liquid air (1) [3]

- (ii) any **two** of the oxides, C, S, P and Si, mentioned (1)  
 carbon dioxide and sulfur dioxide escape/are gases (1)  
  
 phosphorus oxide **or** silicon(IV) oxide react with calcium oxide/  
 phosphorus oxide **or** silicon(IV) oxide are acidic and calcium oxide is basic (1)  
  
 to form a slag **or** calcium silicate **or** calcium phosphate (1)  
  
 must have correct equation for one of the above reactions (1) [5]

- (b) (i) lattice/rows/regular arrangement of cations/positive ions/ $\text{Fe}^{2+}$  (1)  
 mobile/free/delocalised/sea of electrons (1) [2]

- (ii) the rows of ions/ions can move past each other (1)  
 without the metal breaking/bonds are not directional/not rigid (1) [2]

- (iii) carbon particles/atoms different size (1)  
 prevents movement of rows, etc. (1) [2]

[Total: 14]

- 5 (a) faster reaction rate (1)  
 higher collision rate (1)  
 greater yield **or** favour RHS (1)  
 pressure favours products because it has lower volume/fewer product molecules (1) [4]

- (b) higher temperature favour endothermic reaction (1)  
 this is the back reaction/left hand side/reactants (1)  
 reduce yield (1) [3]

- (c) (i) greater surface area (1) [1]

- (ii) increase reaction rate (1)  
 can use a lower temperature to have an economic rate (1)  
 and not decrease yield (by increasing temperature). [2]

Page 4	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0620	33

- (d) lower the temperature (1)  
 only ammonia will liquefy (1)  
**OR**  
 add water (1)  
 only ammonia will dissolve (1)  
**OR**  
 increase pressure (1)  
 only ammonia will liquefy (1) [2]

- (e) second line  $+3 \times 155 = +465$   
 third line  $-3 \times 280 = (-)840$   
 fourth line  $-3 \times 565 = (-)1695$   
 all **three** correct (2)  
 two correct (1)
- $1170 + 465 = 1635$   
 $840 + 1695 = 2535$   
 both numerically correct (1)  
 exothermic reaction with some reasoning (1) [4]

[Total: 16]

- 6 (a) (i) C and H only (1) [1]  
 (ii) only single bonds (1) [1]
- (b) (i)  $C_nH_{2n+2}$  (1) [1]  
 (ii)  $C_{14}H_{30}$  (1)  
 $(14 \times 12) + 30 = 198$  (g) (1) [2]
- (c) (i)  $C_9H_{20} + 14 O_2 \rightarrow 9CO_2 + 10H_2O$  (2) [2]  
 (ii) Volume ratio  

$C_xH_y(g)$	$+ O_2(g)$	$\rightarrow$	$CO_2(g)$	$+$	$H_2O(l)$	
20	160		100			all in $cm^3$ mole ratio
1	8		5			
$C_5H_{12}$	$+ 8O_2$	$\rightarrow$	$5CO_2$	$+$	$6H_2O$	

 For evidence of method (1)  
 for equation as above (2) [3]
- (d) (i) alkanes in petrol/fuel/solvent (1)  
 alkenes to make alcohols/plastics/polymers/solvents (1)  
 hydrogen to make ammonia/fuel/fuel cells, etc. (1) [3]
- (ii) a correct equation for example:  
 $C_{10}H_{22} \rightarrow C_8H_{16} + C_2H_4 + H_2$  (1) [1]

Page 5	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0620	33

(e) (i) light or lead tetraethyl/catalyst/high temperature (1) [1]

(ii)  $\text{CH}_3\text{-CHCl-CH}_3$  (1) [1]

[Total: 16]

7 (a) bauxite (1) [1]

(b) electrolyte alumina/aluminium oxide dissolved in molten cryolite (1)  
 use cryolite to reduce mp/comparable idea/temperature of electrolyte 900 to 1000°C (1)  
 electrodes carbon (1)  
 aluminium formed at cathode/ $\text{Al}^{3+} + 3\text{e} \rightarrow \text{Al}$  (1)  
 oxygen formed at anode/ $2\text{O}^{2-} \rightarrow \text{O}_2 + 4\text{e}$  (1)  
 anode burns/reacts to carbon dioxide/ $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$  (1) [6]

(c) (i) food containers/window frames/cooking foil/cars/bikes/drink cans (1) [1]

(ii)  $4\text{OH}^- \rightarrow \text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}$  (2) [2]

$4\text{Al} + 3\text{O}_2 \rightarrow 2\text{Al}_2\text{O}_3$  (2) [2]

[Total: 12]