

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.Write in dark blue or black pen.You may use an HB pencil for any diagrams or graphs.Do not use staples, paper clips, glue or correction fluid.DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

0

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units. Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use
--------------------

Total

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 7 printed pages and 1 blank page.



IB14 06\_0620\_52/3RP © UCLES 2014

[Turn over

1 You are going to investigate the temperature rise produced when different lengths of magnesium ribbon react with excess dilute sulfuric acid.

### Read all the instructions below carefully before starting the experiments.

#### Instructions

You are going to carry out five experiments.

(a) Experiment 1

Using a measuring cylinder, pour 20 cm<sup>3</sup> of dilute sulfuric acid into the beaker. Measure the initial temperature of the acid and record it in the table below. Add the 2 cm length of magnesium ribbon to the acid in the beaker, and stir the mixture with the thermometer. Measure the highest temperature reached and record it in the table. Remove the thermometer and rinse out the beaker with water.

(b) Experiment 2

Repeat Experiment 1, using the 3 cm length of magnesium ribbon. Record the initial and highest temperatures in the table.

(c) Experiment 3

Repeat Experiment 1, using the 4 cm length of magnesium ribbon. Record the temperatures in the table.

(d) Experiment 4

Repeat Experiment 1, using the 6 cm length of magnesium ribbon. Record the temperatures in the table.


(e) Experiment 5

Repeat Experiment 1, using the 7 cm length of magnesium ribbon. Record the temperatures in the table.

Complete the table.

Experiment	initial temperature /°C	highest temperature /°C	change in temperature /°C
1			
2			
3			
4			
5			

3



4

(i)	(i)	Which experiment gave the greatest change in temperature?
	(ii)	Suggest why the change in temperature was greatest in this experiment.
(j)		at difference would be observed if Experiment 1 was repeated using an equal mass of gnesium powder? Explain your answer.
		[2]

(k) Draw a diagram of apparatus you could use to collect and measure the volume of gas given off in the reaction.

[2]

[Total: 21]

2 You are provided with mixture E. E consists of two solids, F and G. Solid F is water-soluble and solid G is insoluble.

Carry out the following tests on **E**, recording all of your observations in the table. Conclusions must **not** be written in the table.

		tests	observations
test	<u>s or</u>	the mixture <b>E</b>	
(a)	De	scribe the appearance of the mixture.	[1]
(b)		ce a little of mixture <b>E</b> in a test-tube.	
	Heat the mixture gently at first then more strongly for about 1 minute. After 1 minute, test the gas given off with damp pH indicator		
	paper. Leave the test-tube to cool.		[3]
Add the rest of mixture <b>E</b> to about 10 cm <sup>3</sup> of distilled water in a boiling tube. Stopper the boiling tube and shake the contents for about a minute. Filter the contents of the boiling tube. Keep the residue and the filtrate for the following tests.			
tests on the filtrate			
(c)	(i)	To about 1 cm <sup>3</sup> of the filtrate, add about 1 cm <sup>3</sup> of aqueous sodium hydroxide. Gently heat the mixture. Test the gas given off with damp pH indicator paper.	[1]
	(ii)	Add about 1 cm <sup>3</sup> of silver nitrate solution to the second portion of the filtrate followed by about 1 cm <sup>3</sup> of dilute nitric acid.	[1]

tests on the residue		
<ul> <li>(d) Using a spatula, transfer a little of the residue from the filter paper to a test-tube. Using a teat pipette, add about 2 cm<sup>3</sup> of dilute hydrochloric acid to the residue. Test the gas given off.</li> </ul>	[2]	
Add an equal volume of distilled water to the solution in the test-tube. Shake the contents and divide into two portions.		
<ul> <li>(e) (i) Add several drops of aqueous sodium hydroxide to the first portion of the solution. Now add excess aqueous sodium hydroxide.</li> </ul>	[3]	
(ii) Add several drops of aqueousammonia to the second portion. Now add excess aqueous ammonia.	[2]	
(f) Explain your observations in test (b).		
	[2]	
(g) What conclusions can you draw about solid F?		
	[2]	
(h) What conclusions can you draw about solid G?		

# **BLANK PAGE**

## NOTES FOR USE IN QUALITATIVE ANALYSIS

### **Test for anions**

anion	test	test result
carbonate (CO <sub>3</sub> <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO₃⁻) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2–</sup> ) [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

### Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al <sup>3+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH <sub>4</sub> +)	ammonia produced on warming	-
calcium (Ca2+)	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

# Test for gases

gas	test and test results
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue
carbon dioxide (CO <sub>2</sub> )	turns limewater milky
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper
hydrogen (H <sub>2</sub> )	'pops' with a lighted splint
oxygen (O <sub>2</sub> )	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.