



### UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

**COMBINED SCIENCE** 

0653/51

Paper 5 Practical Test

May/June 2012

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials:

As listed in Instructions to Supervisors.

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Chemistry practical notes for this paper are printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
Total		

This document consists of 10 printed pages and 2 blank pages.



1 You are going to test the effect of light on the chemical composition of leaves. You are provided with two leaves from two separate plants of the same species.

For Examiner's

Leaf **A** has been removed from a plant that has been in strong light for a period of 24 hours.

Leaf **B** has been removed from a plant that has been kept in the dark for 48 hours.

- Half fill a 250 cm<sup>3</sup> beaker with water. Heat the water with a Bunsen burner until it is boiling.
- Turn off the Bunsen burner.
- Pick up leaf **A** with a pair of tweezers (forceps) and hold it in the hot water for 15 seconds.
- Using the tweezers place the leaf into the bottom of the large test-tube labelled **A**.
- Repeat the above procedure for leaf **B**, but place the leaf into the bottom of the large test-tube labelled **B**.
- Cover the leaves with alcohol and place the tubes **A** and **B**, into the beaker of hot water (water bath) for 5 minutes. The Bunsen burner must **not** be lit.
- Carefully remove both tubes from the water bath and pour off the alcohol into the waste beaker.

( ) ( )	Suggest what substance from the leaf has dissolved in the alcohol to produce the colour you have described.
	colour

(a) (i) Describe the colour of the alcohol that you have poured off from the tubes.

(ii)	•	Remove the leaves from the tubes with tweezers and hold them in the wate
		path for 2 to 3 seconds

Place the leaves into the separate petri dishes (or onto the white tiles) labelled
A and B. Use the tweezers to lay them flat and cover both leaves with iodine solution. Leave for about 2 minutes.

substance [2]

- Wash off the excess iodine solution with cold water. Pour any excess liquid into the waste beaker.
- Observe and record the colour of the leaves in Table 1.1.
- The colour of each leaf suggests a substance that may be present or absent in it. Use your observation to make a conclusion about each leaf. Record your conclusion in Table 1.1.

# Table 1.1

	leaf A	leaf B
colour of leaf after 2 minutes		
conclusion		

[3]

(b)	(i)	Explain why leaves <b>A</b> and <b>B</b> were placed in boiling water before being placed alcohol.	in
			 [1]
			ניו
	(ii)	Why must the Bunsen burner be turned off before the tubes containing leaves a alcohol are immersed in the water bath?	nd
			[1]
(	iii)	Explain how light has caused the difference in colour, shown by the iodine te between leaves ${\bf A}$ and ${\bf B}$ .	st,
			••••
			••••
			[3]

# **BLANK PAGE**

**2 (a)** You are going to find the cross-sectional area of a piece of resistance wire using the apparatus in Fig. 2.1 and Fig. 2.2.

For Examiner's Use




Fig. 2.1

- (i) Construct the circuit shown in Fig. 2.1. You will be connecting the crocodile clips to the resistance wire.
  - Connect one crocodile clip at the 0 cm point on the rule, with most of the length of the wire pointing along the rule. This will become the point **X** shown on Fig. 2.2.




Fig. 2.2

- Complete the circuit by pressing the other crocodile clip firmly on to the resistance wire at point Y shown in Fig. 2.2. Make sure that the length, 1, between point X and point Y is 20 cm.
- Close the switch and read the current, *I*, on the ammeter and the potential difference, *V*, on the voltmeter. Record these readings in Table 2.1. [1]
- Open the switch.

For Examiner's Use

length of wire,	potential difference, V/volts	current, I/amps	resistance, R/ohms
20			
40			
60			
80			
100			

(ii) Repeat the procedure described in (i), adjusting point **Y** so that the length, *l*, is 40 cm. Close the switch and read the current, *I*, on the ammeter and the potential difference, *V*, on the voltmeter. Record these readings in Table 2.1. [1]

Open the switch.

(iii) Repeat (i) for  $l = 60 \,\text{cm}$ ,  $80 \,\text{cm}$  and  $100 \,\text{cm}$ . For each length, l, record the current, l, and the potential difference, l, in Table 2.1. [2]

Open the switch.

(iv) Calculate the resistance, **R**, of each length of the resistance wire and enter the values in Table 2.1.


Use the formula below.

resistance, 
$$R = \frac{V}{I}$$

[1]

(b) (i) Plot a graph of resistance, **R**, (vertical axis) against length of wire, **1**. Draw a straight line of best fit. Your line should pass through the origin, (0,0).

For Examiner's Use



[2]

(ii) Find the gradient of the line and show clearly your working either on the graph or below.

gradient = [2]

(iii) Calculate the cross-sectional area, **C**, of the resistance wire using the formula given below.

For Examiner's Use

$$\mathbf{C} = \frac{0.000049}{\text{gradient}}$$

$$C = _{max} cm^2$$
 [1]

# **BLANK PAGE**

Please turn over for Question 3.

3

Solid mixture **A** contains one cation and two anions. Use the following tests to identify them. Place the mixture A in a 100 cm<sup>3</sup> beaker and add about 30 cm<sup>3</sup> of distilled or (a) (i) • deionised water. Stir the mixture well. Filter the mixture into the large test-tube for use in (b)(i) and (b)(ii). Keep the residue in the filter paper for use in (a)(ii). Describe the colour of the residue and the filtrate. residue filtrate [2] ..... (ii) Scrape a little of the residue (about one-tenth) from (a)(i) into a clean test-tube. Slowly add dilute hydrochloric acid using a teat pipette (or dropper), 1 cm<sup>3</sup> at a time, until it has all dissolved. Record all your observations and make a conclusion about the anion present in the residue. Keep this solution for (a)(iii). observations conclusion [2] (iii) To the solution from (a)(ii) add aqueous sodium hydroxide using a teat pipette, 1 cm<sup>3</sup> at a time, until there is no further change. Record your observations and conclusion about the cation in the residue from (a)(i). observations conclusion [2]

For Examiner's Use

(b)	(i)	Place about 2 cm <sup>3</sup> of the filtrate from <b>(a)(i)</b> into a clean test-tube. Add 5 drops of aqueous silver nitrate.
		Record your observations and conclusion.
		observations
		conclusion
		[2]
	(ii)	Place about 2 cm <sup>3</sup> of the filtrate from <b>(a)(i)</b> into a clean test-tube. Add about 1 cm <sup>3</sup> of aqueous barium chloride.
		Record your observations and conclusion.
		observations
		conclusion
		[0]

For Examiner's Use

#### **CHEMISTRY PRACTICAL NOTES**

#### **Test for anions**

anion	test	test result	
carbonate (CO <sub>3</sub> <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced	
chloride (Cl <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.	
nitrate (NO <sub>3</sub> <sup>-</sup> ) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced	
sulfate (SO <sub>4</sub> <sup>2-</sup> ) [in solution]	acidify then add aqueous barium chloride <i>or</i> aqueous barium nitrate	white ppt.	

## Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia	
ammonium (NH <sub>4</sub> <sup>+</sup> )	ammonia produced on warming	-	
copper(II) (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution	
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess	
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess	
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution	

### **Test for gases**

gas	test and test results	
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue	
carbon dioxide (CO <sub>2</sub> )	turns limewater milky	
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper	
hydrogen (H <sub>2</sub> )	"pops" with a lighted splint	
oxygen (O <sub>2</sub> )	relights a glowing splint	

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.