



# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

#### **CO-ORDINATED SCIENCES**

0654/51

Paper 5 Practical Test

May/June 2011

2 hours

Candidates answer on the Question Paper.

Additional Materials:

As listed in Instructions to Supervisors

#### READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.


Chemistry practical notes for this paper are printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use			
1			
2			
3			
Total			

This document consists of 12 printed pages.



[Turn over

1 You are going to compare the composition of inhaled and exhaled air.




Fig. 1.1

### (a) Analysis of inhaled air.

- Take a 500 cm<sup>3</sup> glass beaker (or glass jar).
- Light a candle (or night light) and place it onto a heatproof tile.
- Set the clock to zero.
- Place the glass beaker over the candle and start the clock (see Fig. 1.1).
- Record in Table 1.1 the time taken in seconds for the flame to go out. Remove the glass beaker from the candle, leave open to the air for one minute.
- Repeat the experiment and again record the time for the flame to go out in Table 1.1.




Fig. 1.2

## (b) Analysis of exhaled air.

- Fill the glass beaker (or glass jar) with water and place into a bowl of water, inverted (see Fig. 1.2).
- Using the straw or tube blow into the glass beaker, until it is full of exhaled air.
- Light the candle.
- Using your hand to prevent the loss of air from the glass beaker, transfer the beaker from the bowl and place it over the lit candle. Immediately start the clock.
- Record in Table 1.1 the time taken in seconds for the flame to go out.
- Repeat the experiment for exhaled air and again record the time taken for the flame to go out in Table 1.1.

© UCLES 2011 0654/51/M/J/11

	inhal	ed air	exhal	ed air
experiment number	1	2	3	4
time taken/s				

[2]

(c)	(i)	Calculate the average times for the flame to go out in inhaled air and exhaled air
		Show all working.

average time taken	for flame to go out in inhaled air	=

avaraga tima takan	for flame to	go out in exhaled air :		
average time taken	ioi liallie lo	qu dul ili exhaled ali -	- 3	•
0		0		

[3]

(ii)	Describe and air.				
		 	 	 	[2]

- (iii) Take two large test-tubes and label them A and B.
  - Place about 10 cm<sup>3</sup> of limewater into each tube.
  - Place a bung over tube A and gently shake it for 30 seconds. This will allow atmospheric air to bubble through the lime water.
  - Using a straw or tube, blow into tube **B** for 30 seconds. This will allow exhaled air to bubble through the limewater.

	Describe and explain the results observed in tube <b>A</b> and tube <b>B</b> .
	[3]
(d)	Describe an experiment to investigate the effect of exercise on breathing rate. Ensure that it is controlled, reliable and that you explain the need for all equipment.
	[5]

For Examiner's

Use

2 You are to investigate how an elastic band stretches when acted on by an increasing force.

For Examiner's Use

(a) Measure and record the width of the elastic band.

Push the pin into the cork leaving just enough showing to enable the elastic band to be hung from it. Clamp the cork firmly and adjust its position to give about 500 mm or more clearance from the bench top as shown in Fig. 2.1.




Fig. 2.1

**(b)** Hang the elastic band from the pin in such a way that masses can be attached, causing the band to stretch.

If you have a carrier for the masses, this should be attached before the initial length of the elastic band is taken. If no carrier is available, you will need to tie various masses together using thread and tie these to the band.

- (i) Measure the length of the elastic band without any masses attached. Record this length in Table 2.1.
- (ii) Attach a mass of 100 g to the band. Measure and record its new length in Table 2.1.
- (iii) Continue to add masses of 100 g until you have a total of 500 g hanging from the band. Measure the length of the band after each addition and record the values in Table 2.1.
- (iv) Complete Table 2.1 by converting each mass into a force and also calculating the total increase in length of the band.

Table 2.1


total mass/g	force/N	length of band/mm	total increase in length/mm
0	0		0
100	1.0		
200			
300			
400			
500			

[4]

Examiner's Use

(c) Plot a graph of total increase in length (vertical axis) against force (horizontal axis).

Before drawing a line, answer question (d).



[3]

(d)	Do you expect the line to pass through the origin?
	Explain your answer.
	[1]
(e)	Draw a suitable best fit line using the points plotted, taking into account your answer in <b>(d)</b> .
(f)	Use your graph to find the total increase in length produced by a mass of 250g.
	total increase in length = mm [1]
(g)	Use your graph to describe the relationship between the force applied and the total increase in length.
	[1]
(h)	A student suggested that the band may have stretched and would not return to its original form when the force is removed.
	Describe how you could test this suggestion.
	[2]

(i)	If masses were ad axes below, sketch	-		break. On the

[1]

For Examiner's Use

© UCLES 2011 0654/51/M/J/11

3

		e provided with a salt <b>Z</b> which contains two cations and one anion. Carry out the g tests.	For Examiner's Use
(a)	Red	cord the appearance of solid <b>Z</b> .	000
	арр	pearance [1]	
(b)		ce about one quarter of solid <b>Z</b> in a hard glass test-tube. Heat strongly. Test any given off with litmus paper.	
	Red	cord all your observations.	
	acti	ion on litmus paper	
	obs	servations	
		[2]	
(c)	(i)	Determine how to make a solution of <b>Z</b> . Make sure you keep some of the solid for test <b>(d)</b> .	
		Place about half of the solid <b>Z</b> into a fresh test-tube.	
		Try to dissolve ${\bf Z}$ in cold water and if it does not dissolve, try cold dilute sulfuric acid and then hot dilute sulfuric acid. <b>Stop</b> as soon as you have found a way of dissolving ${\bf Z}$ .	
		Note the colour of the solution.	
		Keep the solution for further tests. You will need about 10 cm <sup>3</sup> of solution.	
		<b>Z</b> dissolves in	
		colour of solution [2]	
	(ii)	Place about 2 cm³ of the solution from <b>(c)(i)</b> into a large test-tube. Add dilute sodium hydroxide until alkaline. Use litmus paper to check that you have added enough.	
		Record your observation.	
		observation [1]	
		Now carefully bring the mixture to the boil and test any gas given off with litmus paper.	
		Record your observation.	
		observation [1]	
			1

	(iii)	Place a second 2 cm <sup>3</sup> portion of the solution from <b>(c)(i)</b> in a fresh test-tube. Add ammonia solution until in excess.
		Record your observation.
		observation [1]
(d)		ce a small amount of solid <b>Z</b> in a fresh test-tube. Dissolve in about 5 cm <sup>3</sup> of dilute c acid. Use this solution to test for the chloride anion and the sulfate anion.
	Divi	de the solution into <b>two</b> portions <b>before</b> carrying out any tests.
	Des	scribe how you carried out each test and record the result.
	test	for the chloride anion
	resi	ult[2]
	test	for the sulfate anion
	resi	ult[2]

(e)	State th	State the three ions in solid <b>Z</b> , giving a reason for each one.		
	ion 1		E	
	reason			
	ion 2			
	reason			
	ion 3			
	reason			
		[3]		

## **CHEMISTRY PRACTICAL NOTES**

#### **Test for anions**

anion	test	test result
carbonate (CO <sub>3</sub> <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced
chloride (Cl <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
nitrate (NO <sub>3</sub> -) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2-</sup> ) [in solution]	acidify then add aqueous barium chloride <i>or</i> aqueous barium nitrate	white ppt.

# Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
ammonium (NH <sub>4</sub> <sup>+</sup> )	ammonia produced on warming	-
copper(II) (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

# **Test for gases**

gas	test and test results	
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue	
carbon dioxide (CO <sub>2</sub> )	turns limewater milky	
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper	
hydrogen (H <sub>2</sub> )	"pops" with a lighted splint	
oxygen (O <sub>2</sub> )	relights a glowing splint	

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2011 0654/51/M/J/11