

**MARK SCHEME for the October/November 2009 question paper  
for the guidance of teachers**

**0580 MATHEMATICS**

**0580/04**

Paper 4 (Extended), maximum raw mark 130

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

<b>Page 2</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2009</b>	<b>0580</b>	<b>04</b>

### Abbreviations

cao	correct answer only
cso	correct solution only
dep	dependent
ft	follow through
isw	ignore subsequent working
oe	or equivalent
SC	Special Case
soi	seen or implied
www	without wrong working

<b>1 (a) (i)</b>	<b>8.4(0)</b>	<b>B2</b>	<b>B1</b> for 1.2 or 3.6 seen or <b>SC1</b> for figs 84 in answer
<b>(ii)</b>	$\frac{\text{their(i)}}{20} \times 100$ oe <b>42</b> ft www2	<b>M1</b> <b>A1ft</b>	ft their $8.4 \times 5$ After 0 scored <b>SC1 ft</b> for 58% or $\frac{20 - \text{their(i)}}{20} \times 100$ correctly given
<b>(b)</b>	<b>6</b>	<b>B2</b>	<b>M1</b> for $9$ or $8 \div (1 + 8 + 3)$ soi
<b>(c)</b>	$\frac{2.4}{2} \times 3$ oe (= 3.6 seen) or their <b>(a) (i)</b> $\div 7 \times 3$  $\frac{3}{12} \times 9$ oe (= 2.25 seen) <b>1.6(0)</b> cao www3	<b>M1</b>  <b>M1</b> <b>A1</b>	
<b>(d)</b>	$\frac{2.40}{1.25}$ oe <b>1.92</b> www2	<b>M1</b> <b>A1</b>	Implied by figs 192   <b>[11]</b>

<b>2 (a) (i)</b>	Reflection (M), $x = 1$	<b>B1,B1</b>	If extra transformations given in part <b>(a)</b> then zero scored
<b>(ii)</b>	Rotation (R) 180 (centre) (1, 0)	<b>B1</b> <b>B1</b> <b>B1</b>	Must be "rotation". Allow half turn for 180. Allow other clear forms of (1, 0)
<b>(iii)</b>	Enlargement (E) (centre) (6, 4) (scale factor) 3	<b>B1</b> <b>B1</b> <b>B1</b>	Must be "enlargement" Allow other clear forms of (6, 4) e.g. vector Accept 3 : 1 or 1 : 3
<b>(iv)</b>	Shear (H) $y$ -axis invariant oe  (factor) -1	<b>B1</b> <b>B1</b>  <b>B1</b>	Must be "shear" Allow other explanation for invariant but not "parallel to" isw after $y$ -axis invariant seen

<b>Page 3</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2009</b>	<b>0580</b>	<b>04</b>

<b>(b) (i)</b>	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	<b>B2</b>	<b>B1</b> for correct right-hand column in 2 by 2 matrix
<b>(ii)</b>	$\begin{pmatrix} 1 & 0 \\ -1ft & 1 \end{pmatrix}$	<b>B2ft</b>	Ft only their factor in <b>(a) (iv)</b> provided not zero <b>B1ft</b> for left-hand column in 2 by 2 matrix provided shear factor is not zero or <b>SC1</b> for $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ if not ft
			<b>[15]</b>

<b>3 (a) (i)</b>	<b>1</b>	<b>B1</b>	Penalty of -1 in question if any answers given as decimals or percentages (to 3sf) alone, but isw cancelling/conversion after correct answer
<b>(ii)</b>	$\frac{3}{6}$ oe	<b>B1</b>	
<b>(b) (i)</b>	$\frac{2}{30}$ oe www2	<b>B2</b>	<b>M1</b> for $\frac{2}{6} \times \frac{1}{5}$
<b>(ii)</b>	6-12 and 12-6 and 7-11 and 11-7 soi $k \times \frac{1}{6} \times \frac{1}{5}$ for $k = \text{integer}$ $\frac{4}{30}$ oe www3	<b>M1</b> <b>M1</b> <b>A1</b>	Evidence of all pairs adding up to 18 but no extras e.g. $4/6 \times 1/6$ Without seeing the first M, $\frac{4}{6} \times \frac{1}{5}$ oe scores <b>M2</b> , $\frac{2}{6} \times \frac{1}{5}$ oe scores <b>M1</b>
<b>(iii)</b>	$\frac{4}{6} \times \frac{2}{5}$ $\frac{8}{30}$ oe www2	<b>M1</b> <b>A1</b>	
<b>(c)</b>	$\frac{2}{6} + \frac{4}{6} \times \frac{2}{5}$ oe $\frac{18}{30}$ oe cao www2	<b>M1</b> <b>A1</b>	$\frac{2}{6}$ + their <b>(b) (iii)</b>
<b>(d)</b>	<b>4</b>	<b>B2</b>	<b>M1</b> for $(1 + 1 + 6 + 7 + 11 + 12 + x) \div 7 = 6$ or better
			<b>[13]</b>

<b>Page 4</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2009</b>	<b>0580</b>	<b>04</b>

<b>4 (a) (i)</b>	Accurate triangle with 2 arcs seen, 2 mm accuracy for lines AC and BC	<b>B2</b>	<b>SC1</b> if accurate but no arcs or one arc or if AC and BC are wrong way round with arcs
<b>(ii)</b>	Accurate bisector of angle $ACB$ , $2^\circ$ accuracy and both pairs of arcs shown (accept equidistant marks on edges for 1 <sup>st</sup> set of arcs) + must meet $AB$	<b>B2ft</b>	Ft their triangle <b>SC1ft</b> if accurate but no/one pair of arcs or short with arcs In both <b>(ii)</b> and <b>(iii)</b> isw
<b>(iii)</b>	Accurate perpendicular bisector of $AD$ 2 mm accuracy at mid-point and $2^\circ$ for right angle and shows both sets of arcs + must meet $AC$	<b>B2ft</b>	ft their $D$ , which must be on $AB$ <b>SC1ft</b> if accurate but no/one pair of arcs or short with arcs
<b>(iv)</b>	Correct region shaded cao	<b>B1</b>	Dependent on correct triangle, accurate bisectors of angle $ACB$ and side $AD$ with correct $D$
<b>(b) (i)</b>	$(\cos C) = \frac{140^2 + 180^2 - 240^2}{2 \times 140 \times 180}$ oe  <b>- 0.111(1)...or better or 96.37 to 96.38</b>	<b>M2</b>  <b>E1</b>	$(-5600/50400$ or $-14/126)$ Allow use of 7, 9 and 12 <b>M1</b> for correct implicit statement Verification using 96.4 scores <b>M2</b> max Accept $-\frac{1}{9}$ but not a non-reduced fraction
<b>(ii)</b>	$0.5 \times 140 \times 180 \sin$ (their 96.4) oe <b>12521 to 12523 or 12 500 or 12520</b> cao www2	<b>M1</b> <b>A1</b>	$(s = 280)$ , allow use of 7, 9 (31.3...)
<b>(iii)</b>	$(\sin B =) \frac{140 \sin(\text{their } 96.4)}{240}$ oe  <b>35.4 or 35.42 to 35.44</b> cao www3	<b>M2</b> <b>A1</b>	Allow use of 7, 12 <b>M1</b> for correct implicit statement <b>SC2</b> for correct answer by other method

[15]

<b>5 (a) (i)</b>	$(x + 3)(2x + 5) - x(x + 4) = 59$ oe $2x^2 + 6x + 5x + 15 - x^2 - 4x = 59$ oe $x^2 + 7x - 44 = 0$	<b>M1</b> <b>A1</b> <b>E1</b>	Implies <b>M1</b> (allow $11x$ for $6x + 5x$ ) Correct conclusion – no errors or omissions
<b>(ii)</b>	$(x + 11)(x - 4)$	<b>B2</b>	<b>SC1</b> any other $(x + a)(x + b)$ where $a \times b = -44$ or $a + b = 7$
<b>(iii)</b>	<b>-11, 4</b> www ft	<b>B1ft</b>	Strict ft dep on at least <b>SC1</b> in <b>(ii)</b> allow recovery if new working seen
<b>(iv)</b>	$\tan = \frac{(\text{their} + \text{ve root}) + 3}{2(\text{their} + \text{ve root}) + 5}$ oe  <b>28.3</b> (00...) ft www2	<b>M1</b> <b>A1ft</b>	Could be alt trig method oe <b>M1</b> where trig function is explicit ft one of their positive roots $(27.4^\circ (27.40 - 27.41)$ from $x = 11)$

<b>Page 5</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2009</b>	<b>0580</b>	<b>04</b>

<b>(b) (i)</b>	$\frac{2x+5}{x+4} = \frac{x+3}{x} \quad \text{oe}$ $x^2 + 4x + 3x + 12 = 2x^2 + 5x$ $x^2 - 2x - 12 = 0$	<b>M1</b> <b>A1</b> <b>E1</b>	Must be seen. Allow ratio or correct products Correct expansion of brackets seen (allow $7x$ for $4x + 3x$ ) Correct conclusion – no errors or omissions <b>M1</b> must be seen
<b>(ii)</b>	$\frac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(-12)}}{2(1)}$ or $(x-1)^2 - 12 = 1$ (B1) and $x-1 = \pm\sqrt{13}$ (B1) <b>– 2.61, 4.61 final answers</b> www4	<b>B1,B1</b>  <b>B1,B1</b>	In square root <b>B1</b> for $(-2)^2 - 4(1)(-12)$ or better If in form $\frac{p+\sqrt{q}}{r}$ or $\frac{p-\sqrt{q}}{r}$ , <b>B1</b> for $-(-2)$ and $2(1)$ or better If B0, <b>SC1</b> for $-2.6$ and $4.6$ or <b>both</b> answers correct to 2 or more dps rot $-2.6055\dots, 4.6055\dots$
<b>(iii)</b>	<b>26.4 (26.42.... to 26.44....) ft</b>	<b>B1ft</b>	ft $4 \times$ a positive root + 8

[16]

<b>6 (a) (i)</b>	<b>–16</b>	<b>B1</b>	
<b>(ii)</b>	<b>18 to 19</b>	<b>B1</b>	
<b>(b) (i)</b>	<b>–4.3 to –4.2, 1.5 to 1.6</b>	<b>B1,B1</b>	
<b>(ii)</b>	<b>–4.5 to –4.4 , 1.3 to 1.4</b>	<b>B1,B1</b>	
<b>(iii)</b>	<b>–4.5 to –4.4 <math>&lt; x &lt; 1.3</math> to 1.4 ft</b>	<b>B1ft</b>	Ft their <b>(ii)</b> . Allow clear worded explanations and condone $\leq$ signs
<b>(c)</b>	$-\frac{30}{7}$ oe isw conversion	<b>B2</b>	Accept $-4\frac{2}{7}, 30/-7$ <b>M1</b> for $30/7$ oe fract, isw conversion <b>or</b> for $-30/7$ oe soi
<b>(d)</b>	Ruled line passing within 2 mm of $(-5, 30)$ and $(2, 0)$	<b>B2</b>	<b>B1</b> for ruled line parallel to $g(x)$ . By eye ( $21^\circ$ to $25^\circ$ to horizontal if in doubt) allow broken line
<b>(e) (i)</b>	Ruled horizontal line through $(-3, -27)$	<b>B1</b>	No daylight, not chord (allow broken)
<b>(ii)</b>	$y = -27$	<b>B1</b>	
<b>(f)</b>	Ruled lines $x = -3, x = -2, y = 40$  Region enclosed by lines $x = -3, x = -2, y = 40$ and $y = g(x)$	<b>B1</b>  <b>B1</b>	Long enough to be boundary of region – allow broken or solid ruled lines  Allow any clear indication

[15]

<b>Page 6</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2009</b>	<b>0580</b>	<b>04</b>

<b>7 (a) (i)</b>	$\frac{60}{360} \times \pi \times 2 \times 24$ oe <b>25.1 (25.12 to 25.14)</b> www2	<b>M1</b> <b>A1</b>	Accept $8\pi$
<b>(ii)</b>	$\frac{60}{360} \times \pi \times 24^2$ oe <b>301 or 302 or 301.4 to 301.7</b> www2	<b>M1</b> <b>A1</b>	Accept $96\pi$
<b>(b) (i)</b>	$\pi d = \text{their (a) (i)}$ oe <b>4 (3.99 – 4.01)</b> cao www2	<b>M1</b> <b>A1</b>	
<b>(ii)</b>	$24^2 - (\text{their radius})^2$ <b>23.7 (23.66 to 23.67)</b> cao www2	<b>M1</b> <b>A1</b>	Alt trig method for $h$ explicit Accept $\sqrt{560}, 2\sqrt{140}, 4\sqrt{35}$
<b>(iii)</b>	$\frac{1}{3} \times \pi \times (\text{their } r)^2 \times (\text{their } h)$ <b>394 – 398</b> cao www2	<b>M1</b> <b>A1</b>	Not for $h = 24$
<b>(c) (i)</b>	<b>27W</b>	<b>B1</b>	
<b>(ii)</b>	<b>4W</b>	<b>B1</b>	If B0, B0 in (c), <b>SC1</b> for 27 and 4 alone <b>[12]</b>

<b>8 (a)</b>	<b><math>5.5 &lt; t \leq 6</math></b>	<b>B1</b>	Condone poor notation
<b>(b)</b>	4.25, 4.75, 5.25, 5.75, 6.25, 6.75 $(2 \times 4.25 + 7 \times 4.75 + 8 \times 5.25 + 18 \times 5.75 + 10 \times 6.25 + 5 \times 6.75)$ (= 283.5) $\div 50$ or their $\sum f$ <b>5.67</b> www4	<b>M1</b> <b>M1</b> <b>M1</b> <b>A1</b>	At least 5 correct mid-values seen $\sum fx$ where $x$ is in the correct interval allow one further slip Depend on second method After <b>M3</b> allow 5.7 isw conversion to mins/secs and reference to classes
<b>(c) (i)</b>	<b>17, 15</b>	<b>B1</b>	
<b>(ii)</b>	Rectangular bars of heights <b>11.3</b> and <b>15</b>  Correct widths of 1.5 and 1 – no gaps	<b>B1ft</b> <b>B1ft</b>  <b>B1</b>	ft their 17 divided by 1.5 ft their 15 11.3 plot between 11 and 12 include lines and 15 to be touching the 15 line
<b>(iii)</b>	<b>2.5</b> cao	<b>B1</b>	<b>[10]</b>

<b>Page 7</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2009</b>	<b>0580</b>	<b>04</b>

<b>9 (a)</b>	$3(m-3) + 4(m+4) = -7 \times 12$ $3m - 9 + 4m + 16 = -84$ <b>-13</b> www4	<b>M2</b> <b>A1</b> <b>A1</b>	Allow <u>all over</u> 12 at this stage <b>M1</b> for $3(m-3) + 4(m+4)$ seen Allow <u>all over</u> 12 at this stage May be seen in stages
<b>(b) (i)</b>	0.5 oe	<b>B1</b>	
<b>(ii)</b>	$\frac{3(x+3) - 2(x-1)}{(x-1)(x+3)}$ $\frac{x+11}{(x-1)(x+3)}$ final answer	<b>M1</b> <b>A1</b>	If brackets not seen allow $3x + 9 - 2x \pm 2$ as numerator with a correct denominator <b>A1</b> isw incorrect expansion of denominator if correct brackets seen
<b>(iii)</b>	$\frac{x(x+11)}{(x-1)(x+3)} = 1$ ft or $x + 11 = \frac{1}{x} (x-1)(x+3)$ or better ft $x^2 + 11x = x^2 + 3x - x - 3$ $-\frac{1}{3}$ oe cso www3	<b>M1</b> <b>M1</b> <b>A1</b>	Must clear one denominator correctly Ft their <b>(b)(ii)</b> dep on fraction in <b>(ii)</b> with $(x-1)(x+3)$ oe as denominator Depend on previous <b>M1</b> - 0.33(33...)
<b>(c)</b>	$p(q-1) = t$ oe $pq = t + p$ $\frac{t+p}{p}$ oe final answer www3	<b>M1</b> <b>M1</b> <b>M1</b>	Multiplying by $(q-1)$ Ft their first step e.g. $pq$ only term on one side Ft their 2 <sup>nd</sup> step e.g. dividing by $p$ Note: $q-1 = \frac{t}{p}$ is <b>M2</b> and then $q = \frac{t}{p} + 1$ is <b>M1</b>

[13]

<b>10 (a)</b>	$21 + 23 + 25 + 27 + 29 = 125$ $31 + 33 + 35 + 37 + 39 + 41 = 216$	<b>B1</b> <b>B1</b>	
<b>(b)</b>	Cubes	<b>B1</b>	
<b>(c) (i)</b>	$n$ oe	<b>B1</b>	
<b>(ii)</b>	$n^3$ oe	<b>B1</b>	
<b>(d)</b>	$4^2 - 4 + 1 = 13$ www	<b>E1</b>	Allow 16 for $4^2$ , otherwise all must be seen
<b>(e)</b>	$7 \times 43 + 2 + 4 + 6 + 8 + 10 + 12$	<b>B1</b>	All must be seen
<b>(f)</b>	$n(n-1)$ final answer oe	<b>B1</b>	
<b>(g)</b>	$n(n^2 - n + 1)$ + their <b>(f)</b> $n^3 - n^2 + n + n^2 - n = n^3$	<b>M1</b> <b>E1</b>	All must be seen, no errors or omissions

[10]