



Pearson  
Edexcel

Mark Scheme (Results)

Summer 2022

Pearson Edexcel International GCSE

In Science Double Award (4SD0) Paper 1CR

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2022

Question Paper Log Number P70946A

Publications Code 4SD0\_1CR\_2206\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2022

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.



Question number	Answer		Notes	Marks
2 (a)	<b>Apparatus</b>	<b>Unit</b>	<b>ALLOW</b> ml <b>ALLOW</b> words e.g. grams <b>ALLOW</b> weighing machine <b>ALLOW</b> kg  1 mark each correct row/column Mark horizontally or vertically (whichever benefits the candidate)	2
	(gas) syringe	cm <sup>3</sup>		
	(top pan) balance <b>OR</b> (weighing) scales	g <b>OR</b> mg		
(b) (i)	M1 to cool (the water vapour/steam)		<b>ALLOW</b> to keep condenser cool	2
	M2 so the water vapour/steam condenses		<b>ALLOW</b> so the water vapour/steam becomes liquid	
	(ii)	M1 add silver nitrate/AgNO <sub>3</sub> (solution)		<b>IGNORE</b> addition of nitric acid but <b>REJECT</b> addition of hydrochloric/sulfuric acid for M1  M2 dep on use of silver nitrate
(iii)	M1 measure its boiling point		<b>ALLOW</b> boil it	2
	M2 (boiling point is) 100° C			
	OR			
	M1 measure its freezing point		<b>ALLOW</b> freeze it	
	M2 (freezing point is) 0° C			

Total for question 2 = 8 marks

Question number	Answer	Notes	Marks
3 (a) (i)	B		1
(ii)	A and B		1
(iii)	M1 2 and 8  M2 0.25	0.25 without working scores 2  <b>ALLOW</b> M1 for 1.8-2.2 and 8 and <b>ALLOW</b> M2 ECF as long as correctly evaluated to at least 2 SF  (Special case if used ruler and then) 1.4-1.7 and 5.9-6.2 used no M1 but <b>ALLOW</b> M2 ECF as long as correctly evaluated to at least 2 SF	2
(iv)	the dye is the most soluble (in the solvent/water)		1
(b)	Any four from  M1 draw start line in pencil  M2 use same food colourings/use same solvent/use same (type of chromatography) paper  M3 place (spots/samples of) A, B, C, D/food colourings on the start line OWTTE  M4 (place paper in beaker) with start line above solvent OWTTE  M5 (remove paper/stop experiment) when solvent almost reaches top of paper / when spots stop moving OWTTE  M6 mark solvent front (on paper)  M7 (remove paper from beaker and) allow to dry	<b>ALLOW</b> water for solvent throughout <b>ALLOW</b> dye for food colouring throughout  <b>IGNORE</b> length of paper	4

Total for question 3 = 9 marks

Question number	Answer	Notes	Marks
4 (a)	(i) Any one from: Na K Al In	<b>ALLOW</b> names of elements Apply list principle	1
	(ii) Any one from S Cl	<b>ALLOW</b> names of elements	1
(b)	same number / three electrons in the outer shell	<b>ALLOW</b> valence shell	1
(c)	M1 Xe or xenon  M2 as it has a full outer shell (of electrons)	<b>ALLOW</b> has eight electrons in outer shell <b>ACCEPT</b> does not (easily) gain/lose/share electrons M2 dep on M1	2
(d)	(i) M1 (universal indicator turns) blue or purple  M2 because an alkali is produced	<b>ACCEPT</b> OH <sup>-</sup> / hydroxide ions are produced <b>ALLOW</b> sodium hydroxide is a base / a base is produced  <b>ALLOW</b> fizzes/bubbles  <b>ALLOW</b> float <b>ALLOW</b> both disappear/get smaller/dissolve  <b>ALLOW</b> faster/more vigorous reaction for potassium  <b>ALLOW</b> reverse arguments for sodium	2
	(ii) (similarity) any one from: (both) effervesce melt / turn into a sphere move on surface universal indicator turns the same colour  (difference) any one from: potassium gives a lilac flame potassium moves faster potassium effervesces faster		2
	(iii) Example calculation  M1 (moles of hydrogen) $0.036 \div 2$ <b>OR</b> 0.018 mol  M2 $0.018 \times 6.0 \times 10^{23}$ <b>OR</b> $1.08 \times 10^{22}$ molecules  M3 $1.1 \times 10^{22}$		3
		correct answer with no working scores 3 marks  <b>ALLOW</b> ECF M1 $\times 6.0 \times 10^{23}$  <b>ALLOW</b> ECF M2 but must be to 2 sig figs  $2.16 \times 10^{22}$ scores 1 $2.2 \times 10^{22}$ scores 2	

Total for question 4 = 12 marks

Question number	Answer	Notes	Marks
5 (a)	<p>M1 (put the carbonate in the boiling tube) and the limewater in the test tube</p> <p>M2 heat the carbonate and time how long it takes for the limewater to turn cloudy OWTTE</p> <p>M3 repeat with the same mass / amount / number of moles of another carbonate</p> <p>M4 (the carbonate which decomposes the fastest) will turn the limewater cloudy in the least time</p>	<p><b>ACCEPT</b> repeat with another carbonate using same volume of limewater OWTTE</p> <p>To score M4 reference to limewater turning cloudy must be mentioned at least once somewhere in answer</p>	4
(b) (i)	to prevent loss of solid/ $XCO_3$ /carbonate/XO	<b>ALLOW</b> so <u>only</u> carbon dioxide/gas can escape	1
(ii)	0.05		1
(iii)	0.05	<b>ALLOW</b> ECF from (ii)	1
(iv)	<p>M1 <math>7.40 \div 0.05</math></p> <p>M2 148</p>	<p>correct answer with or without working scores 2</p> <p><b>ALLOW</b> ECF from (iii)</p>	2
(v)	<p>M1 <math>A_r</math> of metal = 148-60 <b>OR</b> 88</p> <p>M2 metal is strontium / Sr</p>	<p>If (iv) correct strontium/Sr scores 2 without working</p> <p><b>ALLOW</b> ECF from (iv)</p> <p><b>ALLOW</b> ECF from M1 as long as answer is nearest Group 2 metal</p>	2


Total for question 5 = 11 marks



Question number	Answer	Notes	Marks
6 (a)	M1 shared pair(s) of electrons M2 attracted to (two) nuclei	<b>REJECT</b> nucleus. Must be plural for M2 M2 dep on mention of electrons in M1	2
(b)	a pair of electrons in each bond and no non-bonding electrons.	<b>ALLOW</b> dots, crosses or any combination	1
(c) (i)	Any one from M1 oxygen is a smaller atom/particle than silicon M2 each (atom of) oxygen forms two bonds (to silicon atoms)		1
(ii)	M1 silicon dioxide has a giant (covalent) structure M2 (in melting silicon dioxide) strong/many covalent bonds (need to be broken) M3 (in melting silicon hydride) weak intermolecular forces (of attraction need to be overcome/broken) M4 more (thermal/heat) energy is needed to break the (covalent) bonds (in SiO <sub>2</sub> ) than break/overcome the intermolecular forces (in SiH <sub>4</sub> )	<b>ALLOW</b> description of covalent bonds as long as strong/many mentioned  <b>ALLOW</b> weak intermolecular bonds  Max 2 if contradictions/references to incorrect forces/particles	4
(d)	SiH <sub>4</sub> + 2O <sub>2</sub> → SiO <sub>2</sub> + 2H <sub>2</sub> O all formula correct and equation correctly balanced	<b>IGNORE</b> state symbols <b>ALLOW</b> multiples and fractions	1

Total for question 6 = 9 marks

Question number	Answer	Notes	Marks
7 (a)	<p>Any five from</p> <p>M1 fractional distillation</p> <p>M2 crude oil heated/vapourised</p> <p>M3 reference to (fractionating) column/tower</p> <p>M4 which is hotter at the bottom than at the top</p> <p>M5 shorter hydrocarbons/chains/molecules have lower boiling point (and rise higher/towards the top)</p> <p>M6 fractions/hydrocarbons/gases/vapours/kerosene condense(s) at (levels depending on) their boiling points OWTTE</p>	<p><b>ALLOW</b> boiled</p> <p><b>ALLOW</b> reference to temperature gradient <b>ALLOW</b> the hydrocarbons/gases/vapours cool as they rise up the column</p> <p><b>ACCEPT</b> reverse argument</p> <p><b>ALLOW</b> correct reference to position of kerosene fraction below refinery gases and gasoline fractions or above diesel and fuel oil fractions</p>	5
(b) (i)	<p><math>C_8H_{18}</math></p>		1
(b) (ii)	<p>Any four from</p> <p>M1 fractional distillation/crude oil produces more long-chain hydrocarbons than can be used (directly)</p> <p>M2 cracking produces short chain alkanes</p> <p>M3 short chain alkanes/hydrocarbons are more flammable/can be used as fuels</p> <p>M4 cracking produces alkene(s)</p> <p>M5 alkenes can be used to make polymers</p>	<p><b>ALLOW</b> short(er) chain hydrocarbons are in higher demand/more useful than long(er) chain hydrocarbons ORA</p> <p><b>ALLOW</b> cracking changes long(er) chain hydrocarbons into short(er) chain hydrocarbons</p> <p><b>IGNORE</b> named alkene</p> <p><b>ALLOW</b> named alkene forming (named) polymer e.g. ethene can be used to make poly(ethene)/polymer</p>	4

(c)	<p>A Addition  The only correct answer is A because the reaction between an alkene and a halogen forming a halogenoalkane is addition.  B is not the correct answer since this reaction is not combustion.  C is not the correct answer since this reaction is not decomposition.  D is not the correct answer since this reaction is not substitution.</p>		1
(d) (i)	 <p>M1 single bond between the two carbons <b>and</b> single bonds to three hydrogens and one chlorine  M2 two extension bonds and n</p>	<p>n can be anywhere after brackets  extension bonds do not have to go through brackets  M2 dep on M1</p>	2
(ii)	<p>M1 they are inert/unreactive  M2 (so) they are non-biodegradable/ do not (naturally) break down/decompose (in landfill sites)  OR  M1 when burned  M2 they produce toxic fumes</p>	<p><b>ALLOW</b> take long time to break down (so landfill sites may fill up)</p>	2

Total for question 7 = 15 marks

Question number	Answer	Notes	Marks
8 (a) (i)	zinc + copper(II) sulfate → <b>zinc sulfate + copper</b>	<b>REJECT</b> copper(II) as a product <b>ACCEPT</b> zinc(II) sulfate	1
(ii)	polystyrene is an insulator / prevents heat loss	<b>ALLOW</b> to prevent the cup falling over	1
(iii)	zinc is more reactive / higher in the reactivity series than copper	<b>ALLOW</b> reverse argument	1
(iv)	M1 temperature rise = 28.6 °C  M2 (75 × 4.2 × 28.6 =) 9009	correct answer without working scores 2  ALLOW 9010/9000  <b>IGNORE</b> any sign <b>ALLOW</b> ECF from M1	2
(b) (i)	M1 800 ÷ 1000 OR 0.8 (kJ) M2 0.65 ÷ 65 OR 0.01 (mol) M3 0.8 ÷ 0.01 = -80 (kJ/mol) OR M1 0.65 ÷ 65 OR 0.01 (mol) M2 800 ÷ 0.01 OR 80 000 (J/mol) M3 -80 (kJ/mol)	correct answer without working scores 3          <b>ALLOW</b> ECF but answer must have a - sign  80/+80 scores 2	3
(ii)	M1 zinc/Zn is oxidised because loses electrons M2 silver ion(s)/Ag <sup>+</sup> reduced because gain electrons OR M1 zinc/Zn is oxidised <b>and</b> silver ion(s)/Ag <sup>+</sup> reduced M2 zinc/Zn loses electrons <b>and</b> silver ion(s)/Ag <sup>+</sup> gain electrons		2

Total for question 8 = 10 marks

Question number	Answer	Notes	Marks
9 (a) (i)	M1 copper(II) sulfate (solution) M2 shortest time taken to turn colourless	<b>ALLOW</b> copper sulfate  <b>ALLOW</b> gave greatest increase in rate OWTTE <b>ALLOW</b> made reaction happen fastest OWTTE  M2 dep on M1	2
(ii)	M1 a catalyst provides an alternative pathway M2 of lower activation energy	Any reference to increasing energy/speed of particles scores 0	2
(b) (i)	An explanation with following four points  <b>M1</b> the rate of reaction increases/ the reaction is faster/ the reaction speeds up  <b>M2</b> because the particles gain (kinetic) energy / move faster  <b>M3</b> there are more collisions per unit time  <b>M4</b> more of the collisions are successful / more collisions/particles have energy greater than the activation energy	<b>ACCEPT</b> more frequent collisions OWTTE  No M4 if refer to lower activation energy  there are more frequent successful collisions scores <b>M3</b> and <b>M4</b>	4
(ii)	M1 fewer particles per unit volume  M2 (hence) fewer collisions per unit time	<b>ALLOW</b> particles less tightly packed / particles further apart  <b>ALLOW</b> decrease in the frequency of collisions between particles  Any reference to changing energy/speed of particles scores 0	2

Total for question 9 = 10 marks

Question number	Answer	Notes	Marks
10 (a) (i)	measuring cylinder / burette / pipette	<b>ALLOW</b> syringe	1
(ii)	<b>M1</b> and <b>M2</b> all the points correct $\pm$ half a square	If only one plotting error scores <b>M1</b>	2
(iii)	2 straight lines of best fit, ignoring the anomalous point	Left line does not have to go through/use (0.0, 10.0) if point has not been plotted	1
(iv)	as the volume of sulfuric acid increases the (electrical) conductivity decreases	<b>IGNORE</b> references to gradient/slope/correlation	1
(v)	(the student) forgot to stir the mixture	<b>ALLOW</b> any reference to adding less acid/lower volume (than should have done) OWTTE	1
(b) (i)	M1 barium sulfate has a (giant) ionic structure <b>OR</b> has ionic bonding  M2 ionic substances do not conduct when solid  M3 water has covalent bonding <b>and</b> covalent compounds do not (usually) conduct electricity	<b>ALLOW</b> only conduct when dissolved/molten <b>ALLOW</b> in solid ions cannot move  <b>ALLOW</b> water does not conduct because it is covalent  <b>IGNORE</b> explanations of why covalent do not conduct	3
(ii)	filtration <b>OR</b> filtering		1

Total for question 10 = 10 marks

Question number	Answer	Notes	Marks
11 (a) (i)	M1 $\text{WO}_3(\text{s}) + 3\text{H}_2(\text{g})$ M2 $\text{W}(\text{s}) + 3\text{H}_2\text{O}(\text{g or l})$	<b>ALLOW</b> upper case	2
(ii)	heat again to constant mass OWTTE		1
(iii)	M1 (mass of tungsten =) 1.84g AND (mass of oxygen =) 0.48g M2 (moles of tungsten) = $\frac{1.84}{184}$ or 0.01 AND (moles of oxygen) = $\frac{0.48}{16}$ or 0.03 M3 therefore ratio is 1:3	M2 subsumes M1 <b>ALLOW</b> M2 ECF from incorrect masses  M3 dep on M2 <b>ALLOW</b> ECF from incorrect M2 only if does give 1:3 when rounded	3
(iv)	Any one from M1 use a safety screen M2 position the class some distance from the apparatus OWTTE M3 do the experiment in a fume cupboard	<b>ALLOW</b> heat proof/safety gloves <b>ALLOW</b> tie back hair	1
(b)	Example calculation M1 moles of tungsten oxide = $(2784 \times 10^6 \div 232) = 12\,000\,000$ M2 maximum mass of tungsten = $(12\,000\,000 \times 184) = 2\,208\,000\,000$ g <b>OR</b> 2208 tonnes M3 mass of tungsten (considering 73.5% yield) = $(73.5 \times 2208 \div 100) = 1622.88$ (tonnes)	correct answer without working scores 3 <b>ALLOW</b> any number of significant figures $\geq 2$ throughout <b>ALLOW</b> other correct methods <b>ALLOW</b> working in megamoles  <b>ALLOW</b> ECF $M1 \times 184$  <b>ALLOW</b> ECF from M2	3

Total for question 11 = 10 marks

