

MODEL SOLUTIONS TO IIT JEE 2009**Paper I CODE 0****PART I**

1	2	3	4	5	6	7	8
B	B	C	A	B	A	D	B
9	10	11	12				
B, C	C, D	A, C, D	A, D				
13	14	15	16	17	18		
D	C	B	B	A	B		
19	20						
A – p, q, r, t	A – q, s, t						
B – q, r, s, t	B – s, t						
C – p, q, r	C – p						
D – p, q, r, s	D – r						

Section I

- Atomic mass of Fe

$$= \frac{(54 \times 5) + (56 \times 90) + (57 \times 5)}{100}$$

$$= 55.95$$
- $\frac{an^2}{v^2}$ is the term that corrects for the attractive forces present in a real gas in the van der Waals equation.
- Sb_2S_3 sol is negatively charged.
 \therefore The most effective coagulating agent among the given is $Al_2(SO_4)_3$ due to the highest charge on the cation (Al^{3+}).
- $P_2 = Kx_2$
 $5 \times 0.8 \text{ atm} = 1 \times 10^5 \text{ atm} \times x_2$
 $x_2 = 4 \times 10^{-5}$
Mole fraction of N_2 dissolved in 10 moles of water
 $= 4 \times 10^{-5} \times 10$
 $= 4 \times 10^{-4}$
- P_4O_6 is formed when P_4 is burnt in a limited supply of air. O_2 diluted with N_2 produces that condition.
- Carboxylic acids are more acidic than phenols. Presence of electron donating groups such as $-CH_3$ group decreases the acid strength of carboxylic acids. Presence of electron withdrawing group such as $-Cl$ increases the acid strength of phenol.
- Natural rubber is an elastomer. The intermolecular force of attraction is the weakest for elastomers.
- $-CN$ group has higher priority over $-OH$ and $-Br$ which are given in alphabetical order.

Section II

- Frenkel defect is favoured by a large difference in sizes of cation and anion. It is a dislocation

effect. Trapping of electrons in lattice sites leads to the formation of F-centres. Schottky defects have effect on the physical properties of solids.

10. $[\text{Pt}(\text{en})_2\text{Cl}_2]\text{Cl}_2$ and $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$ exhibit geometrical isomerism.

11. In excess air Na_2O_2 is the main product. Small amount of NaO_2 is formed which is responsible for the yellow colour of Na_2O_2 .

Air always contains moisture which produces small amounts of NaOH .

12. (A) Total number of stereo isomers is 6
 cis d, l and cis l, d (enantiomers),
 trans d, l and trans l, d (enantiomers),
 cis d, d (same as cis l, l) meso (plane of symmetry),
 trans d, d (same as trans l, l) meso (centre of symmetry)

(D) Two enantiomers are possible
 cis d, l and its mirror image cis l, d

Section III





13. Na_2S

Na_2S forms a sulphur bridge in two p-amino-N,N-dimethyl aniline.








14. FeCl_3

FeCl_3 oxidises the above compound to methylene blue

15. $\text{Fe}^{3+} + [\text{Fe}(\text{CN})_6]^{3-} \rightarrow \text{Fe}[\text{Fe}(\text{CN})_6]$



The complete reaction is



Section IV

19. (A) (p) By MOT B_2 is paramagnetic
(q) Boron can be burned to B_2O_3
(r) Boron can be reduced with metals to form metal borides.
(t) In B_2 molecule by MOT 2s and 2p orbitals mix to bring the energy of $\sigma 2p_z$ above that of $\pi 2p_x$ and $\pi 2p_y$ (It is equivalent to say that $\sigma 2p_z$ and $\sigma^* 2s$ interact to bring $\sigma 2p_z$ above the $\pi 2p_x$ and $\pi 2p_y$).
- (B) (q) N_2 can be oxidised to NO by air.
(r) N_2 undergoes reduction to NH_3 .
(s) Bond order in N_2 is 3.
(t) In N_2 molecule also there is mixing of 2s and 2p as in the above case of B_2 .
- (C) (p) O_2^- is paramagnetic by MOT.
(q) O_2^- can be oxidised to O_2 .
(r) O_2^- can be reduced to O_2^{2-} .


- (D) (p) By MOT O_2 is paramagnetic.
(q) O_2 can be oxidised to OF_2 .
(r) O_2 can be reduced to CaO.
(s) Bond order in O_2 is 2.

20. (A) \rightarrow q, s, t
(B) \rightarrow s, t
(C) \rightarrow p
(D) \rightarrow r

Reduction of cyanides with $SnCl_2 / HCl$ or DIBAL-H followed by hydrolysis gives corresponding aldehydes. Cyanides can undergo alkaline hydrolysis to form sodium salt of carboxylic acid and NH_3 . DIBAL -H reduces esters to aldehydes.

Esters can be catalytically reduced to alcohols and they undergo alkaline hydrolysis.

Double bonds undergo catalytic reduction . Primary amines undergo Hofmann's carbylamine reaction with $CHCl_3$ and alcoholic KOH.



PART II

21 22 23 24 25 26 27 28
A B C A D D D C

29 30 31 32
B, C, D A, C B, C B, A


33 34 35 36 37 38
A B B A B D

39	40
A – p, q, s	A – p
B – p, t	B – s, t
C – p, q, r, t	C – r
D – s	D – q, s

Section I

21. $\frac{x-1}{-3} = \frac{y+1}{1} = \frac{z-2}{5} = \mu$
 Q(-3μ + 1, μ - 1, 5μ + 2)
 P(3, 2, 6)
 $\vec{PQ} = [-3\mu - 2, \mu - 3, 5\mu - 4]$
 $[1, -4, 3]$
 $-3\mu - 2 - 4\mu + 12 + 15\mu - 12 = 0$
 $8\mu - 2 = 0 \Rightarrow \mu = \frac{1}{4}$

22.



$$r = \sqrt{3^2 + 2^2 + 11} = \sqrt{24}$$

Equation of AB is

$$x \times 1 + y \times 8 - 3(x + 1) - 2(y + 8) - 11 = 0$$

$$x + 8y - 3x - 3 - 2y - 16 - 11 = 0$$

$$-2x + 6y - 30 = 0$$

$$x - 3y + 15 = 0$$

Let the circle be

$$x^2 + y^2 - 6x - 4y - 11 + \lambda(x - 3y + 15) = 0$$

It passes through (1, 8)

$$1 + 64 - 6 - 32 - 11 + \lambda(1 - 24 + 15) = 0$$

$$16 - 8\lambda = 0$$

$$\lambda = 2$$

$$x^2 + y^2 - 6x - 4y - 11 + 2(x - 3y + 15) = 0$$

$$x^2 + y^2 - 4x - 10y + 19 = 0$$

23. $\int_0^x \sqrt{1 - (f'(t))^2} dt = \int_0^x f(t) dt$

Differentiating w.r.t. x :

$$\sqrt{1 - \left(\frac{dy}{dx}\right)^2} = f(x)$$

$$y^2 = 1 - \left(\frac{dy}{dx}\right)^2$$

$$\left(\frac{dy}{dx}\right)^2 = 1 - y^2$$

$$\frac{dy}{dx} = \pm \sqrt{1 - y^2}$$

$$\pm \frac{dy}{\sqrt{1 - y^2}} = dx$$

Integrating ,

$$(+)\ \sin^{-1} y = x + C$$

$$0 = 0 + C \Rightarrow C = 0$$

$$y = \sin x$$

$$(-)\ \cos^{-1} y = x + C$$

$$\text{But } \frac{\pi}{2} = 0 + C$$

$$\therefore \cos^{-1} y = x + \frac{\pi}{2}$$

$$y = \cos\left(x + \frac{\pi}{2}\right)$$

$$= -\sin x$$

But $f(x)$ is non negative in $[0, 1]$

$$\therefore f(x) = \sin x$$

$$\left. \begin{aligned} \sin \frac{1}{2} &< \frac{1}{2} \\ \sin \frac{1}{3} &< \frac{1}{3} \end{aligned} \right\}$$

$$24. (\bar{z}\bar{z})(\bar{z})^2 + (\bar{z}z)z^2 = 350$$

$$|z|^2 (z^2 + \bar{z}^2) = 350$$

$$(x^2 + y^2) \{2(x^2 - y^2)\} = 350$$

$$x^4 - y^4 = 175$$


$$(x^2 + y^2)(x^2 - y^2) = 175$$

$$x^2 = 16 \Rightarrow x = \pm 4$$

$$y^2 = 9 \Rightarrow y = \pm 3$$

$$\therefore \text{Area of the rectangle} = 8 \times 6 = 48$$

25.



$$\frac{x^2}{9} + \frac{y^2}{1} = 0$$

Auxiliary O is $x^2 + y^2 = 9$

$$A(3, 0)$$

$$B(0, 1)$$

$$\text{Slope of AB} = -\frac{1}{3}$$

$$y = -\frac{1}{3}(x - 3)$$

$$3y = -x + 3$$

$$y = \frac{-x}{3} + 1$$

$$x^2 + \left(\frac{-x}{3} + 1\right)^2 = 9$$

$$x^2 + \frac{x^2}{9} + 1 - \frac{2x}{3} = 9$$

$$9x^2 + x^2 + 9 - 6x = 81$$

$$10x^2 - 6x - 72 = 0$$

$$5x^2 - 3x - 36 = 0$$

$$x = \frac{3 \pm \sqrt{9 + 720}}{10} = \frac{3 \pm 27}{10}$$

$$= 3, -\frac{12}{5}$$

$$y = \frac{-12}{5x - 3} + 1$$

$$= \frac{4}{5} + 1 = \frac{9}{5}$$

$$\text{Area OAM} = \frac{27}{5} \times \frac{1}{2} = \frac{27}{10}$$

$$26. \text{ Given } (\bar{a} \times \bar{b}) \cdot (\bar{c} \times \bar{d}) = 1$$

$|\bar{a} \times \bar{b}| |\bar{c} \times \bar{d}| \cos \gamma = 1$ where γ is the angle between $(\bar{a} \times \bar{b})$ and $(\bar{c} \times \bar{d})$

$$\Rightarrow \sin \alpha \sin \beta \cos \gamma = 1 \quad (\text{since}$$

$$|\bar{a}| = |\bar{b}| = |\bar{c}| = |\bar{d}| = 1 \text{ and we assume that angle}$$

between \bar{a} and \bar{b} is α and that, the angle

between \bar{c} and \bar{d} is β)

$$\Rightarrow \sin \alpha = 1, \sin \beta = 1, \cos \gamma = 1$$

$$\Rightarrow \alpha = \beta = \frac{\pi}{2}, \gamma = 0$$

$\Rightarrow \bar{a}$ and \bar{b} are orthogonal; \bar{c} and \bar{d} are

orthogonal; $\bar{a} \times \bar{b}$ is parallel to $\bar{c} \times \bar{d}$.

$\Rightarrow \bar{a}, \bar{b}, \bar{a} \times \bar{b}$ form a mutually orthogonal triad

$\bar{c}, \bar{d}, \bar{c} \times \bar{d}$ form a mutually orthogonal triad

since $(\bar{a} \times \bar{b})$ is parallel to $\bar{c} \times \bar{d}$,

we have the choices

\bar{a} is parallel to \bar{d} and \bar{b} is parallel to \bar{c}

Note that \bar{a} cannot be parallel to \bar{c} since

$$\bar{a} \cdot \bar{c} = \frac{1}{2}$$

$$27. \sum_{m=1}^{15} \text{Im } z^{2m-1} = \sin \theta + \sin 3\theta + \sin 5\theta + \dots + \sin 29\theta$$

We have $\sin \alpha + \sin(\alpha + \beta) + \sin(\alpha + 2\beta) + \dots + \sin(\alpha + n - 1\beta)$

$$= \frac{\sin\left(\frac{\alpha + \alpha + n - 1\beta}{2}\right) \sin\left(\frac{n\beta}{2}\right)}{\sin \frac{\beta}{2}}$$

Here $\beta = 2\theta$

$$\therefore \sin \theta + \sin 3\theta + \dots + \sin 29\theta$$

$$= \frac{\sin\left(\frac{\theta + \theta + 14 \times 2\theta}{2}\right) \sin\left(\frac{15 \times 2\theta}{2}\right)}{\sin \frac{2\theta}{2}}$$

$$= \frac{\sin^2 15\theta}{\sin \theta}$$

$$= \frac{\sin^2 30^\circ}{\sin 2^\circ} = \frac{1}{4 \sin 2^\circ}$$

$$28. x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 = 10$$


$$(x + x^2 + x^3)^7 = x^7(1 + x + x^2)^7$$

Coefficient of x^3 in $(1 + x + x^2)^7$

$$\begin{aligned}
 &= \text{Coefficient of } x^3 \text{ in } \frac{(1-x^3)^7}{(1-x)^7} \\
 &= \text{Coefficient of } x^3 \text{ in } (1-x^3)^7 (1-x)^{-7} \\
 &= \frac{7 \cdot 8 \cdot 9}{1 \cdot 2 \cdot 3} - 7 \times 1 \\
 &= 84 - 7 = 77
 \end{aligned}$$

Section II

29.



Required area = area of the region ABC
= Area OCBD - Area OABD

$$= e \times 1 - \int_0^1 e^x dx$$

$$= e - \int_0^1 e^x dx$$

$$= e - (e - 1) = 1$$

$$\int_1^e \ln y dy = [y \log y - y]_1^e$$

$$= (e - e) - (0 - 1)$$

$$= 1$$

$$\int_1^e \ln y dy = \int_1^e \ln(1+e-y) dy$$

30. $L = \lim_{x \rightarrow 0} \frac{a - \sqrt{a^2 - x^2} - \frac{x^2}{4}}{x^4} \quad (a > 0) \left(\frac{0}{0} \text{ form} \right)$

$$= \lim_{x \rightarrow 0} \frac{\left(-\frac{1}{2} \right) \frac{(-2x)}{\sqrt{a^2 - x^2}} - \frac{x}{2}}{4x^3}$$

$$= \lim_{x \rightarrow 0} \frac{\frac{1}{\sqrt{a^2 - x^2}} - \frac{1}{2}}{4x^2}$$

It is given that L is finite $\Rightarrow \frac{1}{a} = \frac{1}{2}$

$$\Rightarrow a = 2$$

When $a = 2$

$$\begin{aligned}
 L &= \lim_{x \rightarrow 0} \frac{2 - \sqrt{4 - x^2} - \frac{x^2}{4}}{x^4} \\
 &= \lim_{x \rightarrow 0} \frac{\left(2 - \frac{x^2}{4} \right)^2 - (4 - x^2)}{x^4 \left(2 - \frac{x^2}{4} + \sqrt{4 - x^2} \right)} = \lim_{x \rightarrow 0} \frac{1}{16} \cdot \frac{1}{4} = \frac{1}{64}
 \end{aligned}$$

31. $2 \cos \frac{B+C}{2} \cos \frac{B-C}{2} = 4 \sin^2 \frac{A}{2}$

$$2 \sin \frac{A}{2} \cdot \cos \frac{B-C}{2} = 4 \sin^2 \frac{A}{2}$$

$$\cos \left(\frac{B-C}{2} \right) = 2 \sin \frac{A}{2}$$

$$= 2 \cos \frac{B+C}{2}$$

$$\cos \frac{B}{2} \cos \frac{C}{2} + \sin \frac{B}{2} \sin \frac{C}{2}$$

$$= 2 \left\{ \cos \frac{B}{2} \cos \frac{C}{2} - \sin \frac{B}{2} \sin \frac{C}{2} \right\}$$

$$\cos \frac{B}{2} \cos \frac{C}{2} = 3 \sin \frac{B}{2} \sin \frac{C}{2}$$

$$\tan \frac{B}{2} \tan \frac{C}{2} = \frac{1}{3}$$

$$\Rightarrow \sqrt{\frac{(s-c)(s-a)}{s(s-b)}} \sqrt{\frac{(s-a)(s-b)}{s(s-c)}} = \frac{1}{3}$$

$$\Rightarrow \frac{s-a}{s} = \frac{1}{3}$$

$$3s - 3a = s$$

$$2s - 3a = 0$$

$$a + b + c - 3a = 0$$

$$b + c = 2a$$

$$b + c = 2a \text{ means}$$

$$CA + BA = 2a, \text{ a constant}$$

$$\Rightarrow \text{Locus of A is an ellipse}$$

32. Given $\frac{\sin^4 x}{2} + \frac{\cos^4 x}{3} = \frac{1}{5} \quad \text{--- (1)}$

Dividing by $\cos^4 x$

$$\frac{\tan^4 x}{2} + \frac{1}{3} = \frac{\sec^4 x}{5}$$

$$= \frac{(1 + \tan^2 x)^2}{5}$$

$$\Rightarrow \tan^4 x \left(\frac{1}{2} - \frac{1}{5} \right) - \frac{2}{5} \tan^2 x + \frac{1}{3} - \frac{1}{5} = 0$$

$$\Rightarrow \frac{3}{10} \tan^4 x - \frac{2}{5} \tan^2 x + \frac{2}{15} = 0$$

$$\Rightarrow 9 \tan^4 x - 12 \tan^2 x + 4 = 0$$

$$\Rightarrow (3 \tan^2 x - 2)^2 = 0$$

$$\Rightarrow \tan^2 x = \frac{2}{3} \quad \text{--- (2)}$$

∴ (A) is true

$$\begin{aligned} & \frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} \\ &= \cos^8 x \left\{ \frac{\tan^8 x}{8} + \frac{1}{27} \right\} \\ &= (\cos^2 x)^4 \left\{ \left(\frac{2}{3} \right)^4 + \frac{1}{27} \right\} \\ &= \left(\frac{1}{1 + \tan^2 x} \right)^4 \left\{ \frac{16}{81 \times 8} + \frac{1}{27} \right\} \\ &= \left(\frac{3}{5} \right)^4 \left\{ \frac{2}{81} + \frac{1}{27} \right\} \\ &= \frac{81}{625} \times \frac{5}{81} = \frac{1}{125} \end{aligned}$$

$$\begin{aligned} \text{Equation (2)} \Rightarrow \frac{\sin^2 x}{2} &= \frac{\cos^2 x}{2} = k \\ \Rightarrow \sin^2 x &= 2k \text{ and } \cos^2 x = 3k \end{aligned}$$

$$\therefore 2k + 3k = 1$$

$$\Rightarrow k = \frac{1}{5}$$

$$\begin{aligned} \therefore \frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} &= \frac{(2k)^4}{8} + \frac{(3k)^4}{27} \\ &= k^4 [2 + 3] = 5k^4 = \frac{1}{125} \end{aligned}$$

Section III

33. A symmetric matrix can be written as $\begin{pmatrix} a & d & e \\ d & b & f \\ e & f & c \end{pmatrix}$

But we have five 1s and four 0s.

The three symmetrical pairs can be filled as per the following.

Case 1

2 pairs of 1s and 1 pair of 0s. This is done in 3 ways. The main diagonal is filled using the remaining 1, 0, 0 in 3 ways.

∴ 9 ways.

Case 2

1 pair of 1s and 2 pairs of 0s. This is done in 3 ways. The main diagonal is filled using the remaining 1, 1, 1

∴ Total 3 ways

∴ 9 + 3 = 12 matrices

34. The matrices are

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

(1) (2) (3)

$$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

(4) (5) (6)

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

(7) (8) (9)

$$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}$$

(10) (11) (12)

Determinants of the matrices 1, 2, 3, 6, 9 and 12 are zeros and all the other 6 matrices are non-singular. Each of these six matrices provide a unique solution to the given system.

35. When we observe matrices 1 and 9, since right

hand side is $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$, they vanish for all Δ_i and thus

give infinite number of solutions. Matrices 2, 3, 6 and 12 give inconsistent systems.

36. $P(X = 3) = \frac{5}{6} \cdot \frac{5}{6} \cdot \frac{1}{6} = \frac{25}{216}$

37. $P(X \geq 3) = 1 - P(X = 1 \text{ or } X = 2)$
 $= 1 - \left[\frac{1}{6} + \frac{5}{6} \cdot \frac{1}{6} \right] = 1 - \frac{11}{36} = \frac{25}{36}$

38. $P(X \geq 6 / X > 3) = P\left(\frac{(X \geq 6) \cap (X > 3)}{P(X > 3)}\right)$

$$= \frac{P(X \geq 6)}{P(X \geq 4)}$$

$$= \frac{\left(\frac{5}{6}\right)^5 \cdot \frac{1}{6} + \left(\frac{5}{6}\right)^6 \cdot \frac{1}{6} + \dots}{\left(\frac{5}{6}\right)^3 \cdot \frac{1}{6} + \left(\frac{5}{6}\right)^4 \cdot \frac{1}{6} + \dots}$$

$$= \frac{\left(\frac{5}{6}\right)^5}{\left(\frac{5}{6}\right)^3} = \frac{25}{36}$$

Section IV

39. (A) $\frac{dy}{dx} = \frac{-y}{(x-3)^2}$

$$\frac{dy}{y} = -\frac{dx}{(x-3)^2}$$

$$\ln y = \frac{1}{x-3}$$

$$y = e^{\frac{1}{x-3}}$$

Domain of non zero solution is $D : \mathbb{R} - \{3\}$

Intervals contained in the domain D are

$$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right), \left(0, \frac{\pi}{2}\right), \left(0, \frac{\pi}{8}\right)$$

$\therefore A \rightarrow p, q, s$

(B) $I = \int_1^5 (x-1)(x-2)(x-3)(x-4)(x-5) dx$

$$= \int_{-2}^2 (t+2)(t+1)t(t-1)(t-2) dt$$

$$= 0$$

$$(\because \int_{-a}^a f(x) dx = 0, \text{ if } f(-x) = -f(x))$$

Intervals containing the value $I = 0$ are

$$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right), (-\pi, \pi)$$

$B \rightarrow (p, t)$

(C) $y = \cos^2 x + \sin x$

$$y' = -2\cos x \sin x + \cos x$$

$$= \cos x (-2\sin x + 1) = -\sin 2x + \cos x$$

For extremum, $y' = 0$

$$\Rightarrow \cos x = 0 \text{ or } \sin x = \frac{1}{2}$$

$$y'' = -2\cos 2x - \sin x$$

When $\cos x = 0$, $y'' = 2(1) - 1 > 0$

$\therefore \cos 2 = 0$ gives a local minimum

$$\text{When } \sin x = \frac{1}{2},$$


$$y'' = -2\left(1 - \frac{2}{4}\right) - \frac{1}{2} < 0$$

$$\Rightarrow \sin x = \frac{1}{2} \text{ gives a local maximum}$$

$$\Rightarrow x = n\pi + (-1)^n \frac{\pi}{6}$$

$\therefore C \rightarrow p, q, r, t$

(D)



$$y = \tan^{-1}(\sin x + \cos x)$$

$$y' = \frac{1}{(\sin x + \cos x)^2 + 1} (\cos x - \sin x)$$

$y = f(x)$ is increasing if $y' > 0$

$\Rightarrow \cos x > \sin x$ since denominator > 0

$$\Rightarrow x \in \left(-\frac{3\pi}{4}, \frac{\pi}{4}\right) \cup \left(\frac{5\pi}{4}, \frac{9\pi}{4}\right)$$

Interval in which y is increasing is $\left(0, \frac{\pi}{8}\right)$

$D \rightarrow s$

40. (p) $m = \frac{-h}{k}, a = 2, c = \frac{1}{k}$

$$\frac{1}{k^2} = 4\left(1 + \frac{h^2}{k^2}\right)$$

$$\Rightarrow h^2 + k^2 = \frac{1}{4}$$

\Rightarrow Locus of (h, k) is a circle

\Rightarrow (A)

(q) Difference = a constant 3.

\Rightarrow Locus of z is a hyperbola \Rightarrow (D)

(r) $x = \sqrt{3} \cos 2\theta, y = \sin 2\theta$

$$\frac{x^2}{3} + \frac{y^2}{1} = 1$$

\Rightarrow Ellipse \Rightarrow (C)

(s) Eccentricity = 1 \rightarrow Parabola

Eccentricity $> 1 \rightarrow$ hyperbola

\Rightarrow (B), (D)

(t) $\operatorname{Re}\{(x+1+iy)^2\} = x^2 + y^2 + 1$

$$\Rightarrow (x+1)^2 - y^2 = x^2 + y^2 + 1$$

$$\Rightarrow 2y^2 = 2x$$

$$\Rightarrow y^2 = x$$

\Rightarrow Parabola \Rightarrow (B)

PART III

41 42 43 44 45 46 47 48
B B C A C D A D

49 50 51 52
A C, D A, D B, D

53 54 55 56 57 58
D A B A B D

59	60
A – p, r, s	A – p, t
B – r, s	B – q, s, t
C – p, q, t	C – p, r, t
D – r, s	D – q


Section I

41. $\frac{Q_1}{R_1^2} = \frac{Q_1 + Q_2}{R_2^2} = \frac{Q_1 + Q_2 + Q_3}{R_3^2}$
 $\Rightarrow \frac{Q_2}{Q_1} = 3; \frac{Q_3}{Q_1} = 5;$

42. At 60° , $mg \sin \theta \frac{h}{2} > mg \cos \theta \frac{a}{2}$
 \therefore it will topple at $\theta < 60^\circ$

43. $v^2 = 2gs = 2 \times 10 \times (20 - 12.8) \Rightarrow$
 $v = 12 \text{ m s}^{-1}$
 $v' = \mu \times v = \frac{4}{3} \times 12 = 16 \text{ m s}^{-1}$

44. $y_{CM} = \frac{ma + ma + m \cdot 0 + m(-a) + 6m \cdot 0}{10m} = \frac{a}{10}$



46. $\phi = AB$, increases. By Lenz's law, induced current in direction dc and ab

47. Charged enclosed = $\frac{1}{2}$ that on disc + $\frac{1}{4}$ that on rod + point charge $-7c$
 $\therefore \phi = \frac{-2C}{\epsilon_0}$

48. $T = 8\text{s}$, phase = $\frac{2\pi}{T} \cdot t = \frac{\pi}{3}$
 $\omega = \frac{2\pi}{T} \therefore a = -\omega^2 A \cdot \sin \frac{\pi}{3}$ ($A = 1 \text{ cm}$)
 $= \frac{-\sqrt{3}}{32} \pi^2 \cdot \text{cm s}^{-2}$

Section II

49. Internal forces can convert K.E to P.E (eg. Spring masses system). Since Newton's third law. A couple exerts no force but a torque.

50.

Reading	f	Error	Calculation
(42, 56)	24	0	$0.2 \times \left(\frac{24}{56}\right)^2$
(48, 48)	24	0	$0.2 \times \left(\frac{24}{48}\right)^2$
(60, 40)	24	0	$0.2 \times \left(\frac{24}{40}\right)^2$
(66, 33)	22	-2	$0.2 \times \left(\frac{24}{33}\right)^2$
(78, 39)	26	+2	$0.2 \times \left(\frac{24}{39}\right)^2$

$$51. R_{eq} = 3.2 \text{ K}\Omega \Rightarrow I = \frac{24\text{V}}{3.2\text{K}\Omega} = 7.5 \text{ mA}$$

$$V_{RL} = 7.5 \text{ mA} \times 1.2 \text{ K}\Omega = 9\text{V}$$

$$\text{Effective emf formula} = \frac{\frac{E}{R_1}}{\frac{1}{R_1} + \frac{1}{R_2}} \quad \text{and}$$

$$\frac{\frac{E}{R_2}}{\frac{1}{R_2} + \frac{1}{R_1}} \Rightarrow \text{ratio} = 3$$

$$\therefore \text{Ratio of power} = 9$$

$$52. C_p - C_v = R \text{ for all gases}$$

$$C_v = \frac{3}{2} R \text{ for monoatomic}$$

$$\frac{5}{2} R \text{ for diatomic}$$

Section III

53. High temperature ionizes the gas

$$54. \text{ Total KE} = 3KT = P.E = \frac{e^2}{4\pi\epsilon_0} \cdot \frac{1}{r}$$

$$\therefore T \approx 1.4 \times 10^9 \text{ K}$$

55. Multiply and check nt with Lawson Number

$$56. n \frac{\lambda}{2} = a$$

$$p = \frac{h}{\lambda}$$

$$E = \frac{p^2}{2m} \Rightarrow E \propto \frac{1}{\lambda^2} \propto \frac{1}{a^2}$$

$$57. E = \frac{h^2}{8ma^2} \Big|_{\text{for } n=1} = 8 \times 10^{-3} \text{ eV}$$

$$\left(E = \frac{p^2}{2m} = \frac{\left(\frac{h}{\lambda}\right)^2}{2m} = \frac{\left(\frac{h}{2a}\right)^2}{2m} = \frac{h^2}{8ma^2} \right)$$

$$58. v \propto p, p = \frac{h}{\lambda} \Rightarrow \lambda \propto \frac{1}{n}$$

$$\Rightarrow p \propto h \Rightarrow v \propto n$$

Section IV

59. Unlike charges moving along a circle \Rightarrow no current (say reason 1)

(p) +, - charges are symmetric

$$\therefore E = 0$$

Same reason, $V = 0$

Due to reason 1, $B = 0$ and $\mu = 0$

(q) Unsymmetric distribution of charges about M. Hence $E \neq 0$ and $V = 0$

Due to reason (1), $B = 0$ and $\mu = 0$

(r) Due to symmetry $E = 0, V \neq 0$

Clearly $B \neq 0, \mu \neq 0$

(s) By symmetry, $E = 0$, distances being not commensurate, $V \neq 0$, negative currents reinforce B plus charges oppose but of different magnitude.

(t) Due to lack of symmetry $E \neq 0$. But V can be zero. Due to reason (1) $B = 0 \Rightarrow \mu = 0$

60. (p) Y has constant velocity. Therefore, reaction force is equal to weight.

PE is continuously decreasing. Mechanical energy decreasing due to frictional loss. Torque is variable

(q) Magnetic force between Z and Y is Mg

\therefore Normal reaction is 2 Mg. Since it is moving up gravitational P.E is increasing and thus mechanical energy is increasing. By symmetry, torque is zero

(r) Pulley supports the mass M. So reaction force = $(m_0 + \sqrt{2}M)g$. Since it is moving down gravitational P.E is decreasing and so the mechanical energy is decreasing. Torque is a non-zero constant

(s) Sphere moving down with uniform acceleration. Therefore force $< Mg$. Gravitational P.E of x is increasing and Mechanical energy is conserved. Torque is a non-zero constant

(t) Terminal velocity \Rightarrow net force zero. Gravitational P.E of x is increasing, but mechanical energy is decreasing because of frictional forces. Torque is a non-zero constant.