

**DipIETE – ET / CS (OLD SCHEME)**

Code: DE23/DC23  
Time: 3 Hours

**DECEMBER 2010**

Subject: MATHEMATICS - II  
Max. Marks: 100

**NOTE:** There are 9 Questions in all.

- Question 1 is compulsory and carries 20 marks. Answer to Q.1 must be written in the space provided for it in the answer book supplied and nowhere else.
- The answer sheet for the Q.1 will be collected by the invigilator after half an hour of the commencement of the examination.
- Out of the remaining EIGHT Questions answer any FIVE Questions. Each question carries 16 marks.
- Any required data not explicitly given, may be suitably assumed and stated.

**Q.1 Choose the correct or the best alternative in the following: (2×10)**

- a. If  $x + iy = \sqrt{2} + 3i$ , then  $x^2 + y$  is  
 (A) 7 (B) 5  
 (C) 13 (D)  $\sqrt{2} + 3$
- b. If  $\sin x = \frac{e^{ix} - e^{-ix}}{2i}$ , then  $\cos x$  is equal to,  
 (A)  $\frac{e^{ix} + e^{-ix}}{2}$  (B)  $\frac{e^{-ix} + e^x}{2}$   
 (C)  $\frac{e^x - e^{-x}}{2}$  (D)  $\frac{e^x - e^{-ix}}{2}$
- c. If three vectors  $\vec{a}, \vec{b}, \vec{c}$  are coplanar then,  
 (A)  $(\vec{a} \times \vec{b}) \times \vec{c} = 0$  (B)  $(\vec{a} \times \vec{b}) \cdot \vec{c} = 0$   
 (C)  $(\vec{a} \times \vec{b}) \cdot \vec{c} = 1$  (D)  $(\vec{a} \times \vec{b}) \cdot \vec{c} = -1$
- d. If  $|\vec{A} + \vec{B}| = 30$ ,  $|\vec{A} - \vec{B}| = 20$  and  $|\vec{B}| = 23$ , then  $|\vec{A}|$  is equal to  
 (A) 12 (B) 13  
 (C) 11 (D) 14
- e. If  $A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$ , then  $A^{-1}$  is  
 (A)  $\frac{1}{2} \begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix}$  (B)  $\frac{1}{3} \begin{bmatrix} 2 & 1 \\ -1 & 2 \end{bmatrix}$   
 (C)  $\frac{1}{3} \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$  (D)  $\frac{1}{2} \begin{bmatrix} -2 & 1 \\ 1 & 2 \end{bmatrix}$

f. For what value of  $x$  is the matrix

$$\begin{bmatrix} 3-x & 2 & 2 \\ 1 & 4-x & 1 \\ -2 & -4 & -1-x \end{bmatrix} \text{ singular?}$$

- (A) 1,2,3 (B) 1,-2,-3  
(C) 1,2,5 (D) 1,-2,-5

g. The characteristic roots of the matrix  $\begin{bmatrix} -6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$  is

- (A) 2,3,6 (B) 1,2,3  
(C) 2,2,8 (D) 1,2,6

h. The period of  $|\cos x|$  is

- (A)  $\pi/2$  (B)  $\pi$   
(C)  $3\pi/2$  (D)  $2\pi$

i. The inverse Laplace transform of  $\frac{1}{S(S+2)}$  is

- (A)  $-\frac{1}{2}[e^{2t}-1]$  (B)  $-\frac{1}{2}[e^{-2t}-1]$   
(C)  $-\frac{1}{2}[e^{-2t}-2]$  (D)  $-\frac{1}{2}[e^{-2t}+2]$

j. The solution of differential equation  $\frac{d^2y}{dx^2} + 9y = e^x - \cos 2x$  is

- (A)  $y = c_1 \cos 3x + c_2 \sin 3x + \frac{1}{10}e^x - \frac{1}{5}\cos 2x$   
(B)  $y = c_1 \cos 3x - c_2 \sin 3x - \frac{1}{10}e^{-x} + \frac{1}{5}\cos 2x$   
(C)  $y = c_1 \cos 3x + c_2 \sin 3x + 10e^x - 5\cos 2x$   
(D)  $y = c_1 \cos 3x - c_2 \sin 3x + 5e^x - 5\cos 2x$

---

**Answer any FIVE Questions out of EIGHT Questions.  
Each question carries 16 marks.**

---

**Q.2** a. Show that the roots of the equation  $x^{10} + 11x^5 - 1 = 0$  are  $\left(\frac{\pm\sqrt{5}-1}{2}\right)\left(\cos\frac{2n\pi}{5} + i\sin\frac{2n\pi}{5}\right)$ , where  $n=0,1,2,3,4$ . (8)

b. If  $\sin(\alpha + i\beta) = x + iy$ , then show that,  $\frac{x^2}{\cosh^2 \beta} + \frac{y^2}{\sinh^2 \beta} = 1$ . (8)

**Q.3** a. The centre of a regular hexagon is at the origin and one vertex is given by  $\sqrt{3} + i$  on the Argand plane. Find the complex number represented by the other vertices. (8)

b. Show that the line joining one vertex of parallelogram to the mid-point of an opposite side trisects the diagonal and is trisected there at. (8)

**Q.4** a. Forces of magnitude 5,3,1 Kg acting on the directions  $6i + 2j + 3k$ ,  $3i - 2j + 6k$ ,  $2i - 3j - 6k$  respectively act on a particle which is displaced from the point  $(2, -1, -3)$  to  $(5, -1, 1)$ . Find the work done by the forces, the unit of length being metre. (8)

b. Find an unit vector parallel to the sum of the vectors  $\vec{a} = 2i + 4j - 5k$  and  $\vec{b} = i + 2j + 3k$ . (8)

**Q.5** a. Evaluate  $\begin{vmatrix} x-2 & 2x-3 & 3x-4 \\ x-4 & 2x-9 & 3x-16 \\ x-8 & 2x-7 & 3x-64 \end{vmatrix} = 0$ . (8)

b. Solve the equation using Cramer's Rule.  
 $3x - 4y - z = 2$   
 $6x + 6y + 3z = 7$  (8)  
 $9x - 8y - 5z = 0$

**Q.6** a. Find the values of  $\lambda$  for which the following system of equation is consistent and has nontrivial solution. Solve the equation for all such values of  $\lambda$   
 $(\lambda - 1)x + (3\lambda + 1)y + 2\lambda z = 0$   
 $(\lambda - 1)x + (4\lambda - 2)y + (\lambda + 3)z = 0$  (8)  
 $2x + (3\lambda + 1)y + 3(\lambda - 1)z = 0$

b. Find the characteristic equation of the matrix  $A = \begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$  and hence find the inverse of the matrix A. (8)

**Q.7** a. Find the Laplace transform of  $e^{-3t}(\cos 4t + 3\sin 4t)$ . (8)

b. Find the Inverse Laplace Transform of  $\frac{3s - 2}{s^2 - 4s + 20}$ . (8)

- Q.8** a. Use Laplace transform technique to solve  $\frac{d^2x}{dt^2} + 9x = \cos 2t$ , given that  $x(0) = 1$  and  $x\left(\frac{\pi}{2}\right) = -1$ . (8)
- b. Solve the differential equation  $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = x \cdot e^x \cdot \sin x$ . (8)
- Q.9** a. Show that any real valued function can be uniquely expressed as the sum of an even function and an odd function. (8)
- b. Find the Fourier series for the function  $f(x) = x$  in the interval  $[-\pi, \pi]$ . (8)