**Test Paper Code: BT** 

Student Bounty Com Time: 3 Hours Max. Marks: 300

#### INSTRUCTIONS

#### A. General:

- 1. This Booklet is your Question Paper. It contains **X** pages and has 100 questions.
- 2. The Question Booklet **Code** is printed on the right-hand top corner of this page.
- 3. The Question Booklet contains blank spaces for your rough work. No additional sheets will be provided for rough work.
- 4. Clip board, log tables, slide rule, calculator, cellular phone and electronic gadgets in any form are NOT allowed.
- 5. Write your Name and Registration Number in the space provided at the bottom.
- 6. All answers are to be marked only on the machine gradable Objective Response Sheet (ORS) provided along with this booklet, as per the instructions therein.
- 7. The Question Booklet along with the Objective Response Sheet (ORS) must be handed over to the Invigilator before leaving the examination hall.
- 8. Refer to **Special Instruction/Useful Data** on reverse of this sheet.

#### B. Filling-in the ORS:

- 9. Write your Registration Number in the boxes provided on the upper left-hand-side of the ORS and darken the appropriate bubble under each digit of your Registration Number using a HB pencil.
- 10. Ensure that the code on the Question Booklet and the code on the ORS are the same. If the codes do not match, report to the Invigilator immediately.
- 11. On the lower-left-hand-side of the **ORS**, write your Name, Registration Number, Name of the Test Centre and put your signature in the appropriate box with ballpoint pen. Do not write these anywhere else.

# C. Marking of Answers on the ORS:

- 12. Each guestion has 4 choices for its answer: (A), (B), (C) and (D). Only ONE of them is the correct answer.
- 13. On the right-hand-side of ORS, for each question number, darken with a HB Pencil ONLY one bubble corresponding to what you consider to be the most appropriate answer, from among the four choices.
- 14. There will be **negative marking** for wrong answers.

#### MARKING SCHEME:

- (a) For each correct answer, you will be awarded 3 (Three) marks.
- (b) For each wrong answer, you will be awarded -1 (Negative one) mark.
- (c) Multiple answers to a question will be treated as a wrong answer.
- (d) For each un-attempted question, you will be awarded **0** (Zero) mark.

Name				
Registration Number				

**Special Instructions/ Useful Data** 

Student Bounty Com

				below. Which one of the	
Q.1		values of four nucleic a les has the highest T <sub>m</sub> ?	cid samples are provided	below. Which one of the	
	(B) A = 26; T = (C) A = 17; T =	21; G = 20; C = 28 14; G = 34; C = 26 19; G = 33; C = 31 30; G = 25; C = 25			COM
Q.2	Which one of th made up of	e following is <b>TRUE</b> re	garding organization of h	uman chromosomes? It is	
	(B) extra-chrom	t are acidic proteins. osomal circular DNA. nat consists of DNA and osomal DNA.	basic proteins.		
Q.3	The melting poi	nt of unsaturated fatty a	eid		
	(B) increases wi (C) is higher that	d to the number of doub th increase in the number in that of its correspondition ith increase in the numb	er of double bonds.  ng saturated fatty acid.		
Q.4	Match the horm Group I P. Progesterone Q. Glucagon R. Insulin S. Androgen	ones in Group I with the	<ul><li>2. Implantation of</li><li>3. Stimulates spo</li></ul>	Group II Group II oneogenesis in liver of fertilized ovum ermatogenesis process acose uptake and storage	
	(A) <b>P-2, Q-1, R</b> (C) <b>P-1, Q-4, R</b>		(B) <b>P-3, Q-2, R</b> -(D) <b>P-1, Q-2, R</b> -		
Q.5	The most abund	ant immunoglobulin in l	human blood is		
	(A) IgM.	(B) IgA.	(C) IgD.	(D) IgG.	
Q.6	The process of p	ourification and recovery	of a product in biotechn	ology is known as	
	<ul><li>(A) upstream pr</li><li>(C) incubation.</li></ul>	ocessing.	(B) downstream (D) formulation.		
Q.7		f an enzyme catalyzed ro [3] to Michaelis-Menton o	eaction is 60% of $v_{max}$ , the constant $K_{M}$ is	en the ratio of substrate	
	(A) 1	(B) 1.5	(C) 2	(D) 4	
Q.8	In a DNA replic with <sup>14</sup> N DNA.	ation experiment, 1 μg of 14 The amount (in μg) of 14	of <sup>15</sup> N DNA is allowed to <sup>15</sup> N DNA formed during th	replicate till two generations are second replication process	

(D) form water from exwaen

				S.	
				ide	
				THE	
Q.25	Regenerative medic	ine aims at		Elite	
	(A) discovering sma (C) growing tissues		<ul><li>(B) generating th</li><li>(D) identifying g</li></ul>	erapeutic proteins. enetic mutations.	-
Q.26	Which of the follow	ring is <b>NOT</b> required in	a Polymerase Chain R	eaction?	
	(A) DNA template (B) Mg <sup>++</sup> ion (C) Primers (D) Restriction enzy	/mes			
Q.27	Which one of the fo genome?	llowing processes allow	vs introduction of gene	of interest to a target site in	
	(A) Somatic embryo (B) Organogenesis (C) Gene cloning (D) Southern hybrid				
Q.28	Based on the dissoc is	iation constant $K_d$ , the p	protein - ligand pair tha	t has the strongest interaction	
	(B) avidin and biotin (C) HIV surface pro	lin receptor ( $K_d = 1 \times 10^{-15}$ ). the n ( $K_d = 1 \times 10^{-15}$ ). the n and anti-HIV IgG (calcium ( $K_d = 3 \times 10^{-6}$ )	$(K_{\rm d} = 4 \times 10^{-10}).$		
Q.29	In genetic code, the	codon degeneracy occu	ırs at	position(s).	
	(A) first	(B) second	(C) third	(D) first and third	
Q.30	plant germinated to	•	plants and 240 yellow	1000 seeds taken from a pea pod plants. The parental	
	(A) heterozygous ar (C) heterozygous an	2	<ul><li>(B) homozygous</li><li>(D) homozygous</li></ul>		
Q.31	Which of the follow	ring is <b>FALSE</b> for DNA	Λ?		
	(C) Orientation of o	DNA associate in parall ne strand is 3' to 5' and	other strand is 5' to 3'	s is due to hydrogen bonds.	
Q.32	In 2009, the swine f	lu outbreak was	in nature.		
	(A) sporadic	(B) pandemic	(C) chronic	(D) endemic	

# Student Bounty.com

- Q.33 In angiosperms, the microsporangia develops to form
  - (A) stigma.
- (B) ovule.
- (C) endosperm.
- (D) pollen sacs.
- Q.34 Given the  $pK_a$  values of different acidic sites in cysteine, the principal ionic form in which it exists at pH 7.0, is

- (A)

(B)

(C)

(D)

- Q.35 evolution, anatomical structures develop in different directions to adapt different functions.
  - (A) convergent, homologous.
  - (B) divergent, homologous.
  - (C) convergent, analogous.
  - (D) divergent, analogous.
- Q.36 A model of gene control for the *lac* operon is shown below.

_						
	I	P	O	Z	Y	A

Match the component of *lac* operon in Group I with the function listed in Group II.

## Group I

- K. O
- L. P
- M. Y
- N. Α
- (A) K-2, L-3, M-4, N-1
- (C) K-3, L-2, M-1, N-4

# **Group II**

- 1. Encodes protein  $\beta$ -galactoside permease
- **2.** Provides binding site for RNA polymerase
- 3. Initiates *lac* mRNA synthesis
- **4.** Encodes protein thiogalactoside transacetylase
- (B) K-3, L-2, M-4, N-1
- (D) K-2, L-3, M-1, N-4

$$\begin{array}{c} \text{(A)} \\ \text{HO} & \text{H}_3 \\ \text{HO} & \text{H} \\ \text{H} & \text{Br} \\ \text{CH}_3 \end{array}$$

$$\dot{C}H_3$$
(C)
 $CH_3$ 
 $HO \longrightarrow H$ 
 $Br \longrightarrow H$ 
 $CH_3$ 

Q.47 Match the compounds in Group I with their appropriate spectroscopic data in Group II.

(B)

# Group I

- P. CH<sub>3</sub>COCH<sub>3</sub>
- Q. CH<sub>3</sub>CH<sub>2</sub>COOH
- R. CH<sub>3</sub>COOCH<sub>3</sub>
- (C) P-1; Q-3; R-2

# **Group II**

- 1. two singlets of equal intensity in the <sup>1</sup>H-NMR spectrum
- 2. a band at 1720 cm<sup>-1</sup> in the IR spectrum
- 3. an intense peak at m/z 45 in the mass spectrum
- (A) P-1; Q-2; R-3
- (B) P-2; Q-3, R-1 (D) P-2; Q-1; R-3
- Among the following compounds, the one that is soluble in aqueous NaOH but not in aqueous Q.48 NaHCO<sub>3</sub>, is

Q.49 The major product of the following reaction sequence, is

Q.50 The major product formed in the E-2 elimination reaction of the following compound, is

$$\begin{array}{c} \text{(A)} \\ \text{H} \\ \text{H}_{3}\text{C} \\ \end{array}$$

(B) 
$$H_3C^{(1)}$$
  $D$ 

# Student Bounty Com

Q.51 The most reactive diene towards Diels-Alder reaction, among the following, is

(A)



(B)



(C)




(D)



Q.52 The correct structure of pyrophosphorus acid is

(A)



(B)

(D)

- Q.53 Among the following complexes, the one which exhibits optical isomerism, is (note: en = ethylenediamine)
  - (A) cis-[Co(en)<sub>2</sub>Cl<sub>2</sub>]<sup>+</sup>

(B) cis-[Pt(NH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>]

(C) trans-[Co(en)<sub>2</sub>Cl<sub>2</sub>]<sup>+</sup>

- (D)  $trans-[Pt(NH_3)_2Cl_2]$
- Q.54 The gas that is produced on treating NaCl with conc. H<sub>2</sub>SO<sub>4</sub> is
  - $(A) O_2$
- (B) Cl<sub>2</sub>
- (C) SO<sub>2</sub>
- (D) HCl
- The correct order of the atoms in terms of their first ionization energy is Q.55
  - (A) Li < B < Be < C

(B) Li<Be<B<C

(C) Li>B>Be>C

- (D) Li>Be>B>C
- Q.56 The compound with square planar geometry is
  - $(A) [Ni(CO)_4]$

(B) [Ni(CN)<sub>4</sub>]<sup>2-</sup> (D) [NiCl<sub>4</sub>]<sup>2-</sup>

(C)  $[Ni(PPh_3)_2Cl_2]$ 

Q.57 Match the molecules in Group I with their shape in Group II.

## Group I

 $\mathbf{P}$ .  $ICl_2^-$ 

 $\mathbf{Q}$ .  $\mathbf{H}_2\mathbf{O}$ 

R. PCl<sub>5</sub>

# **Group II**

"AUGENTBOUNTS, COM 1. trigonal bipyramid

2. linear

**3.** V-shaped

4. square pyramid

$$(A) P-3, Q-2, R-4$$

(C) P-2, Q-3, R-1

(B) **P-4**, **Q-3**, **R-1** 

(D) P-4, O-3, R-2

The spin-only magnetic moment of  $[Fe(CN)_6]^{4-}$  is Q.58

(A) 4.9 D

(B) 0 D

(C) 5.92 D

(D) 2.82 D

Q.59 One mole of a gas absorbs 40 J of heat. If the work done on the surrounding by the gas is 20 J, then  $\Delta U$  (in J) for the gas is

(A) 60

(B) 20

(C) -20

(D) -60

For the reaction,  $N_2O_4(g) \Leftrightarrow 2NO_2(g)$ , taking place in a closed container at a constant Q.60 temperature, the rate constant k in terms of  $P_0$  (pressure at time t = 0) and  $P_t$  (pressure at time t) is given by

(A) 
$$\frac{1}{t} \ln \frac{P_0}{2P_0 - P_t}$$
 (B)  $\frac{1}{t} \ln \frac{P_0}{P_t}$  (C)  $\frac{1}{t} \ln \frac{P_0}{P_0 - P_t}$  (D)  $\frac{1}{t} \ln \frac{P_0}{P_0 - 2P_t}$ 

(B) 
$$\frac{1}{t} \ln \frac{P_0}{P_t}$$

$$(C) \frac{1}{t} \ln \frac{P_0}{P_0 - P_t}$$

(D) 
$$\frac{1}{t} \ln \frac{P_0}{P_0 - 2P_t}$$

Q.61  $pK_a$  of acetic acid is 4.80. A 10 mL of 1M solution of acetic acid is mixed with 5 mL of 1 M solution of NaOH. The pH of the resulting solution is

(A) 3.2

(B) 7.0

(C) 4.8

(D) 2.4

The series that corresponds to transition from higher levels to n = 4 in the hydrogen spectrum is Q.62

(A) Paschen

(B) Balmer

(C) Pfund

(D) Brackett

Q.63 For the reaction,  $A \to product$ , match the order of the reaction in Group I with their corresponding linear plots in Group II.

# Group I

P. Zero

**Q.** First

**R.** Second

#### **Group II**

1. ln[A] versus time

2. 1/[A] versus time

**3.** [A] versus time

(A) P-1, Q-2, R-3

(B) **P-2, Q-1, R-3** 

(C) P-3, Q-1, R-2

(D) P-1, Q-3, R-2

$$(A) + 1.06 V$$

$$(B) -1.06 V$$

$$(C) -0.48 \text{ V}$$

(D) 
$$-1.83 \text{ V}$$

Student Bounts, com The depth of a swimming pool filled with clean water (refractive index = 4/3) appears to be Q.65 3 m to a person standing near it. Its actual depth is

Q.66 A semiconductor device that has two p-n junctions is

- (A) rectifier-diode
- (B) photo-diode
- (C) transistor
- (D) solar-cell

Q.67 The resolution of a microscope is directly proportional to the wavelength of the radiation used for its operation. Among the following, maximum possible resolution can be achieved from

- (A) optical microscope with blue light source.
- (B) optical microscope with yellow light source.
- (C) electron microscope operating at 100 kV.
- (D) electron microscope operating at 200 kV.

Longitudinal waves can travel through Q.68

(A) gas only.

(B) gas and liquid only.

(C) gas and solid only.

(D) gas, liquid and solid.

Q.69 The waves,  $y_1 = A \sin(\omega t + kx)$  and  $y_2 = A \cos(\omega t + kx)$ ,

- (A) are in same phase.
- (B) have a phase difference of  $\pi/4$ .
- (C) have a phase difference of  $\pi/2$ .
- (D) have a phase difference of  $\pi$ .

A vertical spring is fixed at its upper end. Same sized blocks of wood (W), glass (G) and Q.70 copper (Cu) are attached to its lower end one at a time and the system is set into vertical oscillations. The three measured time periods are in the order

(A) 
$$T_{Cu} > T_G > T_W$$

(B) 
$$T_W > T_G > T_{Cu}$$
  
(D)  $T_{Cu} > T_W > T_G$ 

(C) 
$$T_G > T_{Cu} > T_W$$

(D) 
$$T_{Cu} > T_W > T_G$$


A neutron collides head-on with a He-atom at rest. Collision is elastic and He-atom recoils with Q.71 a speed of  $2 \times 10^5$  m/s. Then, the initial speed of the neutron is

(A) 
$$0.5 \times 10^5$$
 m/s. (B)  $2 \times 10^5$  m/s. (C)  $5 \times 10^5$  m/s. (D)  $8 \times 10^5$  m/s.

(B) 
$$2 \times 10^5$$
 m/s

(C) 
$$5 \times 10^5$$
 m/s

(D) 
$$8 \times 10^5$$
 m/s.



- (A) all the three layers is same.
- (C) middle layer is maximum.

- (B) top layer is maximum.
- (D) bottom layer is maximum.
- Q.73 Match the actions in Group II that will produce radiations listed in Group I.

# Group I

## Group II

- **P.**  $\gamma$  -rays
- **Q.** UV radiation
- **R.** IR radiation
- (A) **P-3, Q-1, R-2**
- (C) P-1, Q-3, R-2


3. Fusion of two light nuclei.

**2.** A body at 600 K emitting radiation.

(B) **P-3**, **Q-2**, **R-1** (D) **P-1**, **Q-2**, **R-3** 


**1.** H-atom in 1<sup>st</sup> excited state returns to ground state.

Q.74 A rigid conducting wire PQ is moving on conducting rails (as shown in figure) with constant speed v = 6 m/s in a region of uniform field B = 0.2 Wb/m<sup>2</sup>. The magnitude of induced *emf* and direction of induced current are



- (A) 1.8 V, clockwise.
- (C) 3.6 V, clockwise.

- (B) 1.8 V, anti-clockwise.
- (D) 3.6 V, anti-clockwise.
- Q.75 A ball is projected at 30° from ground with an initial velocity of 10 m/s. Taking  $g = 10 \text{ m/s}^2$ , the horizontal range of the ball is
  - (A) 2.5 m
- (B) 5 m
- (C) 8.66 m
- (D) 10 m



 $(A) \vec{B}_P = -\vec{B}_R$ 

(B)  $\vec{B}_P = \vec{B}_R$ 

(C)  $\left| \vec{B}_{Q} \right| \neq 0$ 

Then

(D)  $\left| \vec{B}_P \right| > \left| \vec{B}_Q \right|$ 


An object weighs 50 N on Earth ( $g = 10 \text{ m/s}^2$ ). Its mass on a planet having  $g = 2 \text{ m/s}^2$  will be Q.77

- (A) 1 kg
- (B) 2.5 kg
- (C) 5 kg
- (D) 10 kg

Q.78 An 80 W fan, a 60 W bulb and a 500 W washing machine are operated for 15, 20 and 1 hr, respectively. The total electrical power units consumed are


- (A) 1.2
- (B) 1.7
- (C) 2.4
- (D) 2.9

Q.79 An electron having a velocity  $\vec{v}$  enters a region of uniform magnetic field  $\vec{B}$  as shown in figure. The effect of  $\vec{B}$  on the motion of electron is that it will



- (A) continue to move without any deflection.
- (B) be reflected back.
- (C) be deflected up.
- (D) be deflected down.

In the given circuit, an ideal battery of 15 V and resistances of 4 ohm each are connected as Q.80 shown below. The current (in amperes) through the ammeter A is




(C) 0

(D) -2

 $\alpha + \beta$  is equal to

(B) 1

(A) 2



$$1+(1+x)+\frac{(1+x)^2}{2!}+\frac{(1+x)^3}{3!}+...$$

is  $e^{\frac{1}{2}}$ , then x is

(A) 
$$-\frac{1}{2}$$

Strudent Bounts, com
(D) 
$$\frac{1}{2}$$

Q.90 The minimum value of the function 
$$f(x) = x^4 - 2x^2 + 2$$
 in  $[-1,2]$  is

(D) 
$$-2$$

Q.92 The equation of the line that makes an intercept of 2 with x-axis and is perpendicular to the line 
$$x + y - 1 = 0$$
 is

(A) 
$$x + y - 2 = 0$$

(B) 
$$x + y + 2 = 0$$

(C) 
$$x - y - 2 = 0$$

(D) 
$$x - y + 2 = 0$$

Q.94 If the point 
$$(1, 0, 1)$$
 is one extremity of the diameter of the sphere

$$x^{2} + y^{2} + z^{2} + 2x - 4y + 2z - 6 = 0$$
,

then its other extremity is

(B) 
$$(-3, 0, -3)$$

$$(C)$$
  $(3, -4, 3)$ 

(B) 
$$(-3, 0, -3)$$
 (C)  $(3, -4, 3)$  (D)  $(-3, 4, -3)$ 

Q.95 Let f be the function defined for real x as 
$$f(x) = \begin{cases} \frac{x}{|x|}, & x \neq 0 \\ 1, & x \neq 0 \end{cases}$$
 Then, f is

(A) continuous for all real 
$$x$$
.

(B) right continuous at 
$$x = 0$$
.

(C) a non-negative function for all real 
$$x$$
.

(D) left continuous at 
$$x = 0$$
.

- Q.96 An urn consists of 10 items out of which 4 are defective. Three items are chosen rank from the urn. The probability that exactly 2 from the chosen items are defective, is
  - (A)  $\frac{1}{20}$
- (B)  $\frac{2}{3}$
- (C)  $\frac{7}{10}$
- (D)  $\frac{3}{10}$

- Q.97 The eccentricity of the ellipse  $\frac{x^2}{16} + \frac{y^2}{9} = 1$  is
  - (A)  $\frac{3}{4}$
- (B)  $\frac{1}{2}$
- (C)  $\frac{\sqrt{7}}{3}$
- (D)  $\frac{\sqrt{7}}{4}$

Q.98 Suppose the statement

# "If the flower smells sweet then I will buy it",

is given to be *FALSE*. Then which one of the following is correct.

- (A) The flower does not smell sweet and I bought it.
- (B) The flower does not smell sweet and I did not buy it.
- (C) The flower smells sweet and I bought it.
- (D) The flower smells sweet and I did not buy it.
- Q.99 The values obtained in 20 throws of a die are given in the following frequency table

Value	1	2	3	4	5	6
Frequency	3	3	4	4	2	4

The sample median is

(A)3

- (B) 3.5
- (C)4

- (D) 4.5
- Q.100 The equation of the normal to the curve  $x^2y^3 = 4$  at the point (2, 1) is
  - (A) y = 3x 5

(B) 5y = 3x - 1

(C) 3y = 5 - x

(D) 5v = -x + 7