

# Physics Paper II

**Time Allowed : 75 Minutes]**

**[Maximum Marks : 100**

**Note :** This Paper contains **Fifty (50)** multiple choice questions. Each question carries **Two (2)** marks. Attempt *All* questions.

StudentBounty.com

- |   |  |
|---|--|
| <p>1. Spherical Bessel function of order <math>n</math> is a product of <math>\sqrt{\frac{\pi}{2x}}</math> with :</p> <p>(A) Bessel function of order <math>n + \frac{1}{2}</math></p> <p>(B) Bessel function of order <math>n - \frac{1}{2}</math></p> <p>(C) Bessel function transformed in polar coordinates</p> <p>(D) Spherical harmonic of order <math>n</math></p> <p>2. The angle between vector <math>\hat{i} + \hat{j}</math> and <math>\hat{j} + \hat{k}</math> is (in radian) :</p> <p>(A) <math>\pi</math></p> <p>(B) <math>\pi/2</math></p> <p>(C) <math>\pi/6</math></p> <p>(D) <math>\pi/4</math></p> | <p>3. The cube roots of unity <math>1, \omega, \omega^2</math> form :</p> <p>(A) a cyclic group of order 3</p> <p>(B) a permutation group</p> <p>(C) SU3 group</p> <p>(D) SU2 <math>\times</math> U group</p> <p>4. The value of</p> $1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \dots}}}$ <p>is :</p> <p>(A) <math>\sqrt{2}</math></p> <p>(B) 1.6</p> <p>(C) <math>\sqrt{3}</math></p> <p>(D) 0.8</p> |
|---|--|

5. The real matrix  $A = \begin{bmatrix} a & f & g \\ -f & a & -h \\ -g & h & a \end{bmatrix}$  is

skew symmetric when :

(A)  $a = 0$

(B)  $f = 0$

(C)  $g = h$

(D)  $f = g$

6. The eigenvalues of the matrix

$$\begin{pmatrix} 1 & \omega \\ \omega & 1 \end{pmatrix} \text{ are :}$$

(A) 1

(B)  $\pm \omega$

(C)  $\pm \omega^2$

(D)  $\pm i$

7. A complex function  $f(z)$  is given by :

$$f(z) = \sqrt{z} + \frac{1}{z-a} + \exp(z)$$

The singularities of  $f(z)$  are :

(A) simple pole at  $z = a$

(B) branch point at  $z = 0$

(C) essential singularity at  $z \rightarrow \infty$

(D) all of the three above

8. The determinant of a 3 × 3 symmetric matrix is 36. If two of its eigenvalues are 2 and 3, then the sum of the eigenvalues is :

(A) 30

(B) 10

(C) 11

(D) 31


9. A harmonic oscillator in one dimension is perturbed by the potential  $\alpha x^3$ . The ground state energy of the oscillator to a first order in perturbation is :

(A)  $\frac{\hbar\omega}{2} + \alpha$

(B)  $\frac{3}{2} \hbar\omega + \alpha$

(C)  $\frac{\hbar\omega}{2} + \alpha^3$

(D)  $\frac{\hbar\omega}{9}$



10. A particle moves in one-dimensional potential  $V(x)$ . At  $x = a$ , if  $V$  has a finite discontinuity (jump), then which of the following is true for its wave function  $\phi$  and its first derivative  $\phi'$  at  $x = a$  ?

- (A)  $\phi$  is continuous and  $\phi'$  must be discontinuous
- (B)  $\phi$  is discontinuous and  $\phi'$  must be continuous
- (C) both  $\phi$  and  $\phi'$  are discontinuous
- (D) both  $\phi$  and  $\phi'$  are continuous

11. In quantum mechanics, three dimensional wave function  $\psi(\vec{r})$  of a particle :

- (A) has dimension of (energy  $\times$  time)
- (B) has dimension of (length)<sup>-3/2</sup>
- (C) has dimension of energy
- (D) is dimensionless

12. A system is known to be in state described by the wave

$$\psi(\theta, \phi) = \frac{1}{\sqrt{30}} \{ 5Y_4^0 + Y_6^0 + 2Y_6^{-6} \}$$

where  $Y_l^m(\theta, \phi)$  are spherical harmonics. The probability of finding the system in a state with  $m = 0$  is :

- (A) zero
- (B)  $6/\sqrt{30}$
- (C)  $6/30$
- (D)  $13/15$

13. What is the degeneracy of the third excited state for a particle in 3-dimensional isotropic Harmonic oscillator potential ?

(Note : ground state is not an excited state)

- (A) 10
- (B) 6
- (C) 4
- (D) 3

14. The parity of wave function  $\psi$  is associated with which of the following transformation ?

- (A) Space translation
- (B) Space rotation
- (C) Space inversion
- (D) Space exchange of two particles

15. Which of the following processes involves tunnelling through a potential barrier ?

- (A) Pair production
- (B)  $\alpha$ -decay
- (C)  $\beta$ -decay

16. The variational method in perturbation theory, when applied to obtain the value of the ground state energy :

- (A) gives energy value higher than or equal to the exact ground state energy
- (B) always gives exact ground state energy
- (C) gives energy value lower than the exact ground state energy
- (D) gives energy value which is sometimes higher than or sometimes lower than the exact ground state energy

17. In a scintillation detector, the height of the output pulse is proportional to :

- (A) Energy of the incident photon
- (B) Intensity of the incident photon
- (C) Energy and intensity of the photon
- (D) Does not depend either on

18. If 'N' number of gadgets are connected to a power supply with a capacity of 'X' amperes without overloading then :

- (A) Total current drawn by all the gadgets should be equal to  $\frac{X}{2}$  ampere
- (B) Total current drawn by all the gadgets should be equal to N.X ampere
- (C) Total current drawn by all the gadgets should be equal to X
- (D) Total current drawn by all the gadgets should be equal to  $\frac{X}{4}$

19. If a square wave function generator is coupled to an oscilloscope in a.c. mode, the waveform would be observed on the oscilloscope ?

- (A) A perfect square wave
- (B) Distorted square wave
- (C) A sawtooth wave
- (D) A perfect square wave with change in repetition rate
20. If an oscilloscope is operated in a d.c. mode, one can faithfully measure :
- (A) only a.c. voltage
- (B) only d.c. voltage
- (C) both a.c. and d.c. voltage
- (D) only low frequency a.c. voltage


21. In recording a powder X-ray diffraction pattern :
- (A) the specimen and detector are both rotated
  - (B) the specimen alone is rotated
  - (C) the detector alone is rotated
  - (D) the specimen and the source are both rotated
22. In a Michelson Interferometer, the mirror  $M_2$  is moved such that 800 fringes are counted. The wavelength of the source used was  $6000 \text{ \AA}$ . Through what distance the mirror  $M_2$  must have been moved ?
- (A) 0.24 mm
  - (B) 0.48 mm
  - (C) 0.36 mm
23. For the measurement of pressure in a chamber evacuated by a diffusion pump and a rotary mechanical pump, one would require the following combination of gauges :
- (A) Thermocouple/Pirani
  - (B) Penning-Pirani
  - (C) Thermocouple/Mercury Manometer
  - (D) Pirani/Mercury Manometer
24. The vapour diffusion pump works in the following region of air flow :
- (A) Molecular flow
  - (B) Turbulent flow
  - (C) Lamellar flow

25. A cork is submerged in a pail of water by a spring attached to the bottom of the pail. The pail is held by a child in an elevator. During the initial acceleration as the elevator travels to the next lower floor, will the displacement of the spring :
- (A) increase  
(B) decrease  
(C) remain the same  
(D) indeterminate
26. A satellite is launched into a circular orbit of radius  $R$ . A second satellite is launched into an orbit of radius  $1.01 R$ . Then, the period of the second satellite is :
- (A) larger by 1.5%  
(B) smaller by 1%  
(C) larger by 2%  
(D) 1.5 hrs
27. A sphere of radius  $R$  is released in a liquid of viscosity  $\eta$ , and by Stokes' law its drag is  $6\pi\eta Rv$ . Simultaneously, a second sphere of identical mass but with radius  $2R$  is released. Then the ratio of their terminal velocities is :
- (A)  $\frac{V_R}{V_{2R}} = 1$   
(B)  $\frac{V_R}{V_{2R}} = \frac{3}{2}$   
(C)  $\frac{V_R}{V_{2R}} = 2$   
(D)  $\frac{V_R}{V_{2R}} = \frac{2}{3}$
28. What would be the approximate length of a day if the earth spun so fast that bodies floated on the equator? Take the radius of the earth =  $6 \times 10^6$  m and  $g = 9.8$  m/sec<sup>2</sup>.
- (A) 12 hrs  
(B) 6 hrs  
(C) 3 hrs  
(D) 1.5 hrs

29. An example of a scleronic, holonomic, conservative and unilateral constraint is :

- (A) simple pendulum with rigid support
- (B) simple pendulum with variable length
- (C) a spherical container of fixed radius filled with gas
- (D) an expanding or contracting spherical container of gas

30. A planet has elliptical orbit with sun at the focus as shown in the figure. Which position of the orbit the planet has the highest speed ?



- (A) A
- (B) B

31. If the Lagrangian of a particle of mass  $m$  is :

$$L(\rho, \theta, \dot{\rho}, \dot{\theta}) = \frac{m}{2} (\dot{\rho}^2 \dot{\theta}^2 + \dot{\rho}^2 \operatorname{cosec}^2 \alpha) - mg \rho \cot \alpha$$

then conserved quantities are :

- (A)  $p_\rho$
- (B)  $p_\rho$  and  $p_\theta$
- (C)  $p_\theta$
- (D) none of the above

32. Example of a non-central force is :

- (A) Gravitational force  $-\frac{Gm_1m_2}{r^2}\hat{r}$
- (B) Coulomb force  $\frac{z_1z_2}{r^2}\hat{r}$
- (C) Hooke law  $k\vec{r}$
- (D) dipole-dipole interaction  $\frac{\vec{p}\cdot\vec{r}}{r^3}$

where  $\vec{p}$  is the dipole



33. An infinitely long line-charge has a uniform linear charge density  $\lambda$ . If  $r$  denotes the distance of a point from the wire, then magnitude of the electric field at the point is :

- (A) proportional to  $\frac{1}{r}$
- (B) proportional to  $\frac{1}{r^2}$
- (C) proportional to  $\frac{1}{r^3}$
- (D) independent of  $r$

34. The dispersion relation for electromagnetic waves in a certain medium is  $\omega^2 = \alpha k$ , where  $\alpha$  is constant,  $\omega$  the angular frequency and  $k$  the magnitude of the wave vector. The velocity of the energy propagation by electromagnetic waves in this medium is :

- (A)  $\frac{\alpha}{\omega}$
- (B)  $\frac{2\alpha}{\omega}$
- (C)  $\frac{\alpha}{2\omega}$
- (D)  $\frac{\alpha}{\omega}$

35. The dispersion relation for electromagnetic waves in a medium is  $\omega^2 = \alpha k^2$ , where  $\alpha$  is constant,  $\omega$  the angular frequency and  $k$  the magnitude of the wave vector. Which of the following statements is correct ?

- (A) The phase velocity in the medium is  $\alpha$
- (B) The group velocity in the medium is  $\alpha$
- (C) The medium is dispersive
- (D) The medium is non-dispersive

36. The interaction energy of an electric dipole  $\vec{p}$  in an external electric field  $\vec{E}$  is :

- (A)  $\vec{p} \cdot \vec{E}$
- (B)  $-\vec{p} \cdot \vec{E}$
- (C)  $|\vec{p} \times \vec{E}|$
- (D)  $|\vec{p} \cdot \vec{E}|$

37. In the Young's double slit experiment, the intensity of central maximum is  $I_2$ . If either of the slits is closed, the intensity at the same location is  $I_1$ . The relation between  $I_1$  and  $I_2$  is :

- (A)  $I_2 = 4I_1$
- (B)  $I_2 = 2I_1$
- (C)  $I_2 = I_1$
- (D)  $I_1 = 2I_2$

38. Maxwell introduced an additional term in :

- (A) Gauss's law
- (B) Faraday's law
- (C) Ampere's law
- (D) Coulomb's law

39. The skin depth  $\delta$  of a good metal, for the microwave frequency  $\omega$  follows the relation :

- (A)  $\delta \propto \omega$
- (B)  $\delta \propto \frac{1}{\omega}$
- (C)  $\delta \propto \sqrt{\omega}$

40. For a certain material  $\frac{g}{v} \ll 1$ , where  $g$  and  $\epsilon$  are conductivity and permittivity of the medium.

radiation of frequency  $\omega$ , the material is :

- (A) a good conductor
- (B) a good insulator
- (C) partially insulating
- (D) a semiconductor

41. The entropy of an ideal gas at absolute zero is :

- (A)  $\infty$
- (B) 0
- (C)  $Nk_B$

(D) can not be calculated

42. For a system of  $N$  non-interacting fermions enclosed in a volume 'V' at constant temperature  $T$ , the average occupation number of the ' $r$ 'th energy level is given by :

(A)  $\bar{n}_r = \frac{1}{e^{\beta(\epsilon_r - \mu)} + 1}$

(B)  $\bar{n}_r = \frac{1}{(e^{\beta(\epsilon_r - \mu)} - 1)}$

(C)  $\bar{n}_r = e^{-\beta(\epsilon_r - \mu)}$

(D)  $\bar{n}_r = (e^{\beta(\epsilon_r - \mu)} + 1)$

43. A first order phase transition is characterised by :

- (A) a divergence of the specific heat at  $T_C$ , the critical temperature
- (B) A cusp in the average energy at  $T_C$
- (C) The constancy of entropy in the transition
- (D) A latent heat is involved in the transition process

44. A gas of molecules of mass ' $m$ ' is in thermal equilibrium at an absolute temperature ' $T$ '.

$v_x, v_y, v_z$  are the components of the velocity ' $\vec{v}$ ' of each molecule, then the mean value of  $v^2$  :

(A) 0

(B)  $\frac{1}{2} k_B T$

(C)  $\frac{3}{m} k_B T$

(D)  $N k_B T$

45. The Fermi energy of a free electron gas at absolute zero is of the order of :

(A) electron-volts

(B) MeV

(C) keV

46. Consider an ideal gas of  $N$  molecules enclosed in a volume ' $V$ ' maintained at a temperature ' $T$ '. The correct expression for the entropy of the system is :

(A)  $S = Nk_B \left[ \ln V + \frac{3}{2} \ln T + \sigma \right]$

(B)  $S = Nk_B \left[ \ln \left( \frac{V}{N} \right) + \frac{3}{2} \ln T + \sigma \right]$

(C)  $S = k_B \left[ \ln V + \frac{3}{2} \ln T + \sigma \right]$

(D)  $S = k_B \left( \frac{N}{V} \right) \left[ \ln \left( \frac{V}{N} \right) + \frac{3}{2} \ln T + \sigma \right]$

47. If the temperature of a black body is increased by a factor of 2, the amount of energy/volume radiated increases by a factor of :

- (A) 2
- (B) 4
- (C) 8

48. If the temperature of an electron gas is increased by a factor of 2, the specific heat increases by a factor of :

- (A) 2
- (B) 4
- (C) 8
- (D) 16

49. In Laue X-ray diffraction experiment in the study of single crystal structure, the following X-ray source is used :

- (A) Monochromatic
- (B) Non-monochromatic
- (C) Pulsed Monochromatic
- (D) Bichromatic

50. 5 boys and 3 girls are to stand in a straight line such that no two girls are adjacent. The number of ways in which this can be done is :

- (A) 5 !
- (B) 3 !
- (C) 5 ! × 3 !