

M. PHIL. IN STATISTICAL SCIENCE

---

Thursday 2 June 2005 1.30 to 4.30

---

STATISTICAL THEORY

Attempt **FOUR** questions, not more than **TWO** of which should be from Section B.

There are **TEN** questions in total.

The questions carry equal weight.

**STATIONERY REQUIREMENTS**

Cover sheet  
Treasury Tag  
Script paper

**SPECIAL REQUIREMENTS**

None

<p>You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.</p>
--

**Section A**

**1** Explain briefly the concepts of *profile likelihood* and *conditional likelihood*, for inference about a parameter of interest  $\psi$ , in the presence of a nuisance parameter  $\lambda$ .

Suppose  $Y_1, \dots, Y_n$  are independent, identically distributed from the exponential family density

$$f(y; \psi, \lambda) = \exp\{\psi\tau_1(y) + \lambda\tau_2(y) - d(\psi, \lambda) - Q(y)\},$$

where  $\psi, \lambda$  are both scalar.

Obtain a saddlepoint approximation to the density of  $S = n^{-1} \sum_{i=1}^n \tau_2(Y_i)$ .

Show that use of the saddlepoint approximation leads to an approximate conditional log-likelihood function for  $\psi$  of the form

$$l_p(\psi) + B(\psi),$$

where  $l_p(\psi)$  is the profile log-likelihood, and  $B(\psi)$  is an adjustment which you should specify carefully.

**2** Explain in detail what is meant by a *transformation model*.

What is meant by (i) a *maximal invariant*, (ii) an *equivariant estimator*, in the context of a transformation model?

Describe in detail how an equivariant estimator can be used to construct a maximal invariant. Illustrate the construction for the case of a *location-scale* model.

**3** Let  $Y_1, \dots, Y_n$  be independent, identically distributed  $N(\mu, \mu^2)$ ,  $\mu > 0$ .

Show that this model is an example of a *curved exponential family*, and find a minimal sufficient statistic.

Show that

$$a = \sqrt{n} \frac{(\sum Y_i^2)^{1/2}}{\sum Y_i}$$

is an ancillary statistic.

Assume that  $a > 0$ . Show that the maximum likelihood estimator of  $\mu$  is

$$\hat{\mu} = \frac{(\sum Y_i^2)^{1/2}}{q\sqrt{n}},$$

where  $q = \{(1 + 4a^2)^{1/2} + 1\}/(2a)$ .

Show further that (apart from a constant) the log-likelihood may be written

$$l(\mu; \hat{\mu}, a) = -\frac{n}{2\mu^2} \left( q^2 \hat{\mu}^2 - \frac{2q\mu\hat{\mu}}{a} \right) - n \log \mu,$$

and obtain the  $p^*$  approximation to the (conditional) density of  $\hat{\mu}$ .

How would you approximate  $\text{Prob}(\hat{\mu} \leq t|a)$ , for given  $t$ ?

**4** Explain what is meant by an *M-estimator* of a parameter  $\theta$ , based on a given  $\psi$  function. Show that under appropriate conditions allowing the interchange of order of integration and differentiation, the influence function is proportional to  $\psi$  and derive an expression for the asymptotic variance  $V(\psi, F)$  of the *M-estimator* at a distribution  $F$ .

A location model on  $\mathbb{R}$ , with parameter space  $\mathbb{R}$ , is specified by  $F_\theta(x) = F(x - \theta)$ , and an *M-estimator* is constructed using a  $\psi$  function of the form  $\psi(x, \theta) = \psi(x - \theta)$ .

For the particular choice

$$\psi(x) = \min\{b, \max\{x, -b\}\}, \quad b < \infty :$$

(i) Find the asymptotic variance  $V(\psi, \Phi)$ , where  $\Phi$  is the standard normal distribution;

(ii) Verify that the estimator is *B-robust*, by determining an explicit bound on the influence function.

**5** Let  $Y_1, \dots, Y_n$  be independent, identically distributed from a distribution  $F$ , with density  $f$  symmetric about an unknown point  $\theta$ . Suppose we wish to test  $H_0 : \theta = \theta_0$  against  $H_1 : \theta < \theta_0$ .

Explain how to test  $H_0$  against  $H_1$  using (i) the sign test, *and* (ii) the Wilcoxon signed rank test.

Show that the null mean and variance of the Wilcoxon signed rank statistic are  $\frac{1}{4}n(n+1)$  and  $\frac{1}{24}n(n+1)(2n+1)$  respectively.

What is meant by a one-sample  $U$ -statistic?

State, without proof, a result concerning the asymptotic distribution of a one-sample  $U$ -statistic, and use it to deduce asymptotic normality of the Wilcoxon signed rank statistic.

**6** Write brief notes on *four* of the following:

- (i) Edgeworth expansion;
- (ii) parameter orthogonality;
- (iii) Laplace approximation;
- (iv) Bartlett correction;
- (v) the invariance principle;
- (vi) finite-sample versions of robustness measures;
- (vii) tests based on the empirical distribution function;
- (viii) large-sample likelihood theory.

### Section B

7 Assume that the  $n$ -dimensional observation vector  $Y$  may be written

$$Y = X\beta + \epsilon,$$

where  $X$  is a given  $n \times p$  matrix of rank  $p$ ,  $\beta$  is an unknown vector, and

$$\epsilon \sim N_n(0, \sigma^2 I).$$

Let  $Q(\beta) = (Y - X\beta)^T(Y - X\beta)$ . Find  $\hat{\beta}$ , the least-squares estimator of  $\beta$ , and show that

$$Q(\hat{\beta}) = Y^T(I - H)Y$$

where  $H$  is a matrix that you should define.

If now  $X\beta$  is written as  $X\beta = X_1\beta_1 + X_2\beta_2$ , where  $X = (X_1 : X_2)$ ,  $\beta^T = (\beta_1^T : \beta_2^T)$ , and  $\beta_2$  is of dimension  $p_2$ , state without proof the form of the  $F$ -test for testing  $H_0 : \beta_2 = 0$ .

What is meant by saying that  $\beta_1$  is orthogonal to  $\beta_2$ ? What is the practical relevance of orthogonality?

8 Suppose that  $Y_1, \dots, Y_n$  are independent binomial observations, with

$$Y_i \sim B(t_i, \pi_i) \text{ and } \log(\pi_i/(1 - \pi_i)) = \beta^T x_i, \quad \text{for } 1 \leq i \leq n,$$

where  $t_1, \dots, t_n$  and  $x_1, \dots, x_n$  are given. Discuss carefully the estimation of  $\beta$ .

Your solution should include

(i) the method of checking the fit of the above logistic model,

and

(ii) the method for finding an approximate 95% confidence interval for  $\beta_2$ , the second component of the vector  $\beta$ .

**9** Write a brief account of Bayesian decision theory. Derive the general form of a Bayes rule, in terms of the posterior distribution and loss function.

Let  $X_1, \dots, X_n$ , conditional on  $\theta$ , be a random sample from a  $N(\theta, \sigma^2)$  population and let the prior distribution on  $\theta$  be  $N(\mu, \tau^2)$ , where the values of  $\sigma^2$ ,  $\mu$  and  $\tau^2$  are known. Consider testing  $H_0 : \theta \geq \theta_0$  against  $H_1 : \theta < \theta_0$ , with loss function,

$$L(\theta, \text{accept } H_0) = \begin{cases} 0 & \text{if } \theta \geq \theta_0 \\ 1 & \text{if } \theta < \theta_0 \end{cases}$$

and

$$L(\theta, \text{accept } H_1) = \begin{cases} 1 & \text{if } \theta \geq \theta_0 \\ 0 & \text{if } \theta < \theta_0 \end{cases}.$$

Show that the Bayes test rejects  $H_0$  if

$$\bar{x} < \theta_0 - \frac{\eta(\mu - \theta_0)}{1 - \eta},$$

where  $\eta = \sigma^2 / (n\tau^2 + \sigma^2)$  and  $\bar{x} = n^{-1} \sum_{i=1}^n x_i$ .

**10** Let  $X_1, \dots, X_n$  be independent identically distributed random variables, with joint density  $f(x; \theta)$ , where  $\theta$  is a scalar parameter.

What is meant by the *Fisher information*  $I_n(\theta)$  about  $\theta$  contained in the sample  $X_1, \dots, X_n$ ?

State the Cramer–Rao lower bound for the variance of an arbitrary unbiased estimator of  $\theta$ . Give the condition for this lower bound to be attained.

Let  $X_1, \dots, X_n$  be an independent, identically distributed sample of size  $n \geq 3$  from the exponential density  $\theta e^{-\theta x}$ ,  $x > 0$ , where  $\theta > 0$  is unknown.

Obtain the maximum likelihood estimator  $\hat{\theta}$  of  $\theta$ . Show that it is biased, but a multiple of it is not. Determine whether the Cramer–Rao lower bound is attained by the variance of this unbiased estimator.

State carefully, but without proof, the asymptotic distribution of  $\hat{\theta}$ .

Explain in detail the rationale for the Wald test of the hypothesis  $H_0 : \theta = \theta_0$ , and find the form of the test statistic for the above exponential example.

[Hint: You may assume that the density of  $X_1 + \dots + X_n$  is  $\frac{\theta^n t^{n-1} e^{-\theta t}}{(n-1)!}$ ,  $t > 0$ .]

**END OF PAPER**