

MATHEMATICAL TRIPOS Part IA

Thursday 29 May 2003 9 to 12

PAPER 1

Before you begin read these instructions carefully.

*Each question in Section II carries twice the credit of each question in Section I. In Section I, you may attempt **all four** questions. In Section II, at most **five** answers will be taken into account and no more than **three** answers on each course will be taken into account.*

Complete answers are preferred to fragments.

*Write on **one side** of the paper only and begin each answer on a separate sheet.*

Write legibly; otherwise you place yourself at a grave disadvantage.

At the end of the examination:

*Tie up your answers in bundles, marked **A, B, C, D** and **F** according to the code letter affixed to each question. Attach a blue cover sheet to each bundle; write the code letter in the box marked 'SECTION' on the cover sheet. Do not tie up questions from Section I and Section II in separate bundles.*

You must also complete a green master cover sheet listing all the questions you have attempted.

Every cover sheet must bear your examination number and desk number.

<p>You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.</p>

SECTION I

1B Algebra and Geometry

(a) Write the permutation

$$(123)(234)$$

as a product of disjoint cycles. Determine its order. Compute its sign.

(b) Elements x and y of a group G are *conjugate* if there exists a $g \in G$ such that $g x g^{-1} = y$.

Show that if permutations x and y are conjugate, then they have the same sign and the same order. Is the converse true? (Justify your answer with a proof or counterexample.)

2D Algebra and Geometry

Find the characteristic equation, the eigenvectors $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$, and the corresponding eigenvalues $\lambda_{\mathbf{a}}, \lambda_{\mathbf{b}}, \lambda_{\mathbf{c}}, \lambda_{\mathbf{d}}$ of the matrix

$$A = \begin{pmatrix} i & 1 & 0 & 0 \\ 1 & i & 0 & 0 \\ 0 & 0 & i & 1 \\ 0 & 0 & -1 & i \end{pmatrix}.$$

Show that $\{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}\}$ spans the complex vector space \mathbb{C}^4 .

Consider the four subspaces of \mathbb{C}^4 defined parametrically by

$$\mathbf{z} = s\mathbf{a}, \quad \mathbf{z} = s\mathbf{b}, \quad \mathbf{z} = s\mathbf{c}, \quad \mathbf{z} = s\mathbf{d} \quad (\mathbf{z} \in \mathbb{C}^4, s \in \mathbb{C}).$$

Show that multiplication by A maps three of these subspaces onto themselves, and the remaining subspace into a smaller subspace to be specified.

3B Analysis

Define what it means for a function of a real variable to be *differentiable* at $x \in \mathbb{R}$.

Prove that if a function is differentiable at $x \in \mathbb{R}$, then it is continuous there.

Show directly from the definition that the function

$$f(x) = \begin{cases} x^2 \sin(1/x) & x \neq 0 \\ 0 & x = 0 \end{cases}$$

is differentiable at 0 with derivative 0.

Show that the derivative $f'(x)$ is not continuous at 0.

4C Analysis

Explain what is meant by the *radius of convergence* of a power series.

Find the radius of convergence R of each of the following power series:

$$(i) \sum_{n=1}^{\infty} n^{-2} z^n, \quad (ii) \sum_{n=1}^{\infty} \left(n + \frac{1}{2^n} \right) z^n.$$

In each case, determine whether the series converges on the circle $|z| = R$.

SECTION II

5B Algebra and Geometry

(a) In the standard basis of \mathbb{R}^2 , write down the matrix for a rotation through an angle θ about the origin.

(b) Let A be a real 3×3 matrix such that $\det A = 1$ and $AA^T = I$, where A^T is the transpose of A .

- (i) Suppose that A has an eigenvector \mathbf{v} with eigenvalue 1. Show that A is a rotation through an angle θ about the line through the origin in the direction of \mathbf{v} , where $\cos \theta = \frac{1}{2}(\text{trace} A - 1)$.
- (ii) Show that A must have an eigenvector \mathbf{v} with eigenvalue 1.

6A Algebra and Geometry

Let α be a linear map

$$\alpha : \mathbb{R}^3 \rightarrow \mathbb{R}^3.$$

Define the kernel K and image I of α .

Let $\mathbf{y} \in \mathbb{R}^3$. Show that the equation $\alpha \mathbf{x} = \mathbf{y}$ has a solution $\mathbf{x} \in \mathbb{R}^3$ if and only if $\mathbf{y} \in I$.

Let α have the matrix

$$\begin{pmatrix} 1 & 1 & t \\ 0 & t & -2b \\ 1 & t & 0 \end{pmatrix}$$

with respect to the standard basis, where $b \in \mathbb{R}$ and t is a real variable. Find K and I for α . Hence, or by evaluating the determinant, show that if $0 < b < 2$ and $\mathbf{y} \in I$ then the equation $\alpha \mathbf{x} = \mathbf{y}$ has a unique solution $\mathbf{x} \in \mathbb{R}^3$ for all values of t .

7B Algebra and Geometry

(i) State the orbit-stabilizer theorem for a group G acting on a set X .

(ii) Denote the group of *all* symmetries of the cube by G . Using the orbit-stabilizer theorem, show that G has 48 elements.

Does G have any non-trivial normal subgroups?

Let L denote the line between two diagonally opposite vertices of the cube, and let

$$H = \{g \in G \mid gL = L\}$$

be the subgroup of symmetries that preserve the line. Show that H is isomorphic to the direct product $S_3 \times C_2$, where S_3 is the symmetric group on 3 letters and C_2 is the cyclic group of order 2.

8D Algebra and Geometry

Let $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$ be non-zero vectors in \mathbb{R}^n . What is meant by saying that \mathbf{x} and \mathbf{y} are linearly independent? What is the dimension of the subspace of \mathbb{R}^n spanned by \mathbf{x} and \mathbf{y} if they are (1) linearly independent, (2) linearly dependent?

Define the scalar product $\mathbf{x} \cdot \mathbf{y}$ for $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$. Define the corresponding norm $\|\mathbf{x}\|$ of $\mathbf{x} \in \mathbb{R}^n$. State and prove the Cauchy–Schwarz inequality, and deduce the triangle inequality.

By means of a sketch, give a geometric interpretation of the scalar product $\mathbf{x} \cdot \mathbf{y}$ in the case $n = 3$, relating the value of $\mathbf{x} \cdot \mathbf{y}$ to the angle α between the directions of \mathbf{x} and \mathbf{y} .

What is a unit vector? Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be unit vectors in \mathbb{R}^3 . Let

$$S = \mathbf{u} \cdot \mathbf{v} + \mathbf{v} \cdot \mathbf{w} + \mathbf{w} \cdot \mathbf{u}.$$

Show that

- (i) for any fixed, linearly independent \mathbf{u} and \mathbf{v} , the minimum of S over \mathbf{w} is attained when $\mathbf{w} = \lambda(\mathbf{u} + \mathbf{v})$ for some $\lambda \in \mathbb{R}$;
- (ii) $\lambda \leq -\frac{1}{2}$ in all cases;
- (iii) $\lambda = -1$ and $S = -3/2$ in the case where $\mathbf{u} \cdot \mathbf{v} = \cos(2\pi/3)$.

9F Analysis

Prove the Axiom of Archimedes.

Let x be a real number in $[0, 1]$, and let m, n be positive integers. Show that the limit

$$\lim_{m \rightarrow \infty} \left[\lim_{n \rightarrow \infty} \cos^{2n}(m! \pi x) \right]$$

exists, and that its value depends on whether x is rational or irrational.

[You may assume standard properties of the cosine function provided they are clearly stated.]

10F Analysis

State without proof the *Integral Comparison Test* for the convergence of a series $\sum_{n=1}^{\infty} a_n$ of non-negative terms.

Determine for which positive real numbers α the series $\sum_{n=1}^{\infty} n^{-\alpha}$ converges.

In each of the following cases determine whether the series is convergent or divergent:

$$(i) \sum_{n=3}^{\infty} \frac{1}{n \log n},$$

$$(ii) \sum_{n=3}^{\infty} \frac{1}{(n \log n) (\log \log n)^2},$$

$$(iii) \sum_{n=3}^{\infty} \frac{1}{n^{(1+1/n)} \log n}.$$

11B Analysis

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Define the *integral* $\int_a^b f(x) dx$. (You are not asked to prove existence.)

Suppose that m, M are real numbers such that $m \leq f(x) \leq M$ for all $x \in [a, b]$. Stating clearly any properties of the integral that you require, show that

$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a).$$

The function $g : [a, b] \rightarrow \mathbb{R}$ is continuous and non-negative. Show that

$$m \int_a^b g(x) dx \leq \int_a^b f(x) g(x) dx \leq M \int_a^b g(x) dx.$$

Now let f be continuous on $[0, 1]$. By suitable choice of g show that

$$\lim_{n \rightarrow \infty} \int_0^{1/\sqrt{n}} n f(x) e^{-nx} dx = f(0),$$

and by making an appropriate change of variable, or otherwise, show that

$$\lim_{n \rightarrow \infty} \int_0^1 n f(x) e^{-nx} dx = f(0).$$

12C Analysis

State carefully the formula for integration by parts for functions of a real variable.

Let $f : (-1, 1) \rightarrow \mathbb{R}$ be infinitely differentiable. Prove that for all $n \geq 1$ and all $t \in (-1, 1)$,

$$f(t) = f(0) + f'(0)t + \frac{1}{2!}f''(0)t^2 + \dots + \frac{1}{(n-1)!}f^{(n-1)}(0)t^{n-1} + \frac{1}{(n-1)!} \int_0^t f^{(n)}(x)(t-x)^{n-1} dx.$$

By considering the function $f(x) = \log(1-x)$ at $x = 1/2$, or otherwise, prove that the series

$$\sum_{n=1}^{\infty} \frac{1}{n 2^n}$$

converges to $\log 2$.