

### MATHEMATICAL TRIPOS Part III

Wednesday 13 June 2001 9 to 12

# **PAPER 74**

## ALGEBRAIC NUMBER THEORY

Attempt **FIVE** questions, including at least **ONE** from each section. All questions carry equal weight.

> You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.

 $\mathbf{2}$ 

#### Section A

1 Give the definition of a Dedekind domain. Let  $\mathfrak{o}$  be a Dedekind domain with field of fractions k. Let  $\mathfrak{a}$  be a non-zero fractional ideal in k and define  $\mathfrak{a}^{-1} = \{ x \in k \mid x\mathfrak{a} \subset \mathfrak{o} \}$ . Show that  $\mathfrak{a}^{-1}$  is a fractional ideal and that  $\mathfrak{a}\mathfrak{a}^{-1} = \mathfrak{o}$ .

 $\mathbf{2}$  (i) State and prove the Chinese remainder theorem for  $\mathfrak{o}$  a Dedekind domain.

(ii) Let K/k be a normal extension of algebraic number fields. Let  $\mathfrak{p}$  be a prime of k, whose factorisation in K is  $\operatorname{conorm}_{K/k} \mathfrak{p} = \mathfrak{P}_1^{e_1} \dots \mathfrak{P}_g^{e_g}$ . Show that  $\operatorname{Gal}(K/k)$  acts transitively on the  $\mathfrak{P}_i$ .

**3** Let K/k be a finite extension of algebraic number fields. Define the relative ideal norm and prove that it is multiplicative. Let  $\mathfrak{p}$  be a prime of k, whose factorisation in K is conorm<sub>K/k</sub>  $\mathfrak{p} = \mathfrak{P}_1^{e_1} \dots \mathfrak{P}_g^{e_g}$ . Show that  $[K:k] = \sum e_i f_i$  where  $f_i$  is the degree of  $\mathfrak{O}/\mathfrak{P}_i$  over  $\mathfrak{o}/\mathfrak{p}$ .

[Properties of the norm for elements may be assumed.]

#### Section B

4 Let K/k be an extension of algebraic number fields. Let  $\mathfrak{p}$  be a prime of k and  $\mathfrak{P}$  a prime of K above  $\mathfrak{p}$ .

(i) Let f(X) be a monic polynomial in  $\mathfrak{o}_{\mathfrak{p}}[X]$  and suppose that the reduction mod  $\mathfrak{p}$  factors as  $\tilde{f}(X) = \phi_1(X)\phi_2(X)$  where  $\phi_1, \phi_2$  in  $(\mathfrak{o}/\mathfrak{p})[X]$  are coprime. Show that  $f(X) = f_1(X)f_2(X)$  with  $\tilde{f}_{\nu}(X) = \phi_{\nu}(X)$ .

(ii) Suppose  $\mathfrak{P}^e \mid | \mathfrak{p}$  and  $\mathfrak{p} \mid e$ . Show that if  $\alpha \in \mathfrak{O}_{\mathfrak{P}}$  then  $\operatorname{Tr}_{K_{\mathfrak{P}}/k_{\mathfrak{p}}}(\alpha) \in \mathfrak{p}_{\mathfrak{p}}$ .

5 Let K/k be an extension of algebraic number fields. Define the relative different  $\mathfrak{d}_{K/k}$ . In the case  $k = \mathbf{Q}$  describe the relationship with the discriminant  $d_K$ .

(i) For  $K \supset L \supset k$  show that  $\mathfrak{d}_{K/k} = \mathfrak{d}_{K/L} \mathfrak{d}_{L/k}$ .

(ii) State a relationship between the different and ramification. Hence show that if  $K_1, K_2$  are Galois over **Q** with coprime discriminants, then  $[K_1K_2 : \mathbf{Q}] = [K_1 : \mathbf{Q}][K_2 : \mathbf{Q}]$ .

Paper 74

3

#### Section C

6

Write an essay on the Hilbert class field. Illustrate by computing the Hilbert class field for *either*  $\mathbf{Q}(\sqrt{-23})$  or  $\mathbf{Q}(\sqrt{-30})$ , explaining all necessary working.

[The cubic  $X^3 + aX + b$  has discriminant  $-4a^3 - 27b^2$ .]

7 Let  $m = m_1 m_2^2$  with  $m_1$ ,  $m_2$  coprime square-free positive integers. Suppose  $m_1 \not\equiv \pm m_2 \pmod{9}$ . Show that  $\mathbf{Q}(\sqrt[3]{m})$  has discriminant  $-27m_1^2m_2^2$ . Find a unit in  $\mathbf{Q}(\sqrt[3]{6})$  and show that this field has class number h = 1.

8 Let K/k be a quadratic extension of algebraic number fields with K totally complex and k totally real.

(i) Show that  $[\mathfrak{O}_K^* : \mathfrak{o}_k^* \mu_K] = 1$  or 2, where  $\mathfrak{O}_K^*$ ,  $\mathfrak{o}_k^*$  are the unit groups in K, k, and  $\mu_K$  is the group of roots of unity in K.

(ii) Show that the class number of k divides the class number of K.

[You may assume any properties of the Hilbert class field you require.]

Paper 74