

MATHEMATICAL TRIPOS Part III

Thursday 3 June, 2004 9 to 12

PAPER 58

APPLICATIONS OF DIFFERENTIAL GEOMETRY TO PHYSICS

*Attempt **FOUR** questions.*

*There are **seven** questions in total.*

The questions carry equal weight.

**You may not start to read the questions
printed on the subsequent pages until
instructed to do so by the Invigilator.**

1 Define the Killing form B on the Lie algebra $\mathfrak{g} = \text{Lie}(G)$ of a Lie group G . Give a condition on G that the Killing form B be non-degenerate (i.e. invertible). Explain how, in this case, one may endow G with a bi-invariant pseudo-riemannian metric and show that this metric is an Einstein metric.

Comment briefly, giving illustrations but no proofs, on the signature of the Killing form B and its relation to the topology of the group G .

2 Define the terms *orbit*, *stabilizer*, *transitive*, *simply transitive* and *multiply transitive* for the action of a group G acting on a manifold X . Show that if the action is multiply transitive, then $X = G/H$ for some subgroup $H \subset G$. How is the subgroup H determined?

Show that the space of quantum states of an $(n + 1)$ -state quantum system is complex projective space $\mathbb{C}\mathbb{P}^n$. Show that $\mathbb{C}\mathbb{P}^n \cong U(n + 1)/U(n) \times U(1)$.

Give a description of S^{2n+1} as an S^1 bundle over $\mathbb{C}\mathbb{P}^n$.

3 Define a Poisson manifold $\{P, \omega^{\mu\nu}\}$ and derive the condition that the second rank antisymmetric contravariant tensor field $\omega^{\mu\nu}$ must satisfy in order that the associated Poisson bracket satisfies the Jacobi identity.

Define a symplectic manifold, and show that every symplectic manifold is a Poisson manifold.

Show, by means of an example based on the dual \mathfrak{g}^* of a Lie algebra \mathfrak{g} , that not every Poisson manifold, equipped with a Poisson bracket satisfying the Jacobi identity, need be a symplectic manifold.

4 The metric of three-dimensional Anti-de-Sitter spacetime AdS_3 may be written in globally static coordinates as

$$ds^2 = -(1+r^2)dt^2 + \frac{dr^2}{1+r^2} + r^2d\theta^2,$$

with both t and θ being periodic coordinates of period 2π . By means of the embedding into $\mathbb{E}^{2,2}$ given by

$$\begin{aligned} X^0 &= \sqrt{1+r^2} \sin t, & X^4 &= \sqrt{1+r^2} \cos t, \\ X^1 &= r \cos \theta, & X^2 &= r \sin \theta, \end{aligned}$$

show that the metric coincides, up to a constant factor, with the Killing metric on $SL(2, \mathbb{R})$.

Show how $SO(2, 2)$ acts by isometries and construct a $2 : 1$ homomorphism from $SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$ to $SO(2, 2)$.

What can you say about the group $SO(2, 1)$?

5 Define a principal fibre bundle and show that it admits a global right action of the structural group G . Show also that such a bundle admits a global section if and only if it is trivial.

Illustrate your answer by means of the bundle of pseudo-orthonormal frames of a pseudo-riemannian manifold.

In particular, show that $SO(4, 1)$ is the bundle of orthonormal frames for four-dimensional De-Sitter spacetime dS_4 .

Given that S^3 admits a global frame field, does the bundle of pseudo-orthonormal frames for dS_4 admit a global section?

6 Let $\{M, g\}$ be a pseudo-riemannian manifold with a local pseudo-orthonormal basis for the tangent space \mathbf{e}_a and dual basis ω^a . Establish the equations

$$\begin{aligned} \theta_{ab} + \theta_{ba} &= 0, \\ d\omega^a + \theta^a_b \wedge \omega^b &= 0, \\ R^a_b &= d\theta^a_b + \theta^a_c \wedge \theta^c_b, \end{aligned}$$

where $\nabla \mathbf{e}_a = \mathbf{e}_b \otimes \theta^b_a$, and R^a_b is the curvature 2-form of the Levi-Civita connection.

How does R^a_b change under change of basis?

7 Write a brief essay on integration on manifolds, giving applications to topological conservation laws and the gauge-invariant coupling of a p -brane to a $(p+1)$ -form potential.