

PAPER 76

BRAIN IMAGING METHODS VIA ELECTRO-
AND MAGNETO- ENCEPHALOGRAPHY

*Attempt no more than **FOUR** questions.*

*There are **FIVE** questions in total.*

The questions carry equal weight.

STATIONERY REQUIREMENTS

*Cover sheet
Treasury Tag
Script paper*

SPECIAL REQUIREMENTS

Auxiliary Formulae data sheet

**You may not start to read the questions
printed on the subsequent pages until
instructed to do so by the Invigilator.**

1 In the spherical model of the brain with a dipole excitation $(\mathbf{r}_0, \mathbf{Q})$, the exterior electric and magnetic potentials are given by

$$u^+(\mathbf{r}) = \frac{1}{4\pi\sigma} \mathbf{Q} \cdot \left[2 \frac{\mathbf{P}}{P^3} + \frac{1}{rP} \frac{P\mathbf{r} + r\mathbf{P}}{Pr + \mathbf{r} \cdot \mathbf{P}} \right], \quad r > a$$

and

$$U(\mathbf{r}) = \frac{\mu_0}{4\pi} \frac{\mathbf{Q} \times \mathbf{r}_0 \cdot \mathbf{r}}{rP^2 + P\mathbf{r} \cdot \mathbf{P}}, \quad r > a$$

where $\mathbf{P} = \mathbf{r} - \mathbf{r}_0$ and a is the radius of the sphere. Find the singularities of u^+ and U in \mathbb{R}^3 , as well as their order.

2 Use the Geselowitz formula for the magnetic field together with the potential representation

$$\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \nabla U(\mathbf{r}), \quad r > a$$

to construct a function ϕ in closed form such that

$$U(\mathbf{r}) = \mathbf{Q} \times \mathbf{r}_0 \cdot \nabla_{\mathbf{r}_0} \phi(\mathbf{r}), \quad r > a$$

for a spherical conductor of radius a .

3 The interior electric potential for the spherical model of the brain with a dipole $(\mathbf{Q}, \mathbf{r}_0)$ is given by

$$u^-(\mathbf{r}) = \frac{1}{4\pi\sigma} \mathbf{Q} \cdot \left[\frac{\mathbf{r} - \mathbf{r}_0}{|\mathbf{r} - \mathbf{r}_0|^3} + \frac{\bar{\mathbf{r}}}{a} \frac{\bar{\mathbf{r}} - \mathbf{r}_0}{|\bar{\mathbf{r}} - \mathbf{r}_0|^3} \right] + \frac{1}{4\pi\sigma} \mathbf{Q} \cdot \frac{1}{aR} \frac{R\bar{\mathbf{r}} + \bar{\mathbf{r}}\mathbf{R}}{R\bar{\mathbf{r}} + \bar{\mathbf{r}} \cdot \mathbf{R}}$$

where $\bar{\mathbf{r}} = (a/r)^2 \mathbf{r}$ and $\mathbf{R} = \bar{\mathbf{r}} - \mathbf{r}_0$. Show that u^- can be interpreted in terms of images as

$$u^-(\mathbf{r}) = \Psi(\mathbf{r}; \mathbf{r}_0) + \left(\frac{r}{a}\right)^3 \Psi\left(\mathbf{r}; \left(\frac{r}{a}\right)^2 \mathbf{r}_0\right) + \frac{r}{a} \int_0^{\left(\frac{r}{a}\right)^2} \Psi(\mathbf{r}; t\mathbf{r}_0) dt$$

where

$$\Psi(\mathbf{r}; \mathbf{r}_0) = \frac{1}{4\pi\sigma} \mathbf{Q} \cdot \frac{\mathbf{r} - \mathbf{r}_0}{|\mathbf{r} - \mathbf{r}_0|^3}$$

4 Prove that the Kelvin transformation

$$\mathbf{r} \rightarrow \bar{\mathbf{r}} = \frac{a^2}{r^2} \mathbf{r}$$

leaves the operator $\mathbf{r} \times \nabla_{\mathbf{r}}$ invariant. Compute the effect of Kelvin's transformation on the following system of vector spherical harmonics:

$$\begin{aligned} \mathbf{P} &= \hat{\mathbf{r}}Y \\ \mathbf{B} &= \frac{r}{\sqrt{n(n+1)}} \nabla Y \\ \mathbf{C} &= -\frac{r}{\sqrt{n(n+1)}} \hat{\mathbf{r}} \times \nabla Y \end{aligned}$$

where Y is any scalar spherical harmonic and for convenience, the indices have been dropped.

5 Prove that the function $H_n(\mathbf{r})r^{-(2n+1)}$ is harmonic if and only if the n th degree homogeneous polynomial $H_n(\mathbf{r})$ is harmonic.

END OF PAPER

MATHEMATICAL TRIPOS, PART III

Paper 76: Brain Imaging Methods via Electro- and Magneto- Encephalography

Auxiliary Formulae data sheet

1.

$$F(\mathbf{r}; \mathbf{r}_0) = r|\mathbf{r} - \mathbf{r}_0|^2 + |\mathbf{r} - \mathbf{r}_0|\mathbf{r} \cdot (\mathbf{r} - \mathbf{r}_0)$$

2.

$$(1 - 2\gamma p + p^2)^{-1/2} = \sum_{n=0}^{\infty} p^n P_n(\gamma)$$

 P_n : Legendre polynomials, $|p| < 1$, $|\gamma| \leq 1$

3.

$$\sum_{n=1}^{\infty} \frac{1}{n} p^n P_n(\gamma) = -\ln \frac{\sqrt{1 - 2\gamma p + p^2} + 1 - \gamma p}{2}$$

4.

$$\sum_{n=0}^{\infty} \frac{1}{n+1} p^{n+1} P_n(\gamma) = \ln \frac{\sqrt{1 - 2\gamma p + p^2} + p - \gamma}{1 - \gamma}$$

5.

$$\Delta u(\mathbf{r}) = \left(\frac{\bar{r}}{a}\right)^5 \bar{\Delta} \left[\frac{a}{\bar{r}} u \left(\frac{a^2}{\bar{r}^2} \bar{\mathbf{r}} \right) \right]$$

6.

$$\nabla = \hat{\mathbf{r}} \frac{\partial}{\partial r} + \frac{\hat{\boldsymbol{\theta}}}{r} \frac{\partial}{\partial \theta} + \frac{\hat{\boldsymbol{\phi}}}{r \sin \theta} \frac{\partial}{\partial \phi}$$

7.

$$\int \frac{dr}{|\mathbf{r} - \mathbf{r}_0|} = \ln (|\mathbf{r} - \mathbf{r}_0| + \hat{\mathbf{r}} \cdot (\mathbf{r} - \mathbf{r}_0))$$

8.

$$\frac{|\mathbf{r} - \mathbf{r}_0|\mathbf{r} + r(\mathbf{r} - \mathbf{r}_0)}{F(\mathbf{r}; \mathbf{r}_0)} = \nabla_{\mathbf{r}_0} \ln \frac{|\mathbf{r} - \mathbf{r}_0|}{F(\mathbf{r}; \mathbf{r}_0)}$$

9.

$$P_0(\gamma) = 1, \quad P_1(\gamma) = \gamma, \quad P_2(\gamma) = \frac{3\gamma^2 - 1}{2}$$

10.

$$u(\mathbf{r}) = \frac{1}{4\pi\sigma} \mathbf{Q} \cdot \frac{\mathbf{r} - \mathbf{r}_0}{|\mathbf{r} - \mathbf{r}_0|^3} - \frac{1}{4\pi} \oint_S u^-(\mathbf{r}') \hat{\mathbf{n}}' \cdot \frac{\mathbf{r} - \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|^3} ds(\mathbf{r}'), \quad \forall \mathbf{r} \notin S$$

11.

$$\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \mathbf{Q} \times \frac{\mathbf{r} - \mathbf{r}_0}{|\mathbf{r} - \mathbf{r}_0|^3} - \frac{\mu_0 \sigma}{4\pi} \oint_S u^-(\mathbf{r}') \hat{\mathbf{n}}' \times \frac{\mathbf{r} - \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|^3} ds(\mathbf{r}'), \quad \forall \mathbf{r} \notin S$$

12.

$$\begin{aligned} \sigma \Delta u^-(\mathbf{r}) &= \nabla \cdot \mathbf{J}^p(\mathbf{r}), & \mathbf{r} \in \Omega \\ \partial_n u^-(\mathbf{r}) &= 0, & \mathbf{r} \in \partial\Omega \end{aligned}$$

13.

$$\begin{aligned} \nabla \cdot (f\mathbf{g}) &= (\nabla f) \cdot \mathbf{g} + f \nabla \cdot \mathbf{g} \\ \nabla \times (f\mathbf{g}) &= (\nabla f) \times \mathbf{g} + f \nabla \times \mathbf{g} \\ \nabla \times (\nabla \times \mathbf{g}) &= \nabla \nabla \cdot \mathbf{g} - \Delta \mathbf{g} \end{aligned}$$

14.

$$P_n(\hat{\mathbf{r}} \cdot \hat{\mathbf{r}}_0) = \frac{4\pi}{2n+1} \sum_{m=-n}^n \hat{Y}_n^m(\hat{\mathbf{r}}) \hat{Y}_n^{m*}(\hat{\mathbf{r}}_0)$$

\hat{Y}_n^{m*} : orthonormalised complex for of spherical harmonics.