

MATHEMATICAL TRIPOS Part III

Thursday 29 May 2008 1.30 to 4.30

PAPER 26

CATEGORY THEORY

*You should attempt **one** question from Section 1, and **two** from Section 2.*

*There are **six** questions in total.*

The questions carry equal weight.

STATIONERY REQUIREMENTS

*Cover sheet
Treasury Tag
Script paper*

SPECIAL REQUIREMENTS

None

<p>You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.</p>

SECTION 1

1 Explain what is meant by the terms *monad*, *Eilenberg–Moore category* and *monadic adjunction*. State and prove the Precise Monadicity Theorem, and use it to show that the adjunction formed by the forgetful functor from the category of compact Hausdorff spaces to **Set**, and its left adjoint, is monadic.

[Standard results from general topology, and the existence of the left adjoint, may be assumed.]

2 It has been said that category theory is the one part of mathematics where definitions matter more than theorems. Write a short essay arguing the case *either* for *or* against this assertion, illustrating your argument with examples drawn from the course.

SECTION 2

3 Let \mathcal{C} be a small category and $F: \mathcal{C} \rightarrow \mathbf{Set}$ a functor. Explain what is meant by the arrow category $(A \downarrow F)$, where A is a fixed set.

Show that F may be expressed as the colimit of a diagram of shape $(1 \downarrow F)^{\text{op}}$ in $[\mathcal{C}, \mathbf{Set}]$ (where 1 denotes a one-element set) whose vertices are representable functors. Deduce that if \mathcal{C} has finite limits, the following conditions are equivalent:

- (i) F preserves finite limits.
- (ii) For any set A , $(A \downarrow F)$ has finite limits.
- (iii) $(1 \downarrow F)^{\text{op}}$ is filtered.

[You may assume the result that filtered colimits commute with finite limits in **Set**.]

4 Define a *balanced* category. If $F: \mathcal{C} \rightarrow \mathcal{D}$ is a faithful functor and \mathcal{C} is balanced, prove that F reflects isomorphisms.

Let $((F: \mathcal{C} \rightarrow \mathcal{D}) \dashv (G: \mathcal{D} \rightarrow \mathcal{C}))$ be an adjunction with unit η and counit ϵ . Show that F is faithful if and only if η is a (pointwise) monomorphism. Now suppose that \mathcal{C} is balanced, and that every morphism of \mathcal{D} can be factored as a regular epimorphism followed by a monomorphism. Show that the following are equivalent:

- (i) Both η and ϵ are monomorphisms.
- (ii) F is full and faithful, and its image is closed (up to isomorphism) under regular quotients in \mathcal{D} . (That is, if $FA \rightarrow B$ is regular epic, then B is isomorphic to some FA' .)

Give an example of an adjunction whose unit and counit are both monic, but whose left adjoint is not full.

5 Let \mathcal{C} be a category, and let \mathcal{D} be a full subcategory of the functor category $[\mathcal{C}, \mathcal{C}]$ which is closed under composition and contains the identity functor. Suppose \mathcal{D} has a terminal object T : show that T carries a unique monad structure \mathbf{T} , and that if \mathbf{S} is any monad on \mathcal{C} whose functor part lies in \mathcal{D} then there is a forgetful functor $\mathcal{C}^{\mathbf{T}} \rightarrow \mathcal{C}^{\mathbf{S}}$.

Now let $\mathcal{C} = \mathbf{Set}$, and let \mathcal{D} be the category of all functors $\mathbf{Set} \rightarrow \mathbf{Set}$ which preserve finite coproducts. Show that \mathcal{D} has a terminal object T , and that TA may be identified with the set of all ultrafilters on A .

[Recall that an *ultrafilter* on a set A is a family \mathcal{F} of subsets of A satisfying $(B \in \mathcal{F}, B \subseteq C \Rightarrow C \in \mathcal{F})$, $(B \in \mathcal{F}, C \in \mathcal{F} \Rightarrow (B \cap C) \in \mathcal{F})$ and (for all $B \subseteq A$, exactly one of B and $A \setminus B$ is in \mathcal{F}).]

6 Define the notions of *abelian category* and of *exact sequence* in an abelian category. Prove that every morphism in an abelian category may be factored as an epimorphism followed by a monomorphism. Show also that, in an exact sequence

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0,$$

the morphism $A \rightarrow B$ is split monic if and only if $B \rightarrow C$ is split epic.

Let \mathcal{A} be an abelian category with enough projectives (i.e., such that every object admits an epimorphism from a projective object). Sketch the construction of the left derived functors $L^n F$ of a right exact functor $F: \mathcal{A} \rightarrow \mathcal{B}$, where \mathcal{B} is any abelian category, and write down the long exact sequence in \mathcal{B} induced by an exact sequence $(0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0)$ in \mathcal{A} . [Detailed proofs are not required: in particular, you need not prove that the $L^n F$ are well-defined or functorial.]

Show that an object A of \mathcal{A} is projective if and only if $L^1 F A = 0$ for all right exact $F: \mathcal{A} \rightarrow \mathcal{B}$. [Hint: given A , consider the functor $\mathcal{A}(-, K): \mathcal{A} \rightarrow \mathbf{AbGp}^{\text{op}}$, where $(0 \rightarrow K \rightarrow P \rightarrow A \rightarrow 0)$ is an exact sequence with P projective.] Deduce that the following conditions on \mathcal{A} are equivalent:

- (i) Every subobject of a projective object is projective.
- (ii) For any right exact $F: \mathcal{A} \rightarrow \mathcal{B}$, $L^2 F$ is identically 0.
- (iii) For any right exact $F: \mathcal{A} \rightarrow \mathcal{B}$, $L^1 F$ is left exact.

END OF PAPER