

PAPER 20

CONSTRUCTIVE TOPOLOGY

*Attempt **THREE** questions.*

*There are **five** questions in total.*

*The questions carry equal weight.*

**You may not start to read the questions  
printed on the subsequent pages until  
instructed to do so by the Invigilator.**

**1** Explain what is meant by the terms *nucleus*, *open nucleus* and *closed nucleus* on a frame  $A$ . Assuming the result that the nuclei on  $A$  form a frame  $N(A)$  under pointwise ordering, show that the open and closed nuclei  $o(a)$  and  $c(a)$  are complementary elements of this frame, and that the set of all open or closed nuclei generates  $N(A)$  as a frame.

Now suppose given a frame homomorphism  $h: B \rightarrow A$ . We say two nuclei  $j$  and  $k$  on  $A$  are *B-equivalent* if  $j(h(b)) = k(h(b))$  for all  $b \in B$ . By considering the nucleus  $\bigvee \{o(h(b)) \wedge c(j(h(b))) \mid b \in B\}$  (where  $j$  is an arbitrary nucleus), show that each *B-equivalence class* has a least element. Show also that these least elements form a subframe  $N_B(A)$  of  $N(A)$ , generated by the elements  $\{c(a) \mid a \in A\} \cup \{o(h(b)) \mid b \in B\}$ . Deduce that  $c: A \rightarrow N_B(A)$  is a frame homomorphism, and that it is the pushout of  $c: B \rightarrow N(B)$  along  $h$  in **Frm**.

[You may assume the results that  $o(a)$  is the least nucleus sending  $a$  to 1, and  $c(a)$  is the least sending 0 to  $a$ .]

**2** Explain what is meant by a *preframe*, and define the tensor product of two preframes. Assuming the result that the category **PFrm** of preframes is symmetric monoidal closed, show that the category **Frm** is coreflective in the category **CMon(PFrm)** of commutative monoids in **PFrm**, and show how this result may be used to give descriptions of binary coproducts and filtered colimits in **Frm** in terms of preframes. Hence prove the Tychonoff theorem that an arbitrary product of compact locales is compact.

**3** Define the terms *Hausdorff* and  $T_U$  for locales, and show that every Hausdorff locale is  $T_U$ . Give an example of a space which is Hausdorff as a space, but not  $T_U$  as a locale.

An open locale  $X$  is said to be *totally connected* if every positive open sublocale of  $X$  is connected. Show that the collection of all positive opens in a totally connected locale is a completely prime filter, and that the corresponding point  $p: 1 \rightarrow X$  is a dense sublocale of  $X$ . Conversely, if  $X$  is open and has a point which is a dense sublocale, show that it is totally connected. Show also that any continuous map from a totally connected locale  $X$  to a  $T_U$  locale  $Y$  is constant (i.e., factors through  $X \rightarrow 1$ ). [Hint: show that the composite  $px: X \rightarrow 1 \rightarrow X$  satisfies  $1_X \leq px$ .]

[If you wish, you may assume classical logic when answering this question, but full credit will be given only for constructively valid answers.]

4 Show that the following conditions on a locale  $X$  are equivalent:

- (i) The closure of every open sublocale of  $X$  is clopen (i.e. simultaneously open and closed).
- (ii) Every regular open sublocale of  $X$  (i.e. every sublocale which is the interior of its closure) is clopen.
- (iii) The mapping  $\neg\neg: \mathcal{O}(X) \rightarrow \mathcal{O}(X)$  preserves binary joins.
- (iv) The Boolean part  $X_b$  of  $X$  is a flat sublocale of  $X$ .

A locale with these properties is said to be *extremally disconnected*. Show also that if  $\mathcal{O}(X)$  is the frame  $\text{Idl}(B)$  of ideals of a Boolean algebra  $B$ , then  $X$  is extremally disconnected iff  $B$  is complete.

Now let  $X$  be a compact regular locale, and consider the locale  $\gamma X$  corresponding to the frame  $\text{Idl}(\mathcal{O}(X)_{\neg\neg})$  of ideals of the Boolean algebra of regular open sublocales of  $X$ . Show that  $\gamma X$  is extremally disconnected, and there is a surjection  $\gamma X \rightarrow X$ . [*Joyal's Lemma may be assumed.*]

5 Define a *propositional geometric theory*, and explain how any such theory  $\mathbb{T}$  generates a frame  $\mathcal{O}(X_{\mathbb{T}})$  such that the category  $\mathbf{Sh}(X_{\mathbb{T}})$  contains a canonical model of  $\mathbb{T}$ .

Given a commutative ring  $R$  (with 1), consider the geometric theory  $\mathbb{T}_R$  of ‘prime filters’ of  $R$ , whose primitive propositions have the form  $(r \in \mathcal{F})$ ,  $r \in R$ , and whose axioms are  $(\top \vdash (1 \in \mathcal{F}))$ ,

$$((r \in \mathcal{F}) \wedge (s \in \mathcal{F}) \vdash (rs \in \mathcal{F})),$$

$((rs \in \mathcal{F}) \vdash (r \in \mathcal{F}))$ ,  $((0 \in \mathcal{F}) \vdash \perp)$  and

$$(((r + s) \in \mathcal{F}) \vdash (r \in \mathcal{F}) \vee (s \in \mathcal{F}))$$

for all  $r, s \in R$ . (Thus, classically, a model of  $\mathbb{T}_R$  in  $\mathbf{Set}$  is the complement of a prime ideal of  $R$ .) Show that any ring homomorphism  $R \rightarrow S$  induces a locale map  $X_{\mathbb{T}_S} \rightarrow X_{\mathbb{T}_R}$ . If  $U_r$  denotes the open sublocale of  $X_{\mathbb{T}_R}$  corresponding to the primitive proposition  $(r \in \mathcal{F})$ , show that  $U_r \cup U_s$  is the whole of  $X_{\mathbb{T}_R}$  iff there exist  $a, b \in R$  such that  $ar + bs = 1$ . [*Hint: for the necessity of this condition, consider the homomorphism  $R \rightarrow S$ , where  $S$  is the quotient of  $R$  by the ideal generated by  $r$  and  $s$ . You may assume that  $X_{\mathbb{T}_S}$  is degenerate iff  $0 = 1$  in  $S$ .]*

Hence show that there is a sheaf  $\tilde{R}$  on  $X_{\mathbb{T}_R}$  whose elements of extent  $U_r$ , for each  $r$ , are the elements of the ring of fractions  $R[r^{-1}]$  (that is, equivalence classes of formal fractions  $a/r^n$  where  $a \in R$  and  $n \geq 0$ , where we identify  $a/r^n$  and  $b/r^m$  if there exists  $k \geq 0$  such that  $ar^{m+k} = br^{n+k}$ ). [*You may assume that it suffices to verify the sheaf axiom for coverings of the whole of  $X_{\mathbb{T}_R}$  by a pair of basic opens.*]