

MATHEMATICAL TRIPOS Part III

Monday 3 June 2002 9 to 12

PAPER 44

GALAXIES

*Attempt **THREE** questions*

*There are **four** questions in total*

The questions carry equal weight

**You may not start to read the questions
printed on the subsequent pages until
instructed to do so by the Invigilator.**

1 (a) Show that the gravitational potential energy of a spherical system can be written

$$W = -\frac{G}{2} \int_0^\infty \frac{M^2(r) dr}{r^2},$$

where $M(r)$ is the mass interior to radius r .

(b) Derive the virial theorem for a self-gravitating system of N identical points, each of mass m . Define the gravitational radius R_g in terms of W by the formula:

$$W \equiv -G (mN)^2 / R_g.$$

Find the rms velocity dispersion $\langle v_{eq}^2 \rangle$ for a system in equilibrium [the answer to be phrased in terms of (N, m, R_g)].

(c) Find the rms escape velocity $\langle v_{esc}^2 \rangle$ in the same units and also in terms of $\langle v_{eq}^2 \rangle$.

(d) If a system is not in equilibrium, but collapses from being at rest at infinity and is observed in free fall at a time when its gravitational energy is W (defined above), what is its velocity dispersion $\langle v_{coll}^2 \rangle$ in units of $\langle v_{eq}^2 \rangle$?

2 (a) A popular fit to the radial density distribution resultant from dark matter simulations of halos is the “NFW profile”:

$$\rho_D(r) \equiv \rho_1 \frac{r_1^3}{r(r+r_1)^2}.$$

Derive the mass distribution for this density law and find the circular velocity $v_{NFW,c}(r)$. Make a rough sketch of this function indicating the asymptotic ($r \rightarrow 0$, $r \rightarrow \infty$) dependencies. Estimate the maximum rotation velocity $v_{NFW,c,max}$ in units of (G, ρ_1, r_1) , and the radius at which this maximum occurs.

(b) Does the above derived functional form for $v_c(r)$ agree with observed galaxy rotation curves? If not, how is the difference to be understood?

(c) Assume that a galaxy initially has an NFW profile as above $\rho_D(r)$ and slowly accumulates additional mass until the resulting profile is flat with

$$v'_c = \text{const} = 2 v_{NFW,c,max}.$$

After the other matter $\rho_B(r)$ has settled down and the dark matter has reached a new equilibrium $\rho'_D(r)$ find expressions for the ratios:

$$\frac{\rho'_D(r)}{\rho_D(r)} = f(r); \quad \frac{\rho'_D(r)}{\rho_B(r)} = g(r);$$

in the domain $r \ll r_1$. [Hint: recall “adiabatic invariants”.]

3 We may study the vertical structure of a thin axisymmetric disk by neglecting all radial derivatives and adopting the form $f = f(E_z)$ for the distribution function, where $E_z \equiv \frac{1}{2}v_z^2 + \Phi(z)$. Show that if $f = \rho_0(2\pi\sigma_z^2)^{-1/2}\exp(-E_z/\sigma_z^2)$, the approximate form of Poisson's equation may be written:

$$2\frac{d^2\phi}{d\zeta^2} = e^{-\phi}, \quad \text{where } \phi \equiv \frac{\Phi}{\sigma_z^2}, \quad \zeta \equiv \frac{z}{z_0} \quad \text{and} \quad z_0 \equiv \frac{\sigma_z}{\sqrt{8\pi G\rho_0}}.$$

By solving this equation subject to the boundary conditions

$$\phi(0) = d\phi/d\zeta|_0 = 0,$$

show that the density ρ in the disk is given by

$$\rho(z) = \rho_0 \operatorname{sech}^2\left(\frac{1}{2}z/z_0\right).$$

4 There is a massive black hole (MBH) of mass M_H at the centre of a system of N stars each of mass m . In the vicinity of the black hole, gravitational collisions amongst the stars scatter them into a loss cone (in velocity space) whence, they plunge directly to the Schwarzschild radius of the MBH,

$$R_{sch} \equiv 2GM_H/c^2,$$

and are eaten whole.

- (a) Estimate the angular size of this loss cone in velocity space as a function of r .
- (b) Estimate the rate at which the MBH eats stars.
- (c) Estimate the rate at which the remaining cluster contracts or expands.