

MATHEMATICAL TRIPOS Part III

---

Tuesday 5 June 2001 1.30 to 4.30

---

PAPER 57

NUMERICAL SOLUTION OF DIFFERENTIAL EQUATIONS

Attempt **THREE** questions from Section A and attempt **ONE** question from Section B.

*Each question from Section A carries the weight of 18 points  
and each question from Section B carries the weight of 36 points.*

**You may not start to read the questions  
printed on the subsequent pages until  
instructed to do so by the Invigilator.**

## SECTION A

**1** Prove that any  $\nu$ -stage Runge-Kutta method of order  $2\nu$  is necessarily A-stable. [You may use, without proof, properties of Padé approximants to the exponential.]

**2** The parabolic equation

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \kappa u$$

is given for  $0 \leq x \leq 1, t \geq 0$ , together with an initial condition at  $t = 0$  and zero boundary conditions at  $x = 0$  and  $x = 1$ .

- (a) Prove (by separation of variables or otherwise) that the exact solution of the equation tends to zero as  $t \rightarrow \infty$  for every initial condition if and only if  $\kappa < \pi^2$ .
- (b) The equation is semidiscretized into

$$u'_m = \frac{1}{(\Delta x)^2}(u_{m-1} - 2u_m + u_{m+1}) + \kappa u_m, \quad m = 1, 2, \dots, M,$$

where  $(M+1)\Delta x = 1$ . Find the necessary and sufficient condition on  $\kappa$  that ensures  $\lim_{t \rightarrow \infty} u(t) = 0, m = 1, 2, \dots, M$ , for all possible initial conditions.

**3** Given  $\mathbf{y}' = \mathbf{f}(\mathbf{y})$ , we let  $\mathbf{g}(\mathbf{y}) := \mathbf{y}'' = (\partial \mathbf{f}(\mathbf{y}) / \partial \mathbf{y}) \mathbf{f}(\mathbf{y})$ . Determine the order of the two-step two-derivative method

$$\mathbf{y}_{n+1} - \frac{7}{15} h \mathbf{f}(\mathbf{y}_{n+1}) + \frac{1}{15} h^2 \mathbf{g}(\mathbf{y}_{n+1}) = \mathbf{y}_{n-1} + h \left( \frac{16}{15} \mathbf{f}(\mathbf{y}_n) + \frac{7}{15} \mathbf{f}(\mathbf{y}_{n-1}) \right) + \frac{1}{15} h^2 \mathbf{g}(\mathbf{y}_{n-1}).$$

**4** Prove that any positive-definite  $9 \times 9$  matrix with the sparsity structure

$$\begin{bmatrix} \times & 0 & \times & \times & 0 & 0 & \times & 0 & 0 \\ 0 & \times & 0 & 0 & 0 & 0 & 0 & \times & 0 \\ \times & 0 & \times & 0 & 0 & 0 & 0 & 0 & 0 \\ \times & 0 & 0 & \times & \times & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \times & \times & 0 & 0 & \times & 0 \\ 0 & 0 & 0 & 0 & 0 & \times & \times & 0 & 0 \\ \times & 0 & 0 & 0 & 0 & \times & \times & 0 & \times \\ 0 & \times & 0 & 0 & \times & 0 & 0 & \times & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \times & 0 & \times \end{bmatrix}$$

admits, after re-ordering, Gaussian elimination without fill-in. Describe the procedure that you have used in the proof and carefully state all relevant theorems.

**5** The semi-discretization

$$\begin{aligned}
 u'_m &= \frac{1}{\Delta x} \left( -\frac{3}{2}u_m + 2u_{m+1} - \frac{1}{2}u_{m+2} \right), \quad m = 1, 2, \dots, M-2, \Delta x = \frac{1}{M}, \\
 u'_{M-1} &= \frac{1}{\Delta x} \left( -\frac{3}{2}u_{M-1} + 2u_M - \frac{1}{2}u_1 \right), \\
 u'_M &= \frac{1}{\Delta x} \left( -\frac{3}{2}u_M + 2u_1 - \frac{1}{2}u_2 \right)
 \end{aligned}$$

approximates the solution of the advection equation  $u_t = u_x$ , given with initial conditions  $u(x, 0) = \varphi(x)$ ,  $0 \leq x \leq 1$ , and *periodic* boundary conditions  $u(0, t) = u(1, t)$ ,  $t \geq 0$ . Prove that the method is stable. [*Hint: You might formulate the problem in matrix form,  $\mathbf{u}' = \frac{1}{\Delta x} \mathbf{A} \mathbf{u}$ , say, and show that all eigenvectors of  $\mathbf{A}$  are of the form  $[1, \omega_\ell, \dots, \omega_\ell^{M-1}]^\top$  for some complex numbers  $\{\omega_\ell : \ell = 1, 2, \dots, M\}$ .]*

**6** Determine the range of the real parameter  $\alpha$  such that the multistep method

$$\begin{aligned}
 &\mathbf{y}_{n+3} - (1 + 2\alpha)\mathbf{y}_{n+2} + (1 + 2\alpha)\mathbf{y}_{n+1} - \mathbf{y}_n \\
 &= \frac{1}{6}h[(5 + \alpha)\mathbf{f}(\mathbf{y}_{n+3}) - (4 + 8\alpha)\mathbf{f}(\mathbf{y}_{n+2}) + (11 - 5\alpha)\mathbf{f}(\mathbf{y}_{n+1})]
 \end{aligned}$$

is convergent.

What is the order of the method for different values of  $\alpha$ ? For which values of  $\alpha$  is the method A-stable?

**SECTION B**

**7** Write an essay on the connection between convergence, order and stability in the numerical solution of ordinary differential equations.

**8** Describe in detail the fast Fourier transform and discuss its application to fast solution of the Poisson equation in a square.

**9** Describe the Engquist-Osher method for a single, one dimensional, hyperbolic nonlinear conservation law

$$u_t + \frac{\partial}{\partial x} f(u) = 0.$$

Prove that it is stable, provided that  $f$  is convex, differentiable and possesses a unique stagnation (sonic) point.