

## MATHEMATICAL TRIPOS Part III

Tuesday 5 June 2007 9.00 to 11.00

## PAPER 32

## ROUGH PATH THEORY AND APPLICATIONS

Attempt **THREE** questions.

There are **FOUR** questions in total.

The questions carry equal weight.

STATIONERY REQUIREMENTS

 $\begin{array}{c} \textbf{SPECIAL} \ \textbf{REQUIREMENTS} \\ \textit{None} \end{array}$ 

Cover sheet Treasury Tag Script paper

You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.



- 1 (i) Define  $(G^N(\mathbb{R}^d), \otimes, ^{-1}, e)$ , the free step-N nilpotent group over  $\mathbb{R}^d$ , and give the definition of the Carnot–Carathéodory d distance on  $G^N(\mathbb{R}^d)$ . What is a weak geometric p-rough path?
- (ii) How can the step-2 nilpotent group over  $\mathbb{R}^2$  be identified with the 3-dimensional Heisenberg group  $\mathbb{H}$ ?
- (iii) Since  $\mathbb{H} \cong \mathbb{R}^3$  we can equip  $\mathbb{H}$  with the *Euclidean* distance inherited from  $\mathbb{R}^3$ . Is a Lipschitz path in  $\mathbb{H}$  relative to this Euclidean distance automatically a Lipschitz path relative to the Carnot–Carathéodory distance on  $\mathbb{H}$ ?
- Let x be a Lipschitz continuous  $\mathbb{R}^d$ -valued path. Define  $S_N(x)_{s,t}$ , the step-Nsignature of the path segment  $x|_{[s,t]}$ , as an element in a suitable tensor algebra over  $\mathbb{R}^d$ . State and prove an algebraic relation between the step-N signature of the path segment  $x|_{[s,t]}$  and the path segment  $x|_{[t,u]}$  respectively. Show that the signature is invariant under reparametrisation of the path. More precisely, given  $\psi:[0,1]\to[0,1]$  strictly increasing and continuously differentiable, show that

$$S_N(x)_{0,1} = S_N(x \circ \psi)_{0,1}$$
.

- 3 Nested piecewise linear approximations to d-dimensional Brownian motion and their canonical area converge to Brownian motion and Lévy area in a rough path sense. Give a precise statement of this and sketch a proof with particular focus on martingale arguments.
- 4 Write an essay on the rough path proof of the Stroock-Varadhan support theorem. In particular, explain how the universal limit theorem is used.

## END OF PAPER