

MATHEMATICAL TRIPOS Part III

Monday 4 June 2007 1.30 to 4.30

PAPER 20

SPECTRAL GEOMETRY

*Attempt **THREE** questions.*

*There are **FIVE** questions in total.*

The questions carry equal weight.

STATIONERY REQUIREMENTS

*Cover sheet
Treasury Tag
Script paper*

SPECIAL REQUIREMENTS

None

<p>You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.</p>

1 Give three alternative definitions of the Laplacian acting on the functions on a Riemannian manifold and prove their equivalence.

2 Prove the existence of a 3-parameter family of pairs of bounded planar domains that are not isometric but are isospectral for the Laplacian acting on functions with Dirichlet boundary condition.

3 Define the heat kernel for a compact Riemannian manifold. If $p : N \rightarrow M$ is a finite normal locally isometric covering of Riemannian manifolds with covering transformation group U , obtain an expression for the heat kernel of M in terms of that of N .

Deduce a formula for the heat trace of M when U is a subgroup of a larger group T of isometries of N and hence deduce Sunada's Theorem.

4 Identify, with proof, a pair of Gassman equivalent subgroups U_1, U_2 of $PSL(3, 2)$. Explain briefly why they are not conjugate.

Assuming any result you require about hyperbolic tri-rectangles, show how to construct an uncountable family of pairs of isospectral Riemann surfaces of genus four. Explain briefly why they are isospectral and not isometric.

[You may use the fact that $PSL(3, 2)$ has generators A and D with commutator $C = [D, A]$ of order 7 such that the permutation actions of A and D on the cosets $U_i C^n$ are given by

Generator	cosets of U_1	cosets of U_2
A	(0)(1 2 5)(3 6 4)	(0)(1 4 3)(2 5 6)
D	(1)(0 3)(2 6 4 5)	(4)(2 5)(0 1 6 3)

where, in each case, the coset $U_i C^n$ is denoted by n .]

5 Define the Riemann surfaces referred to as X -pieces and Y -pieces. Identify the parameters that determine them up to isometry, including those involved when two Y -pieces form an X -piece.

Describe how a general closed Riemann surface of genus g may be formed from a certain number of Y -pieces and identify a set of parameters that suffice to determine the surface up to isometry.

Define Teichmüller space and state Wolpert's Theorem. Give three major ingredients of the proof of the theorem with a brief indication of the role that they play.

END OF PAPER