

PAPER 77

SYMMETRIC FUNCTIONS

*Attempt any **FIVE** questions. The questions carry equal weight.*

*The following notation is used throughout the paper. Let  $x = (x_1, x_2, \dots)$  be a set of indeterminates, and let  $n \in \mathbb{N}$ . The set of all homogeneous symmetric functions of degree  $n$  over  $\mathbb{Q}$  is denoted by  $\Lambda^n$ .*

*The vector space direct sum  $\Lambda = \Lambda^0 \oplus \Lambda^1 \oplus \dots$  is the  $\mathbb{Q}$ -algebra of symmetric functions.*

*If  $\lambda$  is a partition of  $n$ , we write  $|\lambda| = n$ . The length of  $\lambda$  will be denoted by  $l(\lambda)$ .*

*If  $S$  is any finite set,  $\mathfrak{S}_S$  will denote the symmetric group of all permutations of  $S$ .*

**You may not start to read the questions  
printed on the subsequent pages until  
instructed to do so by the Invigilator.**

**1** Describe the RSK algorithm. Show that this algorithm is a bijection between  $\mathbb{N}$ -matrices  $A = (a_{ij})_{i,j \geq 1}$  of finite support and ordered pairs  $(P, Q)$  of semistandard Young tableaux of the same shape. Show also that under this correspondence

$j$  occurs in  $P$  exactly  $\sum_i (a_{ij})$  times

and

$i$  occurs in  $Q$  exactly  $\sum_j (a_{ij})$  times.

[Standard facts about insertion paths can be quoted without proof, if clearly stated.]

Suppose that in the RSK algorithm  $A \xrightarrow{RSK} (P, Q)$ , the matrix  $A$  is symmetric (so  $P = Q$ ). Show by induction, or otherwise, that  $\text{tr}(A)$  is the number of columns of  $P$  of odd length. [Standard symmetry properties of the RSK algorithm may be assumed.]

**2** State and prove the Jacobi-Trudi identity, defining carefully all the terms. If  $\mu \subseteq \lambda$ , deduce an expression for the number of standard Young tableaux of shape  $\lambda/\mu$ . [If you use a result about determinants you should state it clearly.]

**3** State and prove ‘Burnside’s Lemma’. Given a finite set  $S$ , and for any subgroup  $G$  of  $\mathfrak{S}_S$ , show that  $Z_G = F_G$  where  $Z_G$  is the cycle indicator and  $F_G$  is the pattern inventory (both of which you should define).

The rank of a finite group  $G$  acting transitively on a set  $T$  is defined to be the number of orbits of  $G$  acting in the obvious way on  $T \times T$ , i.e.  $w.(s, t) = (w.s, w.t)$ . Thus  $G$  is doubly transitive if and only if  $\text{rank } G = 2$ . Let  $\chi$  be the character of the action of  $G$  on  $T$ . Show that  $\langle \chi, \chi \rangle = \text{rank } G$ .

**4** Define the partition set  $\text{Par}$  and explain how it can be endowed with a partial order  $\subseteq$  to become Young’s lattice. Define also the dominance order  $\leq$  and the reverse lexicographic order  $\stackrel{R}{\leq}$  on the set  $\text{Par}(n)$ . Define the elementary and complete homogeneous symmetric functions. Sketch a proof of the ‘fundamental theorem of symmetric functions’.

Write  $\Lambda = \mathbb{Q}[e_1, e_2, \dots]$  as a  $\mathbb{Q}$ -algebra generated by the elementary symmetric functions. If  $\{h_\lambda : \lambda \in \text{Par}\}$  are the complete homogeneous symmetric functions, prove the existence of an involutory automorphism  $\omega : \Lambda \rightarrow \Lambda$  such that  $\omega(e_\lambda) = h_\lambda$  for all partitions  $\lambda$ .

Let  $\lambda$  be a partition of  $n$  of length  $l$ . Define the forgotten symmetric function  $f_\lambda$  by  $f_\lambda = \epsilon_\lambda \omega(m_\lambda)$  where  $\epsilon_\lambda = (-1)^{n-l}$  and  $m_\lambda$  is the monomial symmetric function. Let  $f_\lambda = \sum_\mu a_{\lambda\mu} m_\mu$ . Show that  $a_{\lambda\mu}$  is equal to the number of distinct permutations  $(\alpha_1, \alpha_2, \dots, \alpha_l)$  of the sequence  $(\lambda_1, \lambda_2, \dots, \lambda_l)$  such that

$$\{\alpha_1 + \alpha_2 + \dots + \alpha_i : 1 \leq i \leq l\} \supseteq \{\mu_1 + \mu_2 + \dots + \mu_j : 1 \leq j \leq l(\mu)\}.$$

**5** Let  $\lambda$  and  $\mu$  be partitions such that  $\mu \subseteq \lambda$ . Define the semistandard tableau of (skew) shape  $\lambda/\mu$  and show how we may regard it as a Young diagram of shape  $\lambda/\mu$ . If  $\lambda/\mu$  is a skew shape define the skew Schur function  $s_{\lambda/\mu}$  and the Schur function  $s_\lambda$  of shape  $\lambda$ . Show that for any skew shape  $\lambda/\mu$ ,  $s_{\lambda/\mu}$  is indeed a symmetric function.

Let  $\lambda, \mu$  be partitions with  $|\lambda| = |\mu|$ . Suppose that the  $(\lambda, \mu)$ th Kostka number  $K_{\lambda\mu}$  is non-zero. Prove that  $\lambda \geq \mu$  (in the dominance order) and also that  $K_{\lambda\lambda} = 1$ . Deduce that the Schur functions  $s_\lambda$  form a  $\mathbb{Q}$ -basis for  $\wedge$  ( $\lambda \in \text{Par}$ ).

**6** Let  $\lambda \in \text{Par}$  and suppose  $l(\lambda) \leq n$ . Let  $\delta = (n-1, n-2, \dots, 0)$ . Quoting any result that you need, prove the bialternant relation

$$a_{\lambda+\delta}/a_\delta = s_\lambda(x_1, \dots, x_n).$$

Let  $p$  be a prime, and let  $A_p$  denote the matrix  $[\zeta^{jk}]_{j,k=0}^{p-1}$ , where  $\zeta = e^{2\pi i/p}$ . Show that every minor of  $A_p$  is non-zero (i.e. that every square submatrix  $B$  of  $A_p$  is nonsingular).

Show finally that

$$h_r(x_1, \dots, x_n) = \sum_{k=1}^n x_k^{n-1+r} \prod_{i \neq k} (x_k - x_i)^{-1}$$

[Hint: Consider a particular value for  $\lambda$  in the bialternant.]

**7** State and prove the Cauchy identity (the RSK algorithm may be assumed). Use the identity to demonstrate that the Schur functions form an orthonormal basis for  $\wedge$ . State the dual Cauchy identity.

Let  $F(t) = \sum_{j \geq 0} f_j t^j$  be a formal power series, where  $f_0 = 1$ . Expand the product  $F(t_1)F(t_2) \dots$  as a linear combination of Schur functions  $s_\lambda(t_1, t_2, \dots)$ . The coefficient of  $s_\lambda(t_1, t_2, \dots)$  will be denoted by  $s_\lambda^F$ . Equivalently, if  $R$  is a commutative ring containing  $f_1, f_2, \dots$  and  $\varphi: \wedge \rightarrow R$  is the homomorphism defined by  $\varphi(h_j) = f_j$ , then  $s_\lambda^F = \varphi(s_\lambda)$ . We extend the definition of  $s_\lambda^F$  by defining  $u^F = \varphi(u)$  for any  $u \in \wedge_R$ .

(a) Show that if  $F(t) = \prod_{i \geq 1} (1 - x_i t)^{-1}$ , then  $s_\lambda^F = s_\lambda(x)$ .

(b) What if  $F(t) = \prod_{i \geq 1} (1 + x_i t)$ ?

**8** What does it mean to say that a skew shape  $\lambda/\mu$  is connected? Define a border strip,  $B$  and define the height,  $\text{ht}(B)$  of  $B$ .

Let  $p_\lambda$  be the power sum symmetric function. Let  $\alpha$  be a weak composition of  $n$ . Define a border-strip tableau of shape  $\lambda/\mu$  and type  $\alpha$ . Show that

$$s_\mu p_\alpha = \sum_{\lambda} \chi^{\lambda/\mu}(\alpha) s_\lambda,$$

where  $\chi^{\lambda/\mu}(\alpha) = \sum_T (-1)^{\text{ht}(T)}$ , summed over all border-strip tableaux of shape  $\lambda/\mu$  and type  $\alpha$ . Restricting to  $n$  variables where  $n \geq \ell(\lambda)$ , deduce that  $\chi^\lambda(\alpha) = [x^{\lambda+\delta}] p_\alpha a_\delta$  where  $\delta = (n-1, n-2, \dots, 0)$  and  $a_\delta$  is the Vandermonde determinant. Deduce also the Murnaghan-Nakayama rule. Show also that  $s_\delta$  is a polynomial in the odd power sums  $p_1, p_3, \dots$ , where  $\delta$  is the “staircase shape”  $\delta = (m-1, m-2, \dots, 1)$ .

Finally, let  $0 \leq s \leq n-1$  and  $|\lambda| = n$ . Show that if  $w \in \mathfrak{S}_n$  is an  $n$ -cycle, then

$$\chi^\lambda(w) = \begin{cases} (-1)^s & \text{if } \lambda = (n-s, 1^s) \\ 0 & \text{otherwise} \end{cases} .$$