

MATHEMATICAL TRIPOS Part III

Monday 6 June, 2005 1.30 to 4.30

PAPER 17

TOPICS IN ALGEBRAIC GEOMETRY

*Attempt **ALL** questions.*

*Question 1 carries 10 marks
and the others 30 marks each.*

STATIONERY REQUIREMENTS

*Cover sheet
Treasury Tag
Script paper*

SPECIAL REQUIREMENTS

None

<p>You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.</p>

1 Define the formal completion X/Y of a scheme along a closed subscheme Y .

Suppose $X = \mathbb{A}_{k[q]}^1 = \text{Spec } k[q, t]$ and $Z = \mathbb{G}_{m, k[q]} = X_1 - (t = 0)$. Are the formal schemes $Z_{/(q=0)}$ and $X_{/(q=0)} - (t = 0)$ isomorphic? Explain your answer.

2 Show how torus embeddings may be used to construct a morphism $X \rightarrow \text{Spec } \mathbb{Z}[q]$ such that the fibre Y over $(q = 0)$ is an infinite chain of copies of $\mathbb{P}_{\mathbb{Z}}^1$ and the generic fibre is a copy of $\mathbb{G}_{m, \mathbb{Z}[q, q^{-1}]}$.

3 Take X, Y to be the schemes constructed in question 2. Explain how to construct a formal scheme that is the quotient of X/Y by some non-trivial action of an infinite cyclic group.

4 Suppose that M is a free \mathbb{Z} -module of finite rank. Show that the functor $F : \text{Rings} \rightarrow \text{Sets}$ such that $F(R)$ is the set of locally free rank r quotient modules of $M \otimes_{\mathbb{Z}} R$ is representable.

END OF PAPER