

UNIVERSITY OF LONDON

[II(3)E 2001]

B.ENG. AND M.ENG. EXAMINATIONS 2001

For Internal Students of the Imperial College of Science, Technology and Medicine

This paper is also taken for the relevant examination for the Associateship.

PART II : MATHEMATICS 3 (ELECTRICAL ENGINEERING)

Wednesday 6th June 2001 2.00 - 5.00 pm

Answer EIGHT questions.

[Before starting, please make sure that the paper is complete; there should be 7 pages, with a total of 12 questions. Ask the invigilator for a replacement if your copy is faulty.]

Copyright of the University of London 2001

1. Consider the mapping

$$w = \frac{1}{z - i}$$

from the z -plane to the w -plane, where $z = x + iy$ and $w = u + iv$.

(i) Show that, in the z -plane, the family of circles centred at $(0, 1)$ with radius a

$$x^2 + (y - 1)^2 = a^2$$

maps to another family of circles in the w -plane. What is the radius of this family and where is its centre?

(ii) What is the image in the w -plane of the x -axis ($y = 0$) in the z -plane? Show that the curve that represents this image passes through the origin in the w -plane.

(iii) Show that the family of straight lines $y = cx$ in the z -plane with $c = \text{constant}$ have the image in the w -plane represented by

$$\left(u - \frac{c}{2}\right)^2 + \left(v - \frac{1}{2}\right)^2 = \frac{1}{4}(1 + c^2).$$

2. By choosing a suitable contour C in the upper half of the complex plane, use the contour integral

$$\oint_C \frac{e^{iz} dz}{(z^2 + 4)(z^2 + 1)}$$

to show that

$$\int_{-\infty}^{\infty} \frac{\cos x dx}{(x^2 + 4)(x^2 + 1)} = \frac{\pi}{6} \left(\frac{2e - 1}{e^2}\right).$$

PLEASE TURN OVER

3. (i) Show that if C is a circle of arbitrary radius r centred at the origin, then the value of the complex integral

$$\oint_C \frac{dz}{z}$$

is independent of r . What is this value?

- (ii) Use the Residue Theorem to show that

$$\oint_C \frac{z dz}{(z-1)^2(z-i)} = 0,$$

where the contour C is the circle of radius 2 centred at the origin. What is the answer when C is changed to be the rectangle with vertices at $\pm\frac{1}{2} + 2i$ and $\pm\frac{1}{2} - 2i$?

Recall that the residue of a complex function $f(z)$ at a pole $z = a$ of multiplicity m is given by the expression

$$\lim_{z \rightarrow a} \frac{1}{(m-1)!} \left[\frac{d^{m-1}}{dz^{m-1}} \{(z-a)^m f(z)\} \right].$$

4. The Fourier convolution of the functions $f(t)$ and $g(t)$ is defined by

$$f * g = \int_{-\infty}^{\infty} f(u)g^*(t-u) du$$

where g^* is the complex conjugate of g . If $\bar{f}(\omega)$ and $\bar{g}(\omega)$ are the Fourier transforms of $f(t)$ and $g(t)$ respectively, prove the Fourier convolution theorem

$$\int_{-\infty}^{\infty} e^{-i\omega t} (f * g) dt = \bar{f}(\omega) \bar{g}(\omega).$$

For a function $f(t)$, if $\gamma(t)$ is defined by

$$\gamma(t) = \frac{f * f}{\int_{-\infty}^{\infty} |f(t)|^2 dt}$$

show that

$$\int_{-\infty}^{\infty} \bar{\gamma}(\omega) d\omega = 2\pi.$$

5. (i) A second order ordinary differential equation takes the form

$$\frac{d^2x}{dt^2} + \omega^2x = f(t),$$

where $f(t)$ is an arbitrary piecewise smooth function. It has initial conditions

$$x = \frac{dx}{dt} = 0 \text{ when } t = 0.$$

Use the Laplace convolution theorem to show that

$$x(t) = \frac{1}{\omega} \int_0^t \sin(\omega u) f(t-u) du.$$

- (ii) A third order ordinary differential equation takes the form

$$\frac{d^3x}{dt^3} + 3\frac{d^2x}{dt^2} + 3\frac{dx}{dt} + x = f(t)$$

where $f(t)$ is an arbitrary piecewise smooth function. $x(t)$ and its first two derivatives satisfy the conditions

$$x = \frac{dx}{dt} = \frac{d^2x}{dt^2} = 0 \text{ when } t = 0.$$

Use the shift and convolution theorems to show that

$$x(t) = \frac{1}{2} \int_0^t e^{-u} u^2 f(t-u) du.$$

6. Given that

$$\int_{-\infty}^{\infty} \frac{e^{ix}}{x} dx = i\pi,$$

show that

$$\int_{-\infty}^{\infty} \frac{e^{ipx}}{x} dx = \begin{cases} +i\pi, & p > 0 \\ -i\pi, & p < 0, \end{cases}$$

where p is an arbitrary real number. Hence show that the Fourier transform $\bar{f}(\omega)$ of the function

$$f(t) = \frac{\sin t/2}{t/2}$$

is given by

$$\bar{f}(\omega) = \begin{cases} 2\pi, & -\frac{1}{2} < \omega < \frac{1}{2}, \\ 0, & \omega < -\frac{1}{2}, \quad \omega > \frac{1}{2}. \end{cases}$$

PLEASE TURN OVER

7. (i) The double integral I_1 is given by

$$I_1 = \iint_{R_1} (x+y)^2 \cos(x^2 - y^2) \, dx dy,$$

where R_1 is the finite region in the x - y plane enclosed by the lines $x = 0$, $y = 0$ and $y = 1 - x$.

Show that, by using the transformation,

$$u = x - y, \quad v = x + y,$$

the integral can be written as

$$I_1 = \frac{1}{2} \int_0^1 v^2 \left(\int_{-v}^v \cos(uv) \, du \right) dv.$$

Hence evaluate I_1 .

- (ii) Use the same transformation to evaluate

$$I_2 = \iint_{R_2} (x^2 + y^2) \, dx dy,$$

where R_2 is the interior of the square bounded by $y = \pm x$, $y = \pm (x-1)$.

8. A vector field \mathbf{F} is defined as

$$\mathbf{F} = 2xye^z \mathbf{i} + x^2 e^z \mathbf{j} + (x^2 y e^z + z^2 + 3z) \mathbf{k}.$$

- (i) Find $\text{div } \mathbf{F}$ and $\text{curl } \mathbf{F}$.
 (ii) Find a function $\phi(x, y, z)$ such that $\mathbf{F} = \nabla \phi$.
 (iii) Evaluate

$$\frac{\partial^2}{\partial z^2} (x \mathbf{F} \cdot \mathbf{i} - 2\phi).$$

9. (i) The vector field \mathbf{F} is defined by

$$\mathbf{F} = (y^2 \cos x) \mathbf{i} + (\alpha y \sin x) \mathbf{j},$$

where α is a constant. Find the value of α such that $\text{curl } \mathbf{F} = \mathbf{0}$.

- (ii) Consider the integral

$$I = \int_C (y^2 \cos x \, dx + \beta y \sin x \, dy), \quad (\beta \text{ constant}),$$

where C is a curve joining the points $(0, 0)$ and $(\pi/2, 1)$.

Evaluate I in the following cases:

- (a) C is the line $y = (2/\pi)x$;
 (b) C is the curve $y = \sin x$.

Show that the answers to (a) and (b) are equal for one particular value of β and find that value.

Explain why the value of α found in part (i) is the same as this value of β .

10. P and Q are continuous functions of x and y with continuous first partial derivatives in a simply connected region R with a piecewise smooth boundary C . Green's Theorem in a plane says that

$$\oint_C (P \, dx + Q \, dy) = \iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \, dx \, dy.$$

Find a two-dimensional vector \mathbf{u} , defined in terms of P and Q , to show that Green's Theorem can be re-expressed as the two-dimensional version of the Divergence Theorem

$$\oint_C \mathbf{u} \cdot \mathbf{n} \, ds = \iint_R \text{div } \mathbf{u} \, dx \, dy$$

where \mathbf{n} is the unit normal to the curve C .

If \mathbf{u} is given by $\mathbf{u} = x^2 \mathbf{i} + y^2 \mathbf{j}$ and R is the first quadrant of the circle of unit radius, evaluate the right hand side of the Divergence Theorem to show that

$$\iint_R \text{div } \mathbf{u} \, dx \, dy = 4/3.$$

PLEASE TURN OVER

11. Let A_1, \dots, A_k form a partition of a sample space and B be some event. Use the definition of conditional probability and the theorem of total probabilities to derive Bayes's formula for $P(A_i|B)$.

It is estimated that 5% of optical disks produced by a manufacturer are faulty. A disk may be subjected to an initial diagnostic test. If there is a fault, the test gives a diagnosis of 'faulty' with probability 0.8; if there is no fault the test gives a diagnosis of 'OK' with probability 0.95. If the test gives a diagnosis of 'faulty', the disk is rejected. A disk is chosen at random and tested. What is the probability that

- (i) the test gives a diagnosis of 'OK'?
- (ii) a disk is faulty which has been given a diagnosis of 'OK'?

If the initial test gives the diagnosis 'OK', a further independent test is performed; this test has exactly the same properties as the initial test, except that if there is a fault, the test gives a diagnosis of 'faulty' 99% of the time. If this test gives a diagnosis of 'OK' the disk is accepted for use, otherwise it is rejected.

- (iii) Determine the probability that a faulty disk is accepted for use.

12. Let X and Y be two random variables. The coefficient of correlation between X and Y is given by

$$\rho_{X,Y} = \frac{\text{cov}\{X, Y\}}{[\text{var}\{X\} \text{var}\{Y\}]^{1/2}} = \frac{E\{XY\} - E\{X\}E\{Y\}}{[\text{var}\{X\} \text{var}\{Y\}]^{1/2}}.$$

- (i) What does it measure? How should values of $\rho_{X,Y}$ of -1 , 0 and 1 be interpreted?
- (ii) If X and Y are independent, what is $\rho_{X,Y}$?

Let X and Y have the joint probability density function given by

$$f_{X,Y}(x, y) = \begin{cases} x^{-1}, & 0 \leq y \leq x \leq 1, \\ 0, & \text{otherwise.} \end{cases}$$

- (iii) Calculate $E\{XY\}$ and $E\{X\}$ and $E\{Y\}$, and hence find the value of $\text{cov}\{X, Y\}$.
- (iv) Are X and Y independent?

END OF PAPER

MATHS 3
2001

MATHEMATICS FOR ENGINEERING STUDENTS
EXAMINATION QUESTION / SOLUTION
SESSION : 2000 - 2001

PAPER
3

QUESTION

Please write on this side only, legibly and neatly, between the margins

SOLUTION
14

$$W = \frac{1}{z-i} = \frac{1}{x+i(y-1)} = \frac{x-i(y-1)}{x^2+(y-1)^2} = u+iv$$

$$u = \frac{x}{x^2+(y-1)^2} ; v = \frac{-(y-1)}{x^2+(y-1)^2}$$

$$u^2 + v^2 = \frac{1}{x^2+(y-1)^2} \quad (*)$$


a) The family of circles $x^2+(y-1)^2 = a^2$ map to
 $u^2+v^2 = \frac{1}{4}a^2$ - a family centred at (0,0)
radius a^{-1} .

b) $y=0 \Rightarrow u = \frac{x}{x^2+1}, v = \frac{1}{x^2+1}$

From (*) $\therefore u^2 + v^2 = \frac{1}{x^2+1} = v$

$$\therefore u^2 + (v - \frac{1}{2})^2 = (\frac{1}{2})^2$$

A circle centred $(0, \frac{1}{2})$ radius $\frac{1}{2}$



c) $y = cx \quad u = \frac{x}{x^2+(cx-1)^2}, v = \frac{1-cx}{x^2+(cx-1)^2}$

$$u^2 + v^2 = \frac{1}{x^2+(cx-1)^2} \quad v = \frac{1}{x^2+(cx-1)^2} - cu$$

$$\therefore u^2 + v^2 = v + cu$$

$$\therefore (u - \frac{c}{2})^2 + (v - \frac{1}{2})^2 = \frac{1}{4}(1+c^2)$$

Family of circles centred at $(\frac{c}{2}, \frac{1}{2})$ radius $\frac{1}{2}\sqrt{1+c^2}$.

Setter : J.D. GIBBON

Setter's signature : J.D. Gibbon

Checker : NEARZAM

Checker's signature : [Signature]

3

3

4

5

MATHEMATICS FOR ENGINEERING STUDENTS
 EXAMINATION QUESTION / SOLUTION
 SESSION : 2000 - 2001

PAPER

3

QUESTION

SOLUTION


15

Please write on this side only, legibly and neatly, between the margins

Two simple poles in C at $z=i, 2i$

Cauchy's Thm \Rightarrow

$$\oint_C \frac{e^{iz} dz}{(z^2+4)(z^2+1)} = 2\pi i \{ \text{Sum of Residues} \}$$



Residue at $z=i = -\frac{i}{6} e^{-1}$
 " " $z=2i = \frac{e^{-2}}{(1-4)4i} = \frac{i e^{-2}}{12}$

$$\therefore \oint_C = 2\pi i^2 \left(\frac{e^{-2}}{12} - \frac{e^{-1}}{6} \right) = \frac{\pi}{6} \left(\frac{2e-1}{e^2} \right)$$

Now $\oint_C \frac{e^{iz} dz}{(z^2+4)(z^2+1)} = \int_{-R}^R \frac{e^{ix} dx}{(x^2+4)(x^2+1)} + \int_{H_R} \frac{e^{iz} dz}{(z^2+4)(z^2+1)}$

Now take the limit $R \rightarrow \infty$

$$\lim_{R \rightarrow \infty} \oint_C = \int_{-\infty}^{\infty} \frac{e^{ix} dx}{(x^2+4)(x^2+1)} + \lim_{R \rightarrow \infty} \int_{H_R}$$

Now by Jordan's Lemma $\lim_{R \rightarrow \infty} \int_{H_R} = 0$ provided

- (i) Only singularities in upper $\frac{1}{2}$ -plane are poles \checkmark
- (ii) $f(z) \rightarrow 0$ as $R \rightarrow \infty$ \checkmark $\int_{H_R} e^{imz} f(z) dz = \int_{H_R}$
- (iii) $m > 0$ \checkmark

Moreover

$$\begin{aligned} \int_{-\infty}^{\infty} &= \int_{-\infty}^{\infty} \frac{(\cos x + i \sin x) dx}{(x^2+4)(x^2+1)} \\ &= \int_{-\infty}^{\infty} \frac{\cos x dx}{(x^2+4)(x^2+1)} + i \cdot 0 \text{ by symmetry} \end{aligned}$$

$$\therefore \int_{-\infty}^{\infty} \frac{\cos x dx}{(x^2+4)(x^2+1)} = \frac{\pi}{6} \left(\frac{2e-1}{e^2} \right)$$

Please write on this side only, legibly and neatly, between the margins


$$(ii) \oint_C F(z) dz = 2\pi i \times \{ \text{sum of Residues of } F(z) \text{ in } C \}$$

Residues at $z=i$ (simple)
 $z=1$ (double)

calculated: Res. at $z=i$ is $\frac{i}{(i-1)^2} = \frac{-1}{2}$


" " $z=1$ is $\left[\frac{d}{dz} \left(\frac{z}{z-i} \right) \right]_{z=1}$

$$= \left[\frac{(z-i) - z}{(z-i)^2} \right]_{z=1} = \frac{-1}{-2i} = \frac{1}{2}$$



$$\text{Sum of Residues} = -\frac{1}{2} + \frac{1}{2} = 0$$


If the contour is changed to the box
 then $z=1$ is excluded and we
 have $2\pi i \times (-\frac{1}{2}) = -\pi i$



$$(i) \oint_C \frac{dz}{z} = \int_0^{2\pi} \frac{i r e^{i\theta} d\theta}{r e^{i\theta}}$$

$$= i \int_0^{2\pi} d\theta$$

$$= 2\pi i \quad (\text{independent of } r)$$



Alternatively, by the Residue Theorem $F(z) = \frac{1}{z}$
 has one simple pole at $z=0$ in C

$$\therefore \oint_C \frac{dz}{z} = 2\pi i \times (\text{Res at } z=0)$$

$$\text{Res. of } \frac{1}{z} \text{ at } z=0 = 1$$

$$\therefore \oint_C \frac{dz}{z} = 2\pi i \quad \text{regardless of size of circle.}$$

MATHEMATICS FOR ENGINEERING STUDENTS
EXAMINATION QUESTION / SOLUTION
SESSION : 2000 - 2001

E4

PAPER

3

QUESTION

17

SOLUTION

Please write on this side only, legibly and neatly, between the margins

$$f * g = \int_{-\infty}^{\infty} f(t')g(t-t')dt' \quad (t' \equiv u \text{ in question})$$

$$\text{F.T. } (f * g) = \int_{-\infty}^{\infty} e^{-i\omega t} (f * g) dt$$

$$= \int_{-\infty}^{\infty} e^{-i\omega t} \left(\int_{-\infty}^{\infty} f(t')g(t-t')dt' \right) dt$$

Let $\tau = t - t'$: exchanging the order of integration causes no problem as the domain is doubly infinite

$$\begin{aligned} \therefore \text{FT } (f * g) &= \int_{-\infty}^{\infty} f(t') \left(\int_{-\infty}^{\infty} e^{-i\omega t} g(t-t') dt \right) dt' \\ &= \int_{-\infty}^{\infty} f(t') \left(e^{-i\omega t'} \int_{-\infty}^{\infty} e^{-i\omega \tau} g(\tau) d\tau \right) dt' \\ &= \left(\int_{-\infty}^{\infty} f(t') e^{-i\omega t'} dt' \right) \left(\int_{-\infty}^{\infty} e^{-i\omega \tau} g(\tau) d\tau \right) \\ &= \bar{f}(\omega) \quad \bar{g}(\omega) \end{aligned}$$

10

$$\text{Now } \gamma(t) = \int_{-\infty}^{\infty} f(t')f(t-t')dt' / \int_{-\infty}^{\infty} |f(t)|^2 dt$$

Numerator is a number so we have

$$\bar{\gamma}(\omega) = \text{F.T. } (\gamma(t)) = \left(\int_{-\infty}^{\infty} |f|^2 dt \right)^{-1} [\bar{f}(\omega)\bar{f}(\omega)]$$

from the Convolution Thm.

$$\text{Now } \int_{-\infty}^{\infty} |f(t)|^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} |\bar{f}(\omega)|^2 d\omega \quad (\text{Parseval})$$

$$\therefore \int_{-\infty}^{\infty} \bar{\gamma}(\omega) d\omega = 2\pi$$

5

2

2

1

Setter : J.D. GIBBON

Setter's signature : J.D. Gibbon

Checker : NERBEM

Checker's signature : NERBEM

MATHEMATICS FOR ENGINEERING STUDENTS
 EXAMINATION QUESTION / SOLUTION
 SESSION : 2000 - 2001

PAPER

3

QUESTION

18

SOLUTION

18

Please write on this side only, legibly and neatly, between the margins

a) $\ddot{x} + \omega^2 x = f(t)$

($t' \equiv u$
in question)

Using the Math. formulae $\mathcal{L}(\ddot{x}) = s^2 \bar{x}(s) - s x(0) - \dot{x}(0)$
 $= s^2 \bar{x}(s)$ in this case

$\therefore (s^2 + \omega^2) \bar{x}(s) = \bar{f}(s)$

$\therefore \bar{x}(s) = \bar{f}(s) \bar{g}(s)$ where $\bar{g}(s) = \frac{1}{s^2 + \omega^2}$

Hence $g(t) = \frac{1}{\omega} \sin \omega t$ ($s > 0$)

and $x(t) = \mathcal{L}^{-1}[\bar{f}(s) \bar{g}(s)] = \int_0^t g(t-t') f(t') dt'$
 $= \frac{1}{\omega} \int_0^t \sin(\omega t') f(t-t') dt'$ Convolution
Thm

b) $\ddot{x} + 3\dot{x} + 3x = f(t)$

Now $\int_0^\infty e^{-st} \ddot{x} dt = \int_0^\infty e^{-st} d(\dot{x})$
 $= [\dot{x} e^{-st}]_0^\infty + s \int_0^\infty \dot{x} e^{-st} dt$

For $s > 0$, we have, with I.C.s $\dot{x}(0) = \ddot{x}(0) = x(0)$

$\mathcal{L}(\ddot{x}) = s \mathcal{L}(\dot{x}) = s^3 \bar{x}(s)$

$\therefore (s^3 + 3s^2 + 3s + 1) \bar{x}(s) = \bar{f}(s)$

or $(s+1)^3 \bar{x}(s) = \bar{f}(s) \Rightarrow \bar{x}(s) = \frac{\bar{f}(s)}{(s+1)^3}$

Now $\mathcal{L}(s^{-3}) = \frac{1}{2} t^2$,

$(s+1)^{-3}$

plus the shift theorem, gives

$g(t) = \frac{1}{2} t^2 e^{-t}$

$\therefore x(t) = \frac{1}{2} \int_0^t e^{-t'} t'^2 f(t-t') dt'$

2

2

7 3

2

3

8

3

Setter : J. D. GIBSON

Setter's signature : J. D. Gibson

Checker : NERRANT

Checker's signature : N. Robert

Please write on this side only, legibly and neatly, between the margins

In $\int_{-\infty}^{\infty} \frac{e^{ix}}{x} dx$, put $x=pt$ $dx=pt$ $x=\infty \Rightarrow$


$$\int_{-\infty}^{\infty} \frac{e^{ix}}{x} dx = \int_{-\infty/p}^{\infty/p} \frac{e^{i/p t}}{t} dt = \begin{cases} i\pi & p > 0 \\ -i\pi & p < 0 \end{cases}$$

F.T. $\left(\frac{\sin t/h}{t/h}\right) = \int_{-\infty}^{\infty} e^{-i\omega t} \left(\frac{e^{it/h} - e^{-it/h}}{2it/h}\right) dt$

$$= -i \int_{-\infty}^{\infty} \frac{e^{i(\frac{1}{2}-\omega)t}}{t} dt + i \int_{-\infty}^{\infty} \frac{e^{i(-\omega-\frac{1}{2})t}}{t} dt$$

$$= \begin{cases} -i(i\pi) + i(-i\pi) & \omega < \frac{1}{2} \\ -i(i\pi) + i(-i\pi) & -\frac{1}{2} < \omega \\ -i(-i\pi) + i(-i\pi) & \omega > \frac{1}{2} \\ -i(i\pi) + i(i\pi) & \omega < -\frac{1}{2} \end{cases}$$

$$= \begin{cases} 2\pi & -\frac{1}{2} < \omega < \frac{1}{2} \\ 0 & \omega > \frac{1}{2}, \omega < -\frac{1}{2} \end{cases}$$



3

6


6

Please write on this side only, legibly and neatly, between the margins

(i) Jacobian is given by $J = \begin{vmatrix} u_x & v_x \\ u_y & v_y \end{vmatrix}^{-1}$

$$= \begin{vmatrix} 1 & 1 \\ -1 & 1 \end{vmatrix}^{-1} = (1 - (-1))^{-1} = \underline{\underline{\frac{1}{2}}}$$

We have $u = x - y, v = x + y \Rightarrow x = \frac{1}{2}(u + v); y = \frac{1}{2}(v - u)$
 Thus $x = 0 \Rightarrow u = -v$ and $y = 1 - x \Rightarrow v = 1$
 $y = 0 \Rightarrow u = v$



So: $I_1 = \int_{v=0}^{v=1} \int_{u=-v}^{u=v} v^2 \cos(uv) du dv \cdot \left(\frac{1}{2}\right)$ ← from J

$$= \int_{v=0}^{v=1} \frac{1}{2} v^2 \left[\frac{\sin(uv)}{v} \right]_{u=-v}^{u=v} dv$$

$$= \int_0^1 v \sin(v^2) dv \quad \text{Subst } t = v^2.$$

$$= \frac{1}{2} \int_0^1 \sin t dt = \underline{\underline{\frac{1}{2}(1 - \cos(1))}}$$

(ii) As above, $J = \frac{1}{2}$.

$$x^2 + y^2 = \left(\frac{1}{2}(u+v)\right)^2 + \left(\frac{1}{2}(u-v)\right)^2 = \frac{1}{2}(u^2 + v^2)$$

$$y = \pm x \Rightarrow \underline{u=0}, \underline{v=0}$$

$$y = x - 1 \Rightarrow \underline{u=1}, \quad y = 1 - x \Rightarrow \underline{v=1}$$

and $I_2 = \int_{v=0}^{v=1} \int_{u=0}^{u=1} \frac{1}{2}(u^2 + v^2) \cdot \frac{1}{2} du dv$

$$= \frac{1}{4} \int_0^1 \left[\frac{u^3}{3} + v^2 u \right]_{u=0}^{u=1} dv$$

$$= \frac{1}{4} \int_0^1 \left(\frac{1}{3} + v^2 \right) dv = \frac{1}{4} \left[\frac{1}{3}v + \frac{v^3}{3} \right]_0^1 = \underline{\underline{\frac{1}{6}}}$$

Total
15

Setter : WALTON

Setter's signature : Enbrew Walton

Checker : JACOBS

Checker's signature : R.L. Jacobs

MATHEMATICS FOR ENGINEERING STUDENTS
 EXAMINATION QUESTION / SOLUTION
 SESSION : 2000 - 2001

PAPER
 BEN 9 / Maths
 PART 2

QUESTION

SOLUTION
 22

Please write on this side only, legibly and neatly, between the margins

$$\text{div } \underline{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z} \quad \underline{F} = (F_1, F_2, F_3)$$

(i) $\therefore \text{div } \underline{F} = 2ye^z + x^2ye^z + 2z + 3$ (3)

* $\text{Curl } \underline{F} = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2xye^z & x^2e^z & x^2ye^z + z^2 + 3z \end{vmatrix}$

$$= \underline{i} (x^2e^z - x^2e^z) - \underline{j} (2xye^z - 2xye^z) + \underline{k} (2xe^z - 2xe^z)$$

(3)

$$= 0$$

(ii) $\frac{\partial \phi}{\partial x} = 2xye^z \Rightarrow \phi = x^2ye^z + g_1(y, z)$ (2)

$\frac{\partial \phi}{\partial y} = x^2e^z \Rightarrow \phi = x^2ye^z + g_2(y, z)$

$\frac{\partial \phi}{\partial z} = x^2ye^z + z^2 + 3z \Rightarrow \phi = x^2ye^z + \frac{z^3}{3} + \frac{3z^2}{2} + g_3(x, y)$

Hence $\phi = x^2ye^z + \frac{z^3}{3} + \frac{3z^2}{2} + \text{const.}$ (4)

(iii) $\underline{F} \cdot \underline{i} = 2xye^z$
 $x \underline{F} \cdot \underline{i} = 2x^2ye^z$

$\therefore x \underline{F} \cdot \underline{i} - 2\phi = -\frac{2z^3}{3} - 3z^2 + \text{const.}$ (3)

$$\frac{\partial^2}{\partial z^2} (x \underline{F} \cdot \underline{i} - 2\phi) = -4z - 6$$

15

Setter : ATKINSON
 Checker: J. D. GIBBON

Setter's signature: c. atkinson
 Checker's signature: J.D. Gibbon

Please write on this side only, legibly and neatly, between the margins

$$(i) \underline{F} = \alpha y \sin x \hat{j} + y^2 \cos x \hat{i}$$

$$\Rightarrow \text{Curl } \underline{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & 0 \\ y^2 \cos x & \alpha y \sin x & 0 \end{vmatrix} = \hat{k} \begin{pmatrix} \alpha y \cos x \\ -2y \cos x \end{pmatrix}$$

$$= (\alpha - 2)y \cos x \hat{k}$$

$$= 0 \text{ iff } \alpha = 2$$

3

$$(ii) \underline{I} = \int_{(0,0)}^{(\frac{\pi}{2}, 1)} (y^2 \cos x) dx + (\beta y \sin x) dy$$

(a) Consider C to be $y = \frac{2}{\pi}x$ ($0 \leq x \leq \frac{\pi}{2}$)Then subst for y to get :

$$\underline{I} = \int_0^{\pi/2} \left(\left(\frac{2}{\pi} \right)^2 x^2 \cos x + \beta \left(\frac{2}{\pi} \right) x \sin x \left(\frac{2}{\pi} \right) \right) dx$$

$$= \left(\frac{2}{\pi} \right)^2 \int_0^{\pi/2} (x^2 \cos x + \beta x \sin x) dx$$

2

$$\text{by parts} = \left(\frac{2}{\pi} \right)^2 \left\{ \underbrace{[x^2 \sin x - \beta x \cos x]_0^{\pi/2}}_{(\frac{\pi/2})^2} - \int_0^{\pi/2} (2x \sin x - \beta \cos x) dx \right\}$$

$$= \left(\frac{2}{\pi} \right)^2 \left\{ \left(\frac{\pi}{2} \right)^2 - 2 \underbrace{[-x \cos x]_0^{\pi/2}}_{\text{zero}} + \int_0^{\pi/2} \cos x dx \right\} + \beta \int_0^{\pi/2} \cos x dx$$

$$= \left(\frac{2}{\pi} \right)^2 \left\{ \left(\frac{\pi}{2} \right)^2 + (\beta - 2) \right\} = 1 + \frac{4}{\pi^2} (\beta - 2)$$

3

(b) let $y = \sin x$ and subst. for y : ($0 \leq x \leq \frac{\pi}{2}$)

$$\underline{I} = \int_0^{\pi/2} (\sin^2 x \cos x + \beta \sin^2 x \cos x) dx$$

$$= (1 + \beta) \left[\frac{\sin^3 x}{3} \right]_0^{\pi/2} = \frac{1}{3} (1 + \beta)$$

3

Equating answers to (a) & (b) : $1 + \frac{4}{\pi^2} (\beta - 2) = \frac{1}{3} + \frac{\beta}{3}$

$$\Rightarrow \beta \left(\frac{4}{\pi^2} - \frac{1}{3} \right) = \frac{8}{\pi^2} - \frac{2}{3} = 2 \left(\frac{4}{\pi^2} - \frac{1}{3} \right)$$

2

$$\Rightarrow \beta = 2$$

FINAL PART: $\text{Curl } \underline{F} = 0 \Leftrightarrow \int_C \underline{F} \cdot d\underline{r}$ is path independentSo for $\beta = 2$ the answers to (a) & (b) must agree.

2

Setter : WALTON

Setter's signature : *Chelwa Walton*

Checker : JACOBS

Checker's signature : *R.L. Jacobs*Total
15



Please write on this side only, legibly and neatly, between the margins

Bookwork

$$\oint_C (P dx + Q dy) = \iint_R (Q_x - P_y) dx dy$$

Define a vector $\underline{u} = \underline{\hat{i}}Q - \underline{\hat{j}}P$

$$\therefore \text{div } \underline{u} = Q_x - P_y$$



Unit tangent vector $\underline{\hat{t}}$ defined as

$$\underline{\hat{t}} = \frac{d\underline{r}}{ds} = \underline{\hat{i}} \frac{dx}{ds} + \underline{\hat{j}} \frac{dy}{ds}$$

Unit normal $\underline{\hat{n}}$ satisfies $\underline{\hat{n}} \cdot \underline{\hat{t}} = 0$


$$\text{so } \underline{\hat{n}} = \pm \left(\underline{\hat{j}} \frac{dx}{ds} + \underline{\hat{i}} \frac{dy}{ds} \right)$$

$$\therefore \underline{u} \cdot \underline{\hat{n}} = Q \frac{dy}{ds} + P \frac{dx}{ds}$$

\therefore G.T. can be re-expressed as

$$\oint_C (\underline{u} \cdot \underline{\hat{n}}) ds = \iint_R (\text{div } \underline{u}) dx dy$$

$$\underline{u} = \underline{\hat{i}}x^2 + \underline{\hat{j}}y^2 \Rightarrow \text{div } \underline{u} = 2(x+y)$$



$$\therefore \iint_R \text{div } \underline{u} dx dy = 2 \iint_R (x+y) dx dy$$

$$= 2 \iint_R r(\cos\theta + \sin\theta) r dr d\theta$$

$$= 2 \int_0^1 r^2 dr \int_0^{\pi/2} (\cos\theta + \sin\theta) d\theta$$

$dx dy = r dr d\theta$
Because
 $dx dy = \begin{vmatrix} c & s \\ -rs & rs \end{vmatrix} dr d\theta$
 $= r dr d\theta$

Setter : J.D. GIBBON

Checker: NERRANT

Setter's signature: J.D. Gibbon

Checker's signature: NERRANT

MATHEMATICS FOR ENGINEERING STUDENTS

EXAMINATION QUESTION / SOLUTION

SESSION : 2000 - 2001

PAPER

EE II (3)

QUESTION

Please write on this side only, legibly and neatly, between the margins

SOLUTION

Stats 31

$$P(A_i|B) = P(A_i \cap B) / P(B) = P(B|A_i) P(A_i) / P(B) \text{ and}$$

$$\text{by law of total probabilities } P(B) = \sum_{j=1}^k P(B|A_j) P(A_j).$$

$$\text{Hence } P(A_i|B) = \frac{P(B|A_i) P(A_i)}{\sum_{j=1}^k P(B|A_j) P(A_j)}$$

Let OK = test says 'OK' ; F = faulty disk. Then

$$P(OK|\bar{F}) = 0.95 \quad P(\bar{OK}|F) = 0.8 \quad P(F) = 0.05$$

$$\text{so } P(OK|F) = 0.2 \quad P(\bar{F}) = 0.95$$

$$(i) P(OK) = P(OK|F) P(F) + P(OK|\bar{F}) P(\bar{F})$$

$$= (0.2 \times 0.05) + (0.95 \times 0.95) = \underline{0.9125}$$

$$(ii) P(F|OK) = \frac{P(OK|F) P(F)}{P(OK|F) P(F) + P(OK|\bar{F}) P(\bar{F})} = \frac{0.2 \times 0.05}{0.9125}$$

$$= \frac{0.01}{0.9125}$$

$$= \underline{0.01096} \text{ to 5 dp}$$

$$(iii) P(\bar{OK}_2|F) = 0.99$$

$$P(\text{accepted}|F) = P(OK_1 \cap OK_2|F) \quad (OK_1 \equiv OK)$$

$$= P(OK_1|F) P(OK_2|F)$$

$$= 0.2 \times 0.01$$

$$= \underline{0.002}.$$

Setter : ATWalden

Setter's signature : ATW

Checker : D J H A N D

Checker's signature : DJH

MATHEMATICS FOR ENGINEERING STUDENTS

EXAMINATION QUESTION / SOLUTION

SESSION : 2000 - 2001

PAPER

EEII(3)

QUESTION

Please write on this side only, legibly and neatly, between the margins

SOLUTION

Stats 32

i) The coefficient of correlation is a measure of the strength of the linear relationship between two random variables. If $|\rho_{x,y}| = 1$ they are perfectly linearly related, if $\rho_{x,y} = 0$, they are not linearly related.

4

ii) If X and Y are independent $\rho_{x,y} = 0$.

2

$$(iii) f_{x,y}(x,y) = \begin{cases} x^{-1} & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$E\{XY\} = \int_{x=0}^1 \int_{y=0}^x xyx^{-1} dx dy = \int_{x=0}^1 \int_{y=0}^x y dx dy = \int_{x=0}^1 \left. \frac{y^2}{2} \right|_0^x dx = \int_{x=0}^1 \frac{x^2}{2} dx = \left. \frac{x^3}{6} \right|_0^1 = \frac{1}{6}$$

$$E\{X\} = \int_{x=0}^1 \int_{y=0}^x 1 dx dy = \int_0^1 x dx = \left. \frac{x^2}{2} \right|_0^1 = \frac{1}{2}$$

$$E\{Y\} = \int_{x=0}^1 \int_{y=0}^x \frac{y}{x} dx dy = \int_{x=0}^1 \left. \frac{y^2}{2x} \right|_0^x dx = \int_{x=0}^1 \frac{1}{x} \cdot \frac{x^2}{2} dx = \int_0^1 \frac{x}{2} dx = \left. \frac{x^2}{4} \right|_0^1 = \frac{1}{4}$$

So

$$\text{Cov}\{X,Y\} = \frac{1}{6} - \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{24}$$

7

(iv) $\rho_{x,y} \neq 0$ since $\text{Cov}\{X,Y\} = \frac{1}{24}$. Hence X and Y are not independent.

2

(15)

Setter : AT Walden

Setter's signature : ATW

Checker : DJHAND

Checker's signature : DJH