

UNIVERSITY OF LONDON

[II(3)E 2002]

B.ENG. AND M.ENG. EXAMINATIONS 2002

For Internal Students of the Imperial College of Science, Technology and Medicine

This paper is also taken for the relevant examination for the Associateship.

PART II : MATHEMATICS 3 (ELECTRICAL ENGINEERING)

Wednesday 29th May 2002 2.00 - 5.00 pm

Answer EIGHT questions.

Answers to Section A questions must be written in a different answer book from answers to Section B questions.

[Before starting, please make sure that the paper is complete; there should be 7 pages, with a total of 12 questions. Ask the invigilator for a replacement if your copy is faulty.]

Copyright of the University of London 2002

1. Show that the function

$$u(x, y) = \sin x \cosh y + 2 \cos x \sinh y$$

satisfies Laplace's equation.

By integrating the Cauchy-Riemann equations directly, find the conjugate function $v(x, y)$ and hence show that $w = u + iv$ can be expressed as

$$w = (1 - 2i) \sin z + ic$$

where $z = x + iy$ and c is an arbitrary real constant.

2. (i) Consider a complex function $f(z)$ which can be written in the form

$$f(z) = \frac{1}{g(z)},$$

where $g(z)$ has a simple zero at $z = a$.

Show that the residue of $f(z)$ at $z = a$ is $1/g'(a)$.

If, instead, $g(z)$ has a double zero at $z = a$, show that the residue at $z = a$ is

$$-\frac{2}{3} \frac{g'''(a)}{[g''(a)]^2}.$$

(ii) If

$$f(z) = \frac{1}{z^4 + 1},$$

show that of the four simple poles lying on the unit circle, two are located in the upper half-plane at

$$z = e^{i\pi/4} \text{ and } z = e^{3i\pi/4}.$$

By considering a semi-circular contour in the upper half-plane, show that

$$\int_{-\infty}^{\infty} \frac{dx}{x^4 + 1} = \frac{\pi}{\sqrt{2}}.$$

Recall that the residue of a complex function $f(z)$ at a pole $z = a$ of multiplicity m is given by the expression

$$\lim_{z \rightarrow a} \frac{1}{(m-1)!} \left[\frac{d^{m-1}}{dz^{m-1}} \{(z-a)^m f(z)\} \right].$$

PLEASE TURN OVER

3. Consider the contour integral

$$\oint_C \frac{e^{iz} dz}{z(z^2 + 1)}$$

where the contour C is a semi-circle in the upper half of the complex plane, with an additional small semi-circular deformation below the pole at $z = 0$.

Use the Residue Theorem to show that

$$\int_0^\infty \frac{\sin x dx}{x(x^2 + 1)} = \frac{\pi(e - 1)}{2e}.$$

4. The tent function $\Lambda(t)$, the sinc-function $\text{sinc } t$, and the square wave $\Pi(t)$, are defined respectively by

$$\Lambda(t) = \begin{cases} 1+t, & -1 \leq t \leq 0, \\ 1-t, & 0 \leq t \leq 1, \\ 0, & \text{otherwise,} \end{cases}$$

$$\text{sinc } t = \frac{\sin(t/2)}{(t/2)},$$

and

$$\Pi(t) = \begin{cases} 1, & -1/2 \leq t \leq 1/2, \\ 0, & \text{otherwise.} \end{cases}$$

Show that the Fourier transform of $\Pi(t)$ is given by

$$\bar{\Pi}(\omega) = \text{sinc } \omega$$

and the Fourier transform of $\Lambda(t)$ is given by

$$\bar{\Lambda}(\omega) = \text{sinc}^2 \omega.$$

5. (i) The Laplace transform $\mathcal{L}\{y(t)\}$ of a function $y(t)$ is denoted as

$$\bar{y}(s) = \mathcal{L}\{y(t)\} = \int_0^\infty y(t)e^{-st} dt.$$

Show that, for $Re(s) > 0$,

$$\mathcal{L}\{\dot{y}(t)\} = s\bar{y}(s) - y(0),$$

$$\mathcal{L}\{\ddot{y}(t)\} = s^2\bar{y}(s) - sy(0) - \dot{y}(0),$$

provided $y(t)$ and $\dot{y}(t)$ vanish sufficiently fast as $t \rightarrow \infty$.

- (ii) A function $y(t)$ satisfies the differential equation

$$\ddot{y} + 3\dot{y} + 2y = f(t),$$

subject to the initial conditions $\dot{y}(0) = 0$ and $y(0) = \alpha$, a constant. $f(t)$ is a given function that is not specified here. Using a Laplace transform and the Laplace Convolution Theorem, obtain the solution of this differential equation in the form

$$y(t) = \alpha \left(2e^{-t} - e^{-2t} \right) + \int_0^t \left\{ e^{-(t-u)} - e^{-2(t-u)} \right\} f(u) du.$$

6. If a function $f(t)$ is periodic in time t with fixed period T such that $f(t) = f(t - T)$ with $T > 0$, show that for $s > 0$ its Laplace transform $\bar{f}(s)$ is given by

$$\bar{f}(s) = \frac{1}{1 - \exp(-sT)} \int_0^T f(t) \exp(-st) dt.$$

If $f(t)$ is the periodic square wave of period T

$$f(t) = \begin{cases} 1, & 0 \leq t \leq T/2, \\ 0, & T/2 < t \leq T, \end{cases}$$

show that its Laplace transform $\bar{f}(s)$ is given by

$$\bar{f}(s) = \frac{1}{s} \left(\frac{1 - \exp(-sT/2)}{1 - \exp(-sT)} \right).$$

Explain what happens in the limit $T \rightarrow \infty$.

PLEASE TURN OVER

7. The double integral I_1 is given by

$$I_1 = \int \int_A e^{-\alpha(x^2+y^2)} dx dy ,$$

where A is the finite region enclosed by the curve $x^2 + y^2 = R^2$.

Sketch the region of integration A and, upon using the substitution $x = r \cos \theta$, $y = r \sin \theta$ (polar coordinates), show that

$$I_1 = \frac{\pi}{\alpha} (1 - e^{-\alpha R^2}).$$

Calculate $\lim_{R \rightarrow \infty} I_1$ and hence deduce the value of the integral

$$I_2 = \int_{-\infty}^{+\infty} e^{-\alpha x^2} dx .$$

Hint: relate I_2^2 to I_1 .

8. (i) Show that

$$\operatorname{div}(\operatorname{curl}\mathbf{A}) = 0, \quad \operatorname{curl}(\nabla\phi) = \mathbf{0}$$

for a general three-dimensional vector field \mathbf{A} and scalar field ϕ .

(ii) Suppose that \mathbf{A} satisfies the relation

$$\operatorname{curl}\mathbf{A} = \lambda\mathbf{A}$$

for some scalar λ .

Prove the following results:

(a) $\operatorname{div}\mathbf{A} = 0$;

(b) $\nabla^2\mathbf{A} + \lambda^2\mathbf{A} = \mathbf{0}$;

(c) $\lambda = \frac{\operatorname{curl}\mathbf{A} \cdot \operatorname{curl}(\operatorname{curl}\mathbf{A})}{(\operatorname{curl}\mathbf{A}) \cdot (\operatorname{curl}\mathbf{A})}$.

You may assume the vector identity: $\operatorname{curl}(\operatorname{curl}\mathbf{A}) = \nabla(\operatorname{div}\mathbf{A}) - \nabla^2\mathbf{A}$.

9. (i) Show that

$$\int_C \{(x^2 + 2xy + 3y^2)dx + (x^2 + 6xy + 2y^2)dy\}$$

is independent of the path C , joining the initial point to the final point.

Evaluate the integral for a path C from $(0, 0)$ to $(1, 2)$.

- (ii) Let C be a circle in the $x - y$ plane with centre at the origin and radius 1, described in the counter-clockwise direction. Let R be the region inside the circle.

Evaluate

$$\oint_C \{(2x - y)dx + (x - 2y)dy\}$$

(a) directly, and

(b) using Green's Theorem.

Green's Theorem in the plane states that:

$$\oint_C (Pdx + Qdy) = \iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dxdy.$$

10. Consider a two-dimensional region R bounded by a closed piecewise smooth curve C . Using Green's Theorem in a plane (see below), choose the components of a vector field $\mathbf{v}(x, y)$ in terms of $P(x, y)$ and $Q(x, y)$ to prove the two-dimensional form of Stokes's Theorem

$$\iint_R \mathbf{k} \cdot (\operatorname{curl} \mathbf{v}) dxdy = \oint_C \mathbf{v} \cdot d\mathbf{r} \quad (1)$$

where $\mathbf{r} = xi + yj$.

If $\mathbf{v} = y^2i + x^2j$, and R is the finite region bounded by the hyperbola $y = \frac{1}{4x}$, and the lines $x = 1$ and $y = x$, sketch the region R in the $x - y$ plane and, by evaluating the line integral on the right hand side of (1), or otherwise, show that

$$\iint_R \mathbf{k} \cdot (\operatorname{curl} \mathbf{v}) dxdy = \frac{5}{48}.$$

Green's Theorem in a plane states that for a two-dimensional region R bounded by a closed, piecewise smooth curve C , then

$$\oint_C \{P(x, y)dx + Q(x, y)dy\} = \iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dxdy.$$

PLEASE TURN OVER

SECTION B

[II(3)E 2002]

11. (i) Events E_1 and E_2 are exclusive and E_2 and E_3 are independent. Draw a Venn diagram to illustrate the most general structure for these events. Suppose that $\text{pr}(E_1) = p_1$, $\text{pr}(E_2) = p_2$, $\text{pr}(E_3) = p_3$ and $\text{pr}(E_1 | E_3) = p_{13}$. Express $\text{pr}(E_2 | E_3)$, $\text{pr}(E_1 \cap E_3)$, $\text{pr}(E_3 | E_1)$ and $\text{pr}(E_1 | E_2)$ in terms of p_1 , p_2 , p_3 and p_{13} .
- (ii) The insulating material in a transformer is subject to three types of deterioration, D_1 , D_2 and D_3 . From extensive historical data, the probabilities are well estimated as $\text{pr}(D_1) = 0.1$, $\text{pr}(D_2) = 0.05$, $\text{pr}(D_3) = 0.01$, and $\text{pr}(D_1 \cup D_2) = 0.12$. It is also known that D_1 and D_3 occur independently, and that D_2 and D_3 are exclusive. Compute the probabilities of the following states of deterioration:
- (a) D_1 and D_3 both present;
 - (b) D_1 and D_2 both present;
 - (c) D_1 , D_2 and D_3 all present;
 - (d) D_2 present, given that D_1 is present;
 - (e) D_1 present, given that D_3 is present.
12. (i) The discrete random variables X_1 and X_2 have joint probability function $p(x_1, x_2)$ and marginal probability functions $p_1(x_1)$ and $p_2(x_2)$, respectively. Write down an expression that defines the expected value $E(X_1)$ and prove that $E(X_1 + X_2) = E(X_1) + E(X_2)$.
- (ii) The random variables X_1 and X_2 are independent with distributions $N(\mu_1, \sigma_1^2)$ and $N(\mu_2, \sigma_2^2)$, respectively.
What are the distributions of $X_1 + X_2$ and $2X_1 - 3X_2$?
- (iii) The random variable X_3 has mean μ_3 and variance σ_3^2 , X_4 has mean μ_4 and variance σ_4^2 and X_3 and X_4 are correlated with correlation coefficient ρ . Calculate the mean and variance of $2X_3 - 3X_4$.
- (iv) The energy requirements of two systems, which have to satisfy varying loads, are random variables with distributions well-approximated by $N(1.41, 0.11)$ and $N(1.11, 0.07)$. Because of other demands, which themselves vary randomly, the available supply is distributed as $N(3.66, 0.18)$. The two system requirements and the supply are independent random variables.
What is the probability that the supply can meet the demand from the two systems?

For part (iv) you may use the following table of values of the standard normal distribution function.

x	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
$\Phi(x)$	0.5000	0.5398	0.5793	0.6179	0.6554	0.6915	0.7257	0.7580	0.7881	0.8159
x	1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9
$\Phi(x)$	0.8413	0.8643	0.8849	0.9032	0.9192	0.9332	0.9452	0.9554	0.9641	0.9713
x	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9
$\Phi(x)$	0.9772	0.9821	0.9861	0.9893	0.9918	0.9938	0.9953	0.9965	0.9974	0.9981

END OF PAPER

MATHEMATICS DEPARTMENT

3. TRIGONOMETRIC IDENTITIES AND HYPERBOLIC FUNCTIONS

MATHEMATICAL FORMULAE

1. VECTOR ALGEBRA

$$\mathbf{a} = a_1 \mathbf{i} + a_2 \mathbf{j} + a_3 \mathbf{k} = (a_1, a_2, a_3)$$

$$\mathbf{a} \cdot \mathbf{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$$

Scalar (dot) product:

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

Scalar triple product:

$$[\mathbf{a}, \mathbf{b}, \mathbf{c}] = \mathbf{a} \cdot \mathbf{b} \times \mathbf{c} = \mathbf{b} \cdot \mathbf{c} \times \mathbf{a} = \mathbf{c} \cdot \mathbf{a} \times \mathbf{b} = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

Vector triple product: $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{c} \cdot \mathbf{a})\mathbf{b} - (\mathbf{b} \cdot \mathbf{a})\mathbf{c}$

2. SERIES

$$(1+x)^\alpha = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!} x^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!} x^3 + \dots \quad (\alpha \text{ arbitrary, } |x| < 1)$$

$$e^x = 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} + \dots ,$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + (-1)^n \frac{x^{2n}}{(2n)!} + \dots ,$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + \dots ,$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^n \frac{x^{n+1}}{(n+1)} + \dots \quad (-1 < x \leq 1)$$

4. DIFFERENTIAL CALCULUS

(a) Leibniz's formula:

$$D^n(fg) = f D^n g + (n) Df D^{n-1} g + \dots + (n) D^r f D^{n-r} g + \dots + D^n f g .$$

(b) Taylor's expansion of $f(x)$ about $x = a$:

$$f(a+h) = f(a) + hf'(a) + h^2 f''(a)/2! + \dots + h^n f^{(n)}(a)/n! + \epsilon_n(h),$$

where $\epsilon_n(h) = h^{n+1} f^{(n+1)}(a+\theta h)/(n+1)!$, $0 < \theta < 1$.

(c) Taylor's expansion of $f(x, y)$ about (a, b) :

$$f(a+h, b+k) = f(a, b) + [h f_x + k f_y]_{a,b} + 1/2! [h^2 f_{xx} + 2hk f_{xy} + k^2 f_{yy}]_{a,b} + \dots$$

(d) Partial differentiation of $f(x, y)$:

i. If $y = y(x)$, then $f = F(x)$, and $\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{dy}{dx}$.

ii. If $x = x(t)$, $y = y(t)$, then $f = F(t)$, and $\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$.

iii. If $x = x(u, v)$, $y = y(u, v)$, then $f = F(u, v)$, and

$$\frac{\partial F}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u}, \quad \frac{\partial F}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v}.$$

(e) Stationary points of $f(x, y)$ occur where $f_x = 0$, $f_y = 0$ simultaneously. Let (a, b) be a stationary point: examine $D = [f_{xx} f_{yy} - (f_{xy})^2]_{a,b}$.

If $D > 0$ and $f_{xx}(a, b) < 0$, then (a, b) is a maximum;

If $D > 0$ and $f_{xx}(a, b) > 0$, then (a, b) is a minimum;

If $D < 0$ then (a, b) is a saddle-point.

(f) Differential equations:

- i. The first order linear equation $dy/dx + P(x)y = Q(x)$ has an integrating factor $I(x) = \exp[\int P(x)(dx)]$, so that $\frac{d}{dx}(Iy) = IQ$.
- ii. $P(x, y)dx + Q(x, y)dy = 0$ is exact if $\partial Q/\partial x = \partial P/\partial y$.

5. INTEGRAL CALCULUS

- (a) An important substitution: $\tan(\theta/2) = t$:
 $\sin \theta = 2t/(1+t^2)$, $\cos \theta = (1-t^2)/(1+t^2)$, $d\theta = 2dt/(1+t^2)$.

- (b) Some indefinite integrals:

$$\int (a^2 - x^2)^{-1/2} dx = \sin^{-1}\left(\frac{x}{a}\right), \quad |x| < a.$$

$$\int (a^2 + x^2)^{-1/2} dx = \sinh^{-1}\left(\frac{x}{a}\right) = \ln \left\{ \frac{x}{a} + \left(1 + \frac{x^2}{a^2}\right)^{1/2} \right\}.$$

$$\int (x^2 - a^2)^{-1/2} dx = \cosh^{-1}\left(\frac{x}{a}\right) = \ln \left| \frac{x}{a} + \left(\frac{x^2}{a^2} - 1 \right)^{1/2} \right|.$$

$$\int (a^2 + x^2)^{-1} dx = \left(\frac{1}{a} \right) \tan^{-1}\left(\frac{x}{a}\right).$$

7. LAPLACE TRANSFORMS

Function	Transform	Function	Transform	Function	Transform
$f(t)$	$F(s) = \int_0^\infty e^{-st} f(t) dt$	$a f(t) + b g(t)$	$a F(s) + b G(s)$	$s^2 F(s) - s f(0) - f'(0)$	$s^2 F(s) + b G(s)$
df/dt	$sF(s) - f(0)$	$d^2 f/dt^2$	$-dF(s)/ds$	$t f(t)$	$F(s-a)$
$e^{at} f(t)$	$F(s-a)$	$t f(t)$	$F(s-a)$	$F'(s)/s$	$F'(s)/s$
$(\partial/\partial \alpha) f(t, \alpha)$	$(\partial/\partial \alpha) F(s, \alpha)$	$f_0' f(t) dt$	$F(s)G(s)$	$t^n (n = 1, 2, \dots)$	$n!/s^{n+1}, \quad (s > 0)$
$\int_0^t f(u)g(t-u) du$					$\omega/(s^2 + \omega^2), \quad (s > 0)$
1	$1/s$				
e^{at}	$1/(s-a), \quad (s > a)$			$\sin \omega t$	
$\cos \omega t$	$s/(s^2 + \omega^2), \quad (s > 0)$	$H(t-T) = \begin{cases} 0, & t < T \\ 1, & t > T \end{cases}$		$e^{-sT}/s, \quad (s, T > 0)$	

6. NUMERICAL METHODS

- (a) Approximate solution of an algebraic equation:

If a root of $f(x) = 0$ occurs near $x = a$, take $x_0 = a$ and
 $x_{n+1} = x_n - [f(x_n)/f'(x_n)], \quad n = 0, 1, 2, \dots$

(Newton Raphson method).

- (b) Formulae for numerical integration: Write $x_n = x_0 + nh$, $y_n = y(x_n)$.

i. Trapezium rule (1-strip): $\int_{x_0}^{x_1} y(x) dx \approx (h/2) [y_0 + y_1]$.

ii. Simpson's rule (2-strip): $\int_{x_0}^{x_2} y(x) dx \approx (h/3) [y_0 + 4y_1 + y_2]$.

- (c) Richardson's extrapolation method: Let $I = \int_a^b f(x) dx$ and let I_1, I_2 be two estimates of I obtained by using Simpson's rule with intervals h and $h/2$. Then, provided h is small enough,

$$I_2 + (I_2 - I_1)/15,$$

is a better estimate of I .

8. FOURIER SERIES

If $f(x)$ is periodic of period $2L$, then $f(x+2L) = f(x)$, and

$$f(x) = \frac{1}{2} a_0 + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{L} + \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{L}, \quad \text{where}$$

(Parseval's theorem)

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx, \quad n = 0, 1, 2, \dots, \text{ and}$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx, \quad n = 1, 2, 3, \dots$$

Parseval's theorem

estimates of I obtained by using Simpson's rule with intervals h and $h/2$.

Then, provided h is small enough,

$$\frac{1}{L} \int_{-L}^L [f(x)]^2 dx = \frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2).$$

September 2000

(EE
 Stream)

EXAMINATION QUESTION / SOLUTION

SESSION : 2001-2002

E 1

Please write on this side only, legibly and neatly, between the margins

QUESTION

SOLUTION

15

$$u = \sin x \cosh y + 2 \cos x \sinh y$$

$$u_x = \cos x \cosh y - 2 \sin x \sinh y, \quad u_y = \sin x \sinh y + 2 \cos x \cosh y$$

$$u_{xx} = -\sin x \cosh y - 2 \cos x \sinh y; \quad u_{yy} = \sin x \cosh y + 2 \cos x \sinh y$$

$$\therefore \underline{u_{xx} + u_{yy} = 0}$$

5

Hence \exists a conjugate function v such that (CR equation)

$$v_y = u_x = \cos x \cosh y - 2 \sin x \sinh y \quad 1)$$

$$v_x = -u_y = -\sin x \sinh y - 2 \cos x \cosh y \quad 2)$$

$$\text{Integrate: } 1) \rightarrow v = \cos x \sinh y - 2 \sin x \cosh y + A(x)$$

$$2) \rightarrow v = \cos x \sinh y - 2 \sin x \cosh y + B(y)$$

$$\therefore A = B = \text{const} = c$$

$$\therefore \underline{v = \cos x \sinh y - 2 \sin x \cosh y + c}$$

5

$$\therefore w = u + iv = \sin x \cosh y + 2 \cos x \sinh y + i(\cos x \sinh y - 2 \sin x \cosh y) + ic$$

$$= (1-2i)\sin x \cosh y + (2+i)\cos x \sinh y + c,$$

2

$$= (1-2i)\{\sin x \cosh y + i \cos x \sinh y\} + c,$$

$$\text{Now } \cos(iy) = \frac{1}{2}(e^{i(iy)} + e^{-i(iy)}) = \frac{1}{2}(e^y + e^{-y}) = \cosh y$$

$$\sin(iy) = \frac{1}{2i}(e^{i(iy)} - e^{-i(iy)}) = \frac{1}{2i}(e^{-y} - e^y) = i \sinh y$$

2

$$\therefore w = (1-2i)\{\sin x \cos(iy) + \cos x \sin(iy)\} + c_1$$

$$= (1-2i)\sin(x+iy) + c_1$$

1

$$\approx (1-2i)\sin z + c_1$$

Setter : J. D. GIBBON

Setter's signature: J. D. Gibson

Checker: NEERAJ

Checker's signature: Dr. Neeraj

EXAMINATION QUESTION / SOLUTION

SESSION : 2001-2002

E 2

3

QUESTION

Please write on this side only, legibly and neatly, between the margins

SOLUTION

16

i) $f = \frac{1}{g(z)}$; Expand $g(z)$ as a Taylor series about $z=a$

$$g(z) = g(a) + (z-a)g'(a) + \frac{1}{2}(z-a)^2g''(a) + \frac{1}{6}(z-a)^3g'''(a) \dots$$

i) If g has a simple zero at $z=a$ then $g(a)=0$ butii) if g has a double zero at $z=a$ then $g'(a)=0$ also.

$$\therefore i) f(z) = [(z-a)g'(a) + \dots]^{-1}$$

so Res of $f(z)$ at $z=a$ is $\frac{1}{g'(a)}$

3

ii) Res. of $f(z)$ at $z=a$


with a double pole is $\lim_{z \rightarrow a} \frac{d}{dz} \left\{ \frac{(z-a)^2}{\frac{1}{2}(z-a)^2g''(a) + \frac{1}{6}(z-a)^3g'''(a) \dots} \right\}$

$$= \lim_{z \rightarrow a} \frac{d}{dz} \left[\frac{1}{2}g''(a) + \frac{1}{6}(z-a)g'''(a) + \dots \right]^{-1}$$

4

$$= -\frac{1}{6}g'''(a) \left[\frac{1}{2}g''(a) \right]^{-2} = -\frac{2}{3} \cdot \frac{g'''(a)}{(g''(a))^2}$$

iii) $\int_C \frac{dz}{z^4+1} = \lim_{R \rightarrow \infty} \left(\int_{-R}^R \frac{dx}{x^4+1} + \int_{R_R}^R \frac{dz}{z^4+1} \right)$



z^4 has simple zeros at $\left\{ e^{\pi i/4}, e^{3\pi i/4} \text{ (upper half-plane)}, e^{5\pi i/4}, e^{7\pi i/4} \text{ (lower half-plane)} \right\}$

$$\lim_{R \rightarrow \infty} \int_{R_R}^R \frac{dz}{z^4+1} = 0$$

4

From formula above :

Res. at $e^{\pi i/4}$ is $\frac{1}{4e^{3\pi i/4}}$ Res. at $e^{3\pi i/4}$ is $1/4e^{9\pi i/4}$

$$\therefore \int_{-\infty}^{\infty} \frac{dx}{x^4+1} = \frac{2\pi i}{4} \left\{ e^{-\frac{3\pi i}{4}} + e^{-\frac{9\pi i}{4}} \right\}$$

$$\text{Now } \left\{ e^{-3\pi i/4} + e^{-9\pi i/4} \right\} = \left\{ -e^{\pi i/4} + e^{-\pi i/4} \right\} = -2i \sin \pi/4 = -2i/\sqrt{2}$$

4

$$\therefore \int_{-\infty}^{\infty} \frac{dx}{x^4+1} = \frac{2\pi}{2\sqrt{2}} = \frac{\pi}{\sqrt{2}}$$

4

Setter : J. D. GIBBON

Setter's signature : J.D. Gibbon

Checker: NEERAJ

Checker's signature : Dr. Albert

MATHEMATICS FOR ENGINEERING STUDENTS
 EXAMINATION QUESTION / SOLUTION
 SESSION : 2001-2002


PAPER

E 3

3
 QUESTION

Please write on this side only, legibly and neatly, between the margins

$$\oint_C \frac{e^{iz}}{z(z^2+1)} dz$$



SOLUTION
 17

Within C there are

two simple poles : $z=0$ and $z=i$

i) Res. at $z=0$ is 1

ii) Res. at $z=i$ is $\frac{e^{-1}}{ix^2} = -\frac{1}{2}e^{-1}$] sum is $1 - \frac{1}{2}e^{-1}$

3

$$\oint_C = \int_{-R}^{-r} + \int_Y + \int_r^R + \int_{H_R} \quad \text{Take } R \rightarrow \infty \\ r \rightarrow 0$$

a) By Jordan's Lemma $\int_{H_R} \rightarrow 0$ as $R \rightarrow \infty$ because, for

integrals of the type $\int_{H_R} e^{iax} f(x) dx$ i) $a=1 > 0$

$$ii) f(z) = \frac{1}{z(z^2+1)} \rightarrow 0$$

as $z \rightarrow \infty$ with the degree of denominator > 1 .

iii) $f(z)$ has only poles in upper half-plane.

3

b) Small semi-circle is $z=r e^{i\theta}$, $\pi \leq \theta \leq 2\pi$

$$\lim_{r \rightarrow 0} \int \frac{e^{iz}}{z(z^2+1)} dz = \lim_{r \rightarrow 0} \int_{\pi}^{2\pi} \frac{\exp(ir e^{i\theta}) i e^{i\theta} d\theta}{r^2 e^{2i\theta} + 1}$$

$$= i \int_{\pi}^{2\pi} d\theta = \pi i$$

3

$$c) \lim_{\substack{R \rightarrow \infty \\ r \rightarrow 0}} \left(\int_{-R}^{-r} + \int_r^R \right) \frac{e^{ix}}{x(x^2+1)} = \int_{-\infty}^{\infty} \frac{e^{ix}}{x(x^2+1)} dx$$

$$= \int_0^{\infty} \frac{(e^{ix} - e^{-ix})}{x(x^2+1)} dx = 2i \int_0^{\infty} \frac{\sin x}{x(x^2+1)} dx$$

3

Finally, using the Residue Theorem we have

$$2\pi i (1 - \frac{1}{2}e^{-1}) = \pi i + 2i \int_0^{\infty} \frac{\sin x}{x(x^2+1)} dx$$

3

$$\text{so } \int_0^{\infty} \frac{\sin x}{x(x^2+1)} dx = \frac{\pi}{2} \left(1 - \frac{1}{2}e^{-1} \right)$$

Setter : J.D. GIBBON

Setter's signature : J.D. Gibbon

Checker: N. N. B. S. M.

Checker's signature : Dr. N. N. B. S. M.

EXAMINATION QUESTION / SOLUTION

SESSION : 2001-2002

E4

3

QUESTION

Please write on this side only, legibly and neatly, between the margins

SOLUTION

18

$$\begin{aligned}
 \bar{\pi}(\omega) &= \int_{-\infty}^{\infty} e^{-i\omega t} \pi(t) dt \\
 &= \int_{-1/\omega}^{1/\omega} 1 \cdot e^{-i\omega t} dt + 0 \\
 &= -\frac{1}{i\omega} [e^{-i\omega/2} - e^{i\omega/2}] = \frac{\sin \omega/2}{\omega/2} \\
 &= \text{sinc } \omega
 \end{aligned}$$

5

$$\begin{aligned}
 \bar{\Delta}(\omega) &= \int_{-\infty}^{\infty} e^{-i\omega t} \Delta(t) dt \\
 &= \int_{-1}^0 (1+t) e^{-i\omega t} dt + \int_0^1 (1-t) e^{-i\omega t} dt
 \end{aligned}$$

5

$$\begin{aligned}
 \int e^{-i\omega t} dt &= \frac{i}{\omega} [e^{-i\omega t}] \\
 \int t e^{-i\omega t} dt &= \frac{i}{\omega} \int t d(e^{-i\omega t}) = \frac{i}{\omega} [t e^{-i\omega t}] - \frac{i}{\omega} \int e^{-i\omega t} dt \\
 &= \frac{i}{\omega} [t e^{-i\omega t}] + \frac{1}{\omega^2} [e^{-i\omega t}]
 \end{aligned}$$

$$\begin{aligned}
 \therefore \bar{\Delta}(\omega) &= \frac{i}{\omega} [e^{-i\omega} - e^{i\omega}] + \frac{i}{\omega} [0 + e^{i\omega}] + \frac{1}{\omega^2} [1 - e^{i\omega}] \\
 &\quad - \frac{i}{\omega} [e^{-i\omega} - 0] - \frac{1}{\omega^2} [e^{-i\omega} - 1] \\
 &= \frac{1}{\omega^2} [2 - e^{i\omega} - e^{-i\omega}] = \frac{2}{\omega^2} (1 - \cos \omega)
 \end{aligned}$$

$$\text{But } \cos \omega = 1 - 2 \sin^2 \frac{1}{2} \omega$$

$$\therefore \bar{\Delta}(\omega) = 4 \frac{\sin^2 \frac{1}{2} \omega}{\omega^2} = \text{sinc}^2 \omega$$

5

Setter : J. D. GIBBON

Setter's signature : J.D. Gibbon

Checker: NEERAJ

Checker's signature : Dr. Herbert

EXAMINATION QUESTION / SOLUTION

SESSION : 2001-2002

E5

3

QUESTION

Please write on this side only, legibly and neatly, between the margins

$$\begin{aligned} \mathcal{L}(y) &= \int_0^\infty e^{-st} y dt = \int_0^\infty e^{-st} dy \\ &= [ye^{-st}]_0^\infty + s \int_0^\infty y e^{-st} dt \quad s > 0 \\ &= s \bar{y}(s) - y(0) \end{aligned}$$

SOLUTION
19

$$\begin{aligned} \mathcal{L}(\ddot{y}) &= \int_0^\infty e^{-st} \ddot{y} dt = \int_0^\infty e^{-st} d(\dot{y}) \\ &= [\dot{y} e^{-st}]_0^\infty + s \int_0^\infty e^{-st} \dot{y} dt \\ &= -\dot{y}(0) + s(s \bar{y}(s) - y(0)) \quad s > 0 \\ &= s^2 \bar{y}(s) - \dot{y}(0) - s y(0) \end{aligned}$$

2

3

(ii) Now apply above formulae to $\ddot{y} + 3\dot{y} + 2y = f(t)$

$$\therefore s^2 \bar{y}(s) - \dot{y}(0) - s y(0) + 3s \bar{y}(s) - 3y(0) + 2\bar{y}(s) = \bar{f}(s)$$

$$\therefore (s^2 + 3s + 2) \bar{y}(s) = (s+2) y(0) + \bar{f}(s)$$

$$\therefore \bar{y}(s) = \frac{(s+3) y(0)}{(s+1)(s+2)} + \frac{\bar{f}(s)}{(s+1)(s+2)} = \left(\frac{2}{s+1} - \frac{1}{s+2} \right) y(0) + \frac{\bar{f}(s)}{s+1} - \frac{\bar{f}(s)}{s+2}$$

5

Invert:

$$\begin{aligned} y(t) &= (2e^{-t} - e^{-2t}) y(0) \\ &\quad + \mathcal{L}^{-1}\left(\frac{\bar{f}(s)}{(s+1)}\right) - \mathcal{L}^{-1}\left(\frac{\bar{f}(s)}{s+2}\right) \end{aligned}$$

Convolution Thm: $\mathcal{L}(f * g) = \bar{f}(s) \bar{g}(s)$
(Formula sheet)

5

$$\therefore y(t) = (2e^{-t} - e^{-2t}) y(0) +$$

$$+ \int_0^t (e^{-(t-u)} - e^{-2(t-u)}) f(u) du.$$

Setter: J. D. GIBBON

Setter's signature: J. D. Gibbon

Checker: H. R. B. M.

Checker's signature: Dr. R. B. M.

MATHEMATICS FOR ENGINEERING STUDENTS
 EXAMINATION QUESTION / SOLUTION
 SESSION : 2001-2002

PAPER

E 6

3

Please write on this side only, legibly and neatly, between the margins

QUESTION

SOLUTION

20

$$\int_0^\infty f(t) e^{-st} dt = \int_0^T f(t) e^{-st} dt + \int_T^{2T} f(t) e^{-st} dt + \dots$$

Consider

$$\int_T^{2T} f(t) e^{-st} dt$$

and define $\tau = t - T$, so

$$\begin{aligned} \int_T^{2T} f(t) e^{-st} dt &= \int_0^T e^{-s(\tau+T)} f(\tau+T) d\tau \\ &= e^{-sT} \int_0^T e^{-s\tau} f(\tau) d\tau \end{aligned}$$

invoking periodicity

seen
,
short.

$$\therefore \text{clearly } \int_{nT}^{(n+1)T} = e^{-snT} \int_0^T e^{-st} f(t) dt$$

$$\therefore \int_0^\infty f(t) e^{-st} dt = (1 + e^{-sT} + e^{-2sT} + \dots) \int_0^T$$

$$\text{For } s > 0 \quad = (1 - e^{-sT})^{-1} \int_0^T e^{-st} f(t) dt$$

10

$$\begin{aligned} \int_0^T f(t) e^{-st} dt &= \int_0^{T/2} 1 \cdot e^{-st} dt + 0 = -\frac{1}{s} [e^{-st}]_0^{T/2} \\ &= \frac{1}{s} (1 - e^{-sT/2}) \end{aligned}$$

$$\therefore \bar{f}(s) = \frac{1}{s} \left(\frac{1 - e^{-sT/2}}{1 - e^{-sT}} \right)$$

3

When $T \rightarrow \infty$ ($s > 0$), $\bar{f}(s) \rightarrow \frac{1}{s}$. This is the L.T. of

$f(t) = 1$: the square wave period $\rightarrow \infty$ and $f(t) \rightarrow 1$ uniformly in $0 \leq t \leq \infty$.

2

Setter : J. D. GIBBON

Setter's signature : J. D. Gibbon

Checker: HERBEM

Checker's signature : Herbem

EXAMINATION QUESTION / SOLUTION


SESSION : 2001-2002

E 7

Please write on this side only, legibly and neatly, between the margins

$$x^2 + y^2 = R^2 - \text{circle of radius } R$$

Hence the area of integration



Jacobian: $\frac{\partial x}{\partial r} \frac{\partial y}{\partial \theta} - \frac{\partial x}{\partial \theta} \frac{\partial y}{\partial r} = r(\cos^2 \theta + \sin^2 \theta) = r.$

Region $x^2 + y^2 < R^2 \Rightarrow 0 < \theta < 2\pi, 0 < r < R,$

also $x^2 + y^2 = r^2.$ Therefore

$$I_1 = \int_0^R r dr \int_0^{2\pi} e^{-\alpha r^2} d\theta$$

To calculate I_1 , substitute $r^2 = u:$

$$I_1 = \pi \int_0^{R^2} du e^{-\alpha u} = -\frac{\pi}{\alpha} e^{-\alpha u} \Big|_0^{R^2} = \frac{\pi}{\alpha} (1 - e^{-\alpha R^2})$$

Therefore the limit $I_1(\infty) = \lim_{R \rightarrow \infty} I_1 = \frac{\pi}{\alpha}.$

Observe that

$$I_2^2 = \left[\int_{-\infty}^{+\infty} dx e^{-\alpha x^2} \right]^2 = \int_{-\infty}^{+\infty} dx \int_{-\infty}^{+\infty} dy e^{-\alpha(x^2 + y^2)} = I_1(\infty)$$

Hence $I_2 = \sqrt{\frac{\pi}{\alpha}}$

Setter : Gogolin

Checker: WALTON

Setter's signature: A. Gogolin

Checker's signature: Andrew Walton

QUESTION
21

SOLUTION
21

3

2

4

4

1

4

Total

15

Please write on this side only, legibly and neatly, between the margins

(a) (i) Let $\underline{A} = A_1 \hat{i} + A_2 \hat{j} + A_3 \hat{k}$

Then $\text{Curl } \underline{A} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ A_1 & A_2 & A_3 \end{vmatrix}$

= $\hat{i}(A_{3y} - A_{2z}) - \hat{j}(A_{3x} - A_{1z}) + \hat{k}(A_{2x} - A_{1y})$

So $\text{div}(\text{Curl } \underline{A}) = \frac{\partial}{\partial x}(A_{3y} - A_{2z}) + \frac{\partial}{\partial y}(A_{1z} - A_{3x}) + \frac{\partial}{\partial z}(A_{2x} - A_{1y})$
= 0, as required.

(b) $\text{Curl } \underline{A} = \lambda \underline{A} \quad (*)$

(i) Taking div of both sides:

$$\underbrace{\text{div}(\text{Curl } \underline{A})}_{=0 \text{ by (a)}} = \lambda \text{div} \underline{A} \Rightarrow \underline{\text{div} \underline{A} = 0}$$

(ii) Take curl of both sides:

$$\text{Curl}(\text{Curl } \underline{A}) = \lambda \text{Curl } \underline{A}$$

$$\nabla(\text{div} \underline{A}) - \nabla^2 \underline{A}$$

(given in question)

But $\text{div} \underline{A} = 0 \therefore \nabla^2 \underline{A} = -\lambda \text{Curl } \underline{A} = -\lambda^2 \underline{A}$
Hence $\underline{\nabla^2 \underline{A} + \lambda^2 \underline{A} = 0}$ (using *)

(iii) Using (ii) we have $\text{Curl}(\text{Curl } \underline{A}) = \lambda \text{Curl } \underline{A}$

So RHS of (iii) becomes

$$\frac{(\text{Curl } \underline{A}) \cdot (\lambda \text{Curl } \underline{A})}{(\text{Curl } \underline{A}) \cdot (\text{Curl } \underline{A})} = \underline{\underline{\lambda}}, \text{ as required.}$$

(a) (i)

4 MARKS

$$\left. \begin{array}{l} \text{(a)(ii)} \\ \text{Curl}(\nabla \varphi) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \varphi_x & \varphi_y & \varphi_z \end{vmatrix} \\ = \hat{i}(\varphi_{zy} - \varphi_{yz}) - \hat{j}(\varphi_{zx} - \varphi_{xz}) + \hat{k}(\varphi_{yx} - \varphi_{xy}) \\ = \underline{\underline{0}}, \text{ as required.} \end{array} \right\}$$

Total
15

MATHEMATICS FOR ENGINEERING STUDENTS
 EXAMINATION QUESTION / SOLUTION
 SESSION : 2001-2002

PAPER

3

Please write on this side only, legibly and neatly, between the margins

E 9

QUESTION

SOLUTION
24

(i) We know that $\int_C (f dx + g dy)$ is independent of the path joining the initial point to the final point if $\frac{\partial f}{\partial y} = \frac{\partial g}{\partial x}$. Here, $f = (x^2 + 2xy + 3y^2)$, $g = (x^2 + 6xy + 2y^2)$, so $\frac{\partial f}{\partial y} = 2x + 6y = \frac{\partial g}{\partial x}$, as required.

4

Let the path be $y = 2x$. Then

$$\begin{aligned} \int_C &= \int_0^1 \left\{ (x^2 + 4x^2 + 12x^2) dx + (x^2 + 12x^2 + 8x^2) \cdot 2 dx \right\} \\ &= \int_0^1 59x^2 dx \\ &= \frac{1}{3} \cdot 59. \end{aligned}$$

4

(ii) (a) Put $x = r\cos\theta$, $y = r\sin\theta$

Then $dx = -r\sin\theta d\theta$, $dy = r\cos\theta d\theta$,

$$\begin{aligned} \text{and the integral} &= \int_0^{2\pi} \left\{ -(2r\cos\theta - r\sin\theta)r\sin\theta + (r\cos\theta - 2r\sin\theta)r\cos\theta \right\} d\theta \\ &= \int_0^{2\pi} \left\{ (\cos^2\theta + \sin^2\theta) - 4\sin\theta\cos\theta \right\} d\theta \\ &= \int_0^{2\pi} (1 - 2\sin 2\theta) d\theta \\ &= [\theta + \cos 2\theta]_0^{2\pi} \\ &= 2\pi \end{aligned}$$

4

(b) Apply Green's Theorem with

$$f = 2x-y, \quad g = x-2y.$$

We get

$$\frac{\partial g}{\partial x} = 1, \quad \frac{\partial f}{\partial y} = -1$$

3

$$\therefore \text{The integral} = \iint_R 2 dx dy = 2A \text{ where } A = \text{area of the circle} \\ = 2\pi.$$

Setter : G. D. JAMES

Setter's signature : G. James

Checker: A. J. MESTEL

Checker's signature : A. J. Mestel

EXAMINATION QUESTION / SOLUTION

SESSION : 2001-2002

E 10

3

Please write on this side only, legibly and neatly, between the margins

QUESTION

SOLUTION

80

i) Choose $\underline{v} = \underline{i} P + \underline{j} Q$; $d\underline{v} = \underline{i} dx + \underline{j} dy \Rightarrow \underline{v} \cdot d\underline{r} = P dx + Q dy$


$\underline{i} \cdot \text{curl } \underline{v} = \underline{i} \cdot \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & 0 \end{vmatrix} = Q_x - P_y$. Hence result

ii) $\oint_C (y^2 dx + x^2 dy) = \int_{C_1} + \int_{C_2} + \int_{C_3}$

$$= \frac{1}{4} \int_{\frac{1}{2}}^1 \left(\frac{1}{4x^2} - 1 \right) dx$$

$$= -\frac{1}{4} \left[\frac{1}{4x} + x \right]_{\frac{1}{2}}^1$$

$$= -\frac{1}{4} \left\{ \frac{1}{4} + 1 - \frac{1}{2} - \frac{1}{8} \right\} = -\frac{1}{16}$$



2

$$\int_{C_2} (y^2 dx + x^2 dy) = \int_{\frac{1}{4}}^1 dy = 3/4$$

$$C_2: x=1, y: \frac{1}{4} \rightarrow 1$$

$$\int_{C_3} (y^2 dx + x^2 dy) = 2 \int_{\frac{1}{4}}^{\frac{1}{2}} x^2 dx = \frac{2}{3} [x^3]_{\frac{1}{4}}^{\frac{1}{2}} = -\frac{7}{8} \cdot \frac{2}{3} = -\frac{7}{12}$$

$$C_3: y=x \quad x: 1 \rightarrow \frac{1}{2}$$

$$\text{Total: } \oint_C (y^2 dx + x^2 dy) = -\frac{1}{16} + \frac{3}{4} - \frac{7}{12} = \frac{11}{16} - \frac{7}{12} = \frac{1}{4} \left(\frac{11}{4} - \frac{7}{3} \right) = 5/48$$

2

2

I have checked the answer by doing the area integral over R two different ways:

$$\text{LHS} = 2 \iint_R (x-y) dy dx = 2 \int_{\frac{1}{4}}^1 \left(\int_{\frac{1}{4x}}^x (x-y) dy \right) dx = 2 \int_{\frac{1}{4}}^1 \left[xy - \frac{y^2}{2} \right]_{\frac{1}{4x}}^x dx$$

$$= \int_{\frac{1}{4}}^1 \left(x^2 - \frac{1}{2} + \frac{1}{16x^2} \right) dx = \left[\frac{x^3}{3} - \frac{x}{2} - \frac{x^{-1}}{16} \right]_{\frac{1}{4}}^1 = 5/48$$

Setter: J. D. GIBBON

Setter's signature: J. D. Gibbon

Checker: A. WALTON

Checker's signature: Andrew Walton

EXAMINATION QUESTION / SOLUTION


SESSION : 2001-2002

E II

Please write on this side only, legibly and neatly, between the margins

QUESTION
1SOLUTION
33

(i) Venn diagram - as seen



2

$$\text{pr}(E_2 | E_3) = p_2$$

1

$$\text{pr}(E_1 \cap E_3) = p_{13}p_3$$

1

$$\text{pr}(E_3 | E_1) = p_{13}p_3/p_1$$

1

$$\text{pr}(E_1 | E_2) = 0$$

1

(ii) (a) prob= $\text{pr}(D_1) \times \text{pr}(D_3) = 0.001;$

1

$$(b) \text{prob}= \text{pr}(D_1 \cap D_2) = \text{pr}(D_1) + \text{pr}(D_2) - \text{pr}(D_1 \cup D_2)$$

2

$$= 0.1 + 0.05 - 0.12 = 0.03;$$

2

(c) $\text{pr}(D_1 \cap D_2 \cap D_3) = 0$ since D_2 and D_3 exclusive;

2

$$(d) \text{pr}(D_2 | D_1) = \text{pr}(D_2 \cap D_1)/\text{pr}(D_1) = 0.03/0.1 = 0.30;$$

2

(e) $\text{pr}(D_1 | D_3) = \text{pr}(D_1) = 0.1$, since independent

2

(15)

Setter : MJ CROWDER

Setter's signature : MJ Crowder

Checker : AT WALDEN

Checker's signature : AT Walden

MATHEMATICS FOR ENGINEERING STUDENTS
EXAMINATION QUESTION / SOLUTION

SESSION : 2001-2002

Please write on this side only, legibly and neatly, between the margins

PAPER
EE2 (3)

E 12

QUESTION
2

SOLUTION
34

(i) Expression: $E(X_1) = \sum_{x_1} x_1 p_1(x_1)$

Proof: $E(X_1 + X_2) = \sum_{x_1, x_2} (x_1 + x_2) p(x_1, x_2)$

$$= \sum_{x_1, x_2} x_1 p(x_1, x_2) + \sum_{x_1, x_2} x_2 p(x_1, x_2)$$

$$= \sum_{x_1} x_1 p_1(x_1) + \sum_{x_2} x_2 p_2(x_2) = E(X_1) + E(X_2)$$

(ii) $N(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$

$$N(2\mu_1 - 3\mu_2, 4\sigma_1^2 + 9\sigma_2^2)$$

(iii) mean = $2\mu_3 - 3\mu_4$

$$\text{variance} = 4\sigma_3^2 - 12\rho\sigma_3\sigma_4 + 9\sigma_4^2$$

(iv) prob = $\text{pr}(\text{Supply} \geq \text{Dem1} + \text{Dem2})$,

where Supply - (Dem1 + Dem2) is $N(1.14, 0.36)$,

$$\text{so prob} = 1 - \Phi(1.14/0.6) = 1 - \Phi(1.9) = 0.0287$$

1

3

1

2

1

3

2

2

15

Setter : M J CROWDER

Setter's signature: M J Crowder

Checker: A T WALDEN

Checker's signature: A T Walden