

IMPERIAL COLLEGE LONDON

DEPARTMENT OF ELECTRICAL AND ELECTRONIC ENGINEERING
EXAMINATIONS 2006

Corrected Copy

DISCRETE MATHEMATICS AND COMPUTATIONAL COMPLEXITY

Monday, 22 May 2:00 pm

Time allowed: 2:00 hours

There are FOUR questions on this paper.**Q1 is compulsory.****Answer Q1 and any two of questions 2-4.****Q1 carries 40% of the marks. Questions 2 to 4 carry equal marks (30% each).****Any special instructions for invigilators and information for candidates are on page 1.**

Examiners responsible	First Marker(s) :	G.A. Constantinides,
	Second Marker(s) :	T.J.W. Clarke,

NOTATION

The following notation is used throughout this paper:

\mathbb{R} : The set of real numbers.

\mathbb{Z} : The set of integers.

\mathbb{C} : The set of complex numbers.

\mathbb{N} : The set of natural numbers.

The Questions

1. [Compulsory]

- a) Give an example of a countable infinite set. [2]
- b) i) Draw the digraph of the relation R on the set $\{a, b, c\}$, where xRy iff x immediately precedes y in the alphabet.
ii) State whether this R is symmetric, transitive, and/or reflexive.
iii) List the elements of R^* for this R . [10]
- c) Consider the function $f : \mathbb{C} \rightarrow \mathbb{R}$ given by $f(x) = |x|$.
i) Is this function injective? Justify your answer.
ii) Is this function surjective? Justify your answer. [5]
- d) Let p be the proposition 'I am in an exam', and let q be the proposition 'I am not allowed to talk'. Consider the proposition $q \rightarrow p$. Is it true now? Would it be true if you were revising in the 'silent section' of the library? Briefly justify your answers. [5]
- e) Express the proposition 'there is an EE3 student who finds this exam easy' in symbolic logic, given the following predicates. $P(x)$ is the predicate 'x is an EE3 student', $Q(x)$ is the predicate 'x finds this exam easy'. You should take the set of students studying Discrete Mathematics and Computational Complexity as the universe of discourse. [4]
- f) "This exam is either difficult or I didn't revise properly. But I'm sure I revised properly, so this exam must be difficult". What is the name given to the rule of inference being applied here? [3]
- g) If $f(x)$ and $g(x)$ are both $O(x^2)$, use the results from the lectures to provide a big-O expression for (i) $f(x) + g(x)$ and (ii) $f(x)g(x)$. [4]
- h) Briefly define the term 'polynomial-time reduction'. [7]

2. This question relates to a function $f : A \rightarrow B$, where A and B are finite sets.
- a) Let R denote the range of the function. What relationship exists between R and B ? In the case where f is a surjection, what more can be deduced about this relationship? [2]
 - b) Prove that the cardinality of the range of f is at most the cardinality of its domain. *Hint:* you may assume the pigeonhole principle without proof. [10]
 - c) Given that f is a surjection, and that A and B are finite sets, what can be deduced about the cardinalities of A and B ? Prove this result. [3]
 - d) Prove that f has an inverse iff it is a bijection. [15]

3. a) Let P be the proposition $p \wedge (q \vee r) \vee \neg(p \vee (q \vee r))$. Replacing all occurrences of $(q \vee r)$ by $(q \wedge r)$ gives the proposition $P^* = p \wedge (q \wedge r) \vee \neg(p \vee (q \wedge r))$.

Determine whether each of the following compound propositions is a tautology, and provide a suitable proof of your answer in each case.

- i) $q \wedge r \rightarrow q \vee r$.
- ii) $P \rightarrow P^*$.
- iii) $P^* \rightarrow P$.

[14]

- b) You ask two lecturers, G and T, for help, but they try to confuse you. G says 'If T is telling the truth, then so am I'. T says 'at least one of us is lying'. Let p be the proposition 'G is telling the truth'. Let q be the proposition 'T is telling the truth'.

- i) Express G's statement using appropriate logical connectives.
- ii) Express T's statement using appropriate logical connectives.
- iii) By considering possibilities consistent with the truth values of p and q , and the statements made by G and T, deduce who, if anyone, is a liar. Fully explain your answer.

[16]

4. a) Define what is meant by the statements $f(x)$ is $O(g(x))$, $f(x)$ is $\Omega(g(x))$, and $f(x)$ is $\Theta(g(x))$, using appropriate symbolic logic. [3]
- b) Prove that $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$ is $\Theta(x^n)$ when $a_i \geq 0$ for $0 \leq i < n$ and $a_n > 0$. [9]
- c) Derive the number of each of the following type of operation performed by a call to the procedure `proc1(n)` of Figure 4.1, in terms of n :
- i) assignments (including for-loop initialization and incrementation assignments),
 - ii) multiplication,
 - iii) incrementation,
 - iv) comparison.
- [9]
- d) Given that the above operation types are responsible for the run-time of this code, and that each such operation takes $\Theta(1)$ time, derive a big-Theta expression of the form $\Theta(n^k)$ for the execution time of `proc1`. Hence derive a suitable big-O expression for the execution time of `proc2`, expressing your answer in terms of c , an integer constant. [9]

```

proc1(n)
{
  for i = 1 to n {
    t = 2*i
    for j = 1 to t
      a[i][j] = a[i][j]*2
    }
}

proc2(n)
{
  for i = 1 to c
    proc2( floor(n/2) )
  proc1(n)
}

```

Figure 4.1 Two procedures

1. a) \mathbb{Z} 

(ii) Not symmetric
Not transitive
Not reflexive

(iii) (a, b) (b, c) (a, c)

c) (i) $f(x) = f(y)$
 $|x| = |y|$

Consider $x = 1$, $y = -1$
Then $|x| = |y|$ but $x \neq y$, so not injective.

(ii) ~~Let $y = f(x)$~~

(ii) There is no complex # with a negative magnitude \Rightarrow
not surjective.

d) True now: $p \Rightarrow q$
False then: $p \Rightarrow q$

TRUE \Rightarrow TRUE \therefore TRUE
TRUE \Rightarrow FALSE \therefore FALSE
FALSE \Rightarrow TRUE \therefore FALSE

e) $\exists x (P(x) \wedge Q(x))$

f) The disjunctive syllogism

g) (i) $f(x) + g(x)$ is ~~$O(\max(|f(x)|, |g(x)|))$~~ $O(\max(x^2, x^2)) = O(x^2)$

(ii) $f(x)g(x)$ is $O(x^4)$

h) A polynomial time algorithm that transposes ~~over~~ a given instance of one decision problem into an instance of another, such that the answer to the 1st problem is "YES" iff the answer to the 2nd problem is also "YES".

2.

1) a) $R \subseteq B$.

when f is a surjection, $R = B$.

b) ~~$|A| \leq |B|$~~
 ~~$|A| > |B|$~~

b) We want to prove that $|f(A)| = |R| \leq |A|$.

Assume $|f(A)| > |A|$
let us define a function $g: f(A) \rightarrow A$ by
 $g(b) = a$ for some $a \in A$ such that $f(a) = b$,
so $f(g(b)) = b$.

g is an injection since $g(b) = g(c) \Rightarrow f(g(b)) = f(g(c)) \Rightarrow b = c$.

But by the pigeonhole principle on g , it cannot be an injection.

c) $|A| > |B|$

Since f is surjective $f(A) = B$ so $|f(A)| = |B|$.
However $|f(A)| \leq |A|$ for part (b).

$$|B| = |f(A)| \leq |A|$$

d)

First, prove that if $f: A \rightarrow B$ is a bijection, then it has an inverse f^{-1} .

$$\text{Consider } f^{-1} = \bigcup_{(a,b) \in f} \{(a,b)\} \subseteq B \times A$$

here f^{-1} is a relation from B to A .

As f is an injection, no more than one element of A for each element of B .

As f is a surjection, no less than one element of A for each element of B .

So f^{-1} is a function.

Next, prove that if $f: A \rightarrow B$ has inverse $f^{-1}: B \rightarrow A$, then f is a bijection.

If $f(a) = f(b)$ then $f^{-1}(f(a)) = f^{-1}(f(b)) \Rightarrow a = b$, so f is an injection.

Since for any $b \in B$ $a = f^{-1}(b) \in A$, we have $f(a) = f(f^{-1}(b)) = b$. f is a surjection.

3.

AA. a) (i) $q \wedge r \rightarrow q \vee r$

q	r	$q \wedge r$	$q \vee r$	$q \wedge r \rightarrow q \vee r$	F	F
F	F	F	F	T		
F	T	F	T	T		
T	F	F	T	T		
T	T	T	T	T		

✓ TAUTOLOGY

B (ii) $P \rightarrow P^*$, $P^* \rightarrow P$ — Neither are tautologies
& (iii)

P	q	r	P	P*	$P \rightarrow P^*$	$P^* \rightarrow P$
F	F	F	F	T	T	F
F	F	T	T	T	T	T
F	T	F	T	T	T	T
F	T	T	T	F	F	T
T	F	F	F	F	T	T
T	F	T	T	F	F	T
T	T	F	T	F	F	T
T	T	T	T	T	T	T

As a counter-example, consider P, q & r false for $P^* \rightarrow P$ and P false q & r true for $P \rightarrow P^*$.

b) (i) $q \rightarrow p$

(ii) $\neg p \vee \neg q$

(iii) Consider the case that q & T are both telling the truth.

(1) \rightarrow	p	q	$q \rightarrow p$	$\neg p \vee \neg q$
(2) \rightarrow	F	F	T	T
(3) \rightarrow	F	T	F	T (*)
(4) \rightarrow	T	F	T	T
	T	T	T	F

3/7
6/10

#36) (iii) [contd.]

Consider each possibility in the truth table.

- (1) Contradiction: G & T both lying but what they say is true.
- (2) No contradiction
- (3) Contradiction: T is lying but what he says is true.
- (4) Contradiction: T is telling the truth but what he says is false.

Only one possibility is consistent: G is a liar, while T tells the truth.

4. a) $f(x)$ is $O(g(x)) \equiv \exists c \in \mathbb{R}^+ \exists k \in \mathbb{R}^+ \forall x (x > k) \rightarrow (|f(x)| \leq c|g(x)|)$
 $f(x)$ is $\Omega(g(x)) \equiv \exists c \in \mathbb{R}^+ \exists k \in \mathbb{R}^+ \forall x (x > k) \rightarrow (|f(x)| \geq c|g(x)|)$
 $f(x)$ is $\Theta(g(x)) \equiv \exists c_1 \in \mathbb{R}^+ \exists c_2 \in \mathbb{R}^+ \exists k \in \mathbb{R}^+ \forall x (x > k) \rightarrow (c_1|g(x)| \leq |f(x)| \leq c_2|g(x)|)$

b) Need to prove

(i) $f(x)$ is $O(x^n)$

(ii) $f(x)$ is $\Omega(x^n)$

(i) $|f(x)| \leq |a_n x^n| + |a_{n-1} x^{n-1}| + \dots + |a_1 x| + |a_0|$
 $= |x^n| (|a_n| + |a_{n-1}|/|x| + \dots + |a_0|/|x^n|)$
 $\leq |x^n| (|a_n| + |a_{n-1}| + \dots + |a_0|)$ for $x > 1$

So with $c = |a_n| + \dots + |a_0| (> 0)$ and $k = 1$
 $f(x)$ is $O(x^n)$

(ii) $|f(x)| = |a_n x^n + \dots + a_0|$
 $\geq |a_n x^n|$ for $x \geq 0$
 $= |x^n| a_n$ since $a_n > 0$

So with $c = a_n$ and $k = 1$ (say)
 $f(x)$ is $\Omega(x^n)$

Thus $f(x)$ is $\Theta(x^n)$

5/7
8/10

4. b. c) We can derive a $\theta(\cdot)$ expression for all the operations - first count # ops

There are n iterations of the outer loop &
 $\sum_{i=1}^n 2i = 2 \sum_{i=1}^n i = n(n+1)$ iterations of the inner loop

(i) assignments of a[i][j] : $n(n+1)$
 assignments due to init of i : 1
 " " " " of j : n
 assignments due to increment of i : n
 assignments due to t : n
 TOTAL = $2n(n+1) + 3n + 1$
 $= \underline{2n(n+2) + 1} = n(2n+5) + 1$

(ii) mult of i : n
 mult of a : $n(n+1)$
 TOTAL = $n(n+2)$

(iii) increment of i : n
 " " of j : $n(n+1)$
 TOTAL = $3n(n+2)$

(iv) comparison of j : $n+1$
 j : $n(n+1) + \frac{1}{2}n = n(n+2)$
 TOTAL = $n(n+2) + n + 1$
 $= \underline{n(n+3) + 1}$

d) Total run time is $\Theta(n(2n+5))$

$\Theta(1)[n(2n+5) + 1] + \Theta(1)[n(n+2)]$
 $+ \Theta(1)[n(n+2)] + \Theta(1)[n(n+3) + 1]$
 $= \underline{\Theta(n^2)}$

6/7
9/10

4.18d) [contd]

Denote the execution time of $\text{proc}(n)$ by $g(n)$. Then

$$g(n) \leq c g(n/2) + d n^2$$

This is a D&C recurrence.

If $c < 2^2 = 4$ run time is $O(n^2)$

If $c = 2^2 = 4$ run time is $O(n^2 \log n)$

If $c > 2^2 = 4$ run time is ~~$O(n^2)$~~ $O(n^{\log_2 c})$



7/7
10/10

