

**M2PM3 COMPLEX ANALYSIS: PROBLEMS 3, 5.2.2009**

Q1 *The Gamma function (real case).*

Define

$$\Gamma(x) := \int_0^{\infty} t^{x-1} e^{-t} dt \quad (x > 0).$$

(i) Show that the integral converges for  $x > 0$ , but not for  $x \leq 0$ .

(ii) Show that

$$\Gamma(x+1) = x\Gamma(x) \quad (x > 0).$$

(iii) Show that

$$\Gamma(n+1) = n! \quad (n = 0, 1, 2, \dots).$$

Interpret  $\Gamma$  as providing a continuous extension of the factorial.

(iv) Show that

$$\Gamma(1/2) = \sqrt{\pi}.$$

Q2. *The Gamma function (complex case).*

For  $z \in \mathbf{C}$ , define

$$\Gamma(z) := \int_0^{\infty} t^{z-1} e^{-t} dt \quad (x = \Re z > 0).$$

(i) Show that the integral converges for  $\Re z > 0$ .

(ii) Show that

$$\Gamma(z+1) = z\Gamma(z) \quad (x = \Re z > 0).$$

(iii) Use (ii) to extend the domain of definition of  $\Gamma(z)$  from  $\Re z > 0$  to  $\Re z > -1$ .

(iv) By repeated use of (iii) (or by induction), extend the domain of definition of  $\Gamma(z)$  to  $\Re z > -n$  ( $n = 1, 2, 3, \dots$ ), and so to all of the complex plane  $\mathbf{C}$ . Show also that  $\Gamma$  so extended has complex values (i.e., in the complex plane  $\mathbf{C}$  rather than the extended complex plane  $\mathbf{C}^*$ ) except for  $z = 0, -1, -2, \dots$

(v) Find  $\Gamma(-2.5)$  and  $\Gamma(3.5)$ .

Q3 *The Riemann zeta function ( $\sigma > 1$ ).*

(i) For real  $\sigma$ , define (where convergent)

$$\zeta(\sigma) := \sum_{n=1}^{\infty} 1/n^{\sigma}.$$

Find the set of  $\sigma$  for which the series is convergent.

(ii) For  $s = \sigma + i\tau \in \mathbf{C}$ , define (where convergent)

$$\zeta(s) := \sum_{n=1}^{\infty} 1/n^s.$$

Find the half-plane of convergence of this Dirichlet series, and its half-plane of absolute convergence.

Q4 *The Riemann zeta function* ( $\sigma > 0$ ).

(i) By using the Alternating Series Test, or otherwise, show that

$$\sum_{n=1}^{\infty} (-1)^{n-1} / n^{\sigma}$$

converges for  $\sigma > 0$  but not for  $\sigma \leq 0$ . Find the half-planes of convergence and of absolute convergence of the Dirichlet series

$$\sum_{n=1}^{\infty} (-1)^{n-1} / n^s.$$

(ii) By considering the sums over  $n$  odd and even separately, or otherwise, show that

$$\sum_{n=1}^{\infty} (-1)^{n-1} / n^s = (1 - 2^{1-s}) \zeta(s) \quad (\sigma = \Re s > 1).$$

(iii) Deduce that defining

$$\zeta(s) := \frac{1}{(1 - 2^{1-s})} \sum_{n=1}^{\infty} (-1)^{n-1} / n^s$$

defines a function  $\zeta$  for  $\sigma = \Re s > 0$ ,  $s \neq 1$ .

(iv) Show that as  $s \rightarrow 1$ ,

$$\zeta(s) \sim \frac{1}{s-1}.$$

(You may assume that  $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots = \log 2$ . Hint: use L'Hospital's Rule.)

NHB