

M2PM3 SOLUTIONS 4. 19.2.2009

Q1. *Triangle Lemma.*

Draw the line joining z_1 and z_2 , and produce it until it meets triangle Δ at points Z_1, Z_2 say. Then

$$|z_1 - z_2| \leq |Z_1 - Z_2|,$$

with equality iff both z_1, z_2 are *on* Δ rather than inside it (so $z_1 = Z_1, z_2 = Z_2$). There are two cases.

(i) Z_1, Z_2 lie on different sides of the triangle. Let Z_3 be the vertex in which these sides meet. Then by the Triangle Inequality,

$$|Z_1 - Z_2| \leq |Z_1 - Z_3| + |Z_2 - Z_3| \leq L_1 + L_2 \leq L,$$

where L_1, L_2 are the lengths of the sides containing Z_1, Z_2 . Combining,

$$|z_1 - z_2| \leq L.$$

(ii) Z_1, Z_2 lie on the same side, of length L_{12} say. Then

$$|Z_1 - Z_2| \leq L_{12} \leq L,$$

and the result follows as in (i).

Q2. *Holomorphy and Conjugation.*

As f is holomorphic at z_0 , $g(z) := f(z_0 + z)$ is holomorphic at the origin. So

$$\frac{g(h) - g(0)}{h} \rightarrow g'(0) = f'(z_0) \quad (h = k + i\ell \rightarrow 0).$$

So replacing ℓ by $-\ell$,

$$\frac{g(\bar{h}) - g(0)}{\bar{h}} \rightarrow g'(0) = f'(z_0) \quad (h \rightarrow 0).$$

So

$$\frac{g(\bar{h}) - g(0)}{h} = \frac{g(\bar{h}) - g(0)}{\bar{h}} \cdot \frac{\bar{h}}{h} \sim g'(0) \cdot \bar{h}/h \quad (h \rightarrow 0).$$

But $\bar{h}/h = (k - i\ell)/(k + i\ell)$. For h real ($\ell = 0$) this is 1; for h imaginary ($k = 0$) this is -1. So this does not have a limit as $h \rightarrow 0$ (recall that this means $|h| \rightarrow 0$: NO restriction on the argument of h). So $g(h)$ is not differentiable w.r.t. \bar{h} at 0. So $f(z)$ is not differentiable w.r.t. \bar{z} .

Q3. *Union of Domains.*

Suppose $\bigcup_i D_i = G \cup H$ with G, H disjoint and open. We have to show one of G, H is empty. Now

$$D_j = (D_j \cap G) \cup (D_j \cap H).$$

The union on the RHS is disjoint (as G, H are), and the sets on RHS are open. As D_j is connected, one of these sets must be empty: say, $D_j \cap H$ is empty,

i.e. $D_j \subset G$. Similarly, each D_k is contained in one of G, H . But if $D_k \subset H$, $D_j \cap D_k$ non-empty contradicts $G \cap H$ empty. So all the $D_k \subset G$. So H is empty.

Alternative Proof [assuming equivalence of connectedness and polygonal, or arcwise, connectedness].

Take $z_0 \in \bigcap_i D_i$. We can join any point in any D_j to z_0 by a path [e.g., a polygonal arc] lying in D_j , so in $\bigcup_i D_i$. So we can join any two points in $\bigcup_i D_i$ by such a path, by joining the paths linking each to z_0 . So $\bigcup_i D_i$ is polygonally connected, so connected.

Q4 *Connected Components.*

Let z be any point in S . Let $\{C_i\}$ be the class of all connected subsets of S containing z . This class is non-empty (as $\{z\}$ is connected). By Q3, $C := \bigcup_i C_i$ is a connected subset of S containing z . By construction (as the union of all ...), C is maximal. So C is a component, and contains z . If C' is another component containing z , C' must be one of the C_i , so $C' \subset C$. But C' is maximal (it is a component). So $C \subset C'$, so $C = C'$. So each point z lies in a unique (connected) component, called the (*connected*) *component containing* z .

Q5. Write a_n for the coefficient of z^n .

(i) $a_{n+1}/a_n = -n/(n+1) = -1/(1+1/n) \rightarrow -1$, so $|a_{n+1}/a_n| \rightarrow 1$. So by the Ratio Test, the radius of convergence is 1. So the function is holomorphic in the unit disc $D := \{z : |z| < 1\}$.

Note. The sum function is $\log(1+z)$. This has a singularity at $z = -1$ (a branch point).

(ii) $a_{5n} = 1$, $a_{5n+k} = 0$ ($k = 1, 2, 3, 4$). $\limsup |a_n|^{1/n} = 1$. So the radius of convergence is 1, and the region of holomorphy is again D .

Note. The sum function is $1/(1-z^5)$. This has 5 singularities on the unit circle, at the 5 fifth roots of unity.

(iii) $a_n = 1/n^n$, $a_n^{1/n} = 1/n \rightarrow 0$. So the radius of convergence is infinite. The sum function is holomorphic throughout the complex plane \mathbf{C} (is an *entire* function, or an *integral* function).

NHB