

SECTION A

1. Write down the Taylor series about x = 2 for the function

$$f(x) = x^{-1}.$$

State whether this Taylor series converges to f(x) for:

a)
$$x = 1$$
, b) $x = 4$. [5 marks]

2. Find the solutions of the following differential equations:

(i)
$$e^{y} \frac{dy}{dx} = x$$
 with $y(1) = 0$,
(ii) $x \frac{dy}{dx} + 2y = x$ with $y(1) = 0$.

[8 marks]

3. Solve the differential equation

$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 3y = 0$$

with the initial conditions y(0) = 2, y'(0) = 1.

[5 marks]

4. Show, by taking limits along two different paths to the origin (0,0), that

$$\lim_{(x,y)\to(0,0)} \frac{xy}{x^2 + xy + y^2}$$

does not exist.

[4 marks]

CONTINUED



5. Work out all first and second partial derivatives of

$$f(x,y) = x^4 - 6x^2y^2 + y^4,$$

and verify that

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x},$$
$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0$$

[5 marks]

6. Suppose that x = x(t), y = y(t), and z = z(t) are functions of t such that

$$x(0) = 2, y(0) = -1, z(0) = 0.$$

Suppose that the derivatives satisfy

$$x'(0) = y'(0) = 1, \ z'(0) = -1.$$

Then work out

$$\frac{dF}{dt}(0)$$

where F(t) = f(x(t), y(t), z(t)), and

$$f(x, y, z) = x^2 y + \sin(xyz).$$

[5 marks]

7. Find the gradient vector $\nabla f(1, 1, 1)$, where

$$f(x, y, z) = \frac{xyz}{x^2 + y^2 + z^2}.$$

Find also the derivative of f in the direction $\mathbf{i} - 2\mathbf{j} - 2\mathbf{k}$ at the point (1, 1, 1). [5 marks]

8. Locate and classify all stationary points of the function

$$f(x,y) = x^2y - 2xy + y^2 - 15y.$$

[8 marks]

Paper Code MATH102 Page 3 of 5

CONTINUED



9. Find the linear approximation near (x, y) = (2, 1) to the function

$$f(x,y) = \frac{1}{x^2 - y^2}.$$

[4 marks]

10. By changing the order of integration, compute

$$\int_0^1 \int_y^1 e^{y/x} dx dy.$$

[6 marks]

SECTION B

11.

In this question, let

$$f(y) = \frac{1}{1+y}, \quad g(x) = \frac{1}{1+x^2}, \quad h(x) = \tan^{-1}(x).$$

- (i) Write down the Taylor series of f, g and h, all at 0.
- (ii) Show that if $y \ge 0$, the *n*th remainder term $R_n(y,0)$ of f at 0 satisfies

$$|R_n(y,0)| \le y^{n+1}.$$

Hence show that if $x \ge 0$,

$$\int_0^x R_n(t^2, 0) dt \le \frac{x^{2n+3}}{2n+3}.$$

(iii) Express $h(1) = \tan^{-1}(1)$ in terms of π . If $P_n(x,0)$ denotes the *n*th Taylor polynomial of h at 0, use your calculator to compute $P_{22}(1,0)$ and $\tan^{-1}(1)$ and verify that

$$\left|\tan^{-1}(1) - P_{22}(1,0)\right| \le \frac{1}{23}.$$

[15 marks]

Paper Code MATH102 Page 4 of 5

CONTINUED



12. Solve the following differential equations with the given boundary conditions:

(i)

$$y'' - 4y' - 5y = 4e^x$$

with y(0) = 1, y'(0) = -1.(ii)

$$y'' - 4y' - 5y = -5x^2 + 2x + 5$$

with y(0) = 1, y'(0) = -1.

[15 marks]

13. Find the maximum and minimum values of the function f(x, y) in the region where $g(x, y) \leq 3$, where f(x, y) and g(x, y) are defined by

$$f(x, y) = xy + x,$$

$$g(x, y) = 3x^{2} + y^{2}.$$

[15 marks]

14.

a) Find the weight of the triangle R bounded by y = x, y = 2x and x = 1, where the density function is $\rho(x, y) = x$

b) Find the centre of mass $(\overline{x}, \overline{y})$ of R.

[15 marks]

END